

Contents

| | |
|--|--------------|
| Title page | <i>i</i> |
| Abstract | <i>ii</i> |
| Declaration | <i>xvi</i> |
| Certificate | <i>xvii</i> |
| Acknowledgements | <i>xix</i> |
| Contents | <i>xix</i> |
| List of tables | <i>xxix</i> |
| List of figures | <i>xxxii</i> |
| List of abbreviations | <i>xxxv</i> |
| List of symbols | <i>xxxvi</i> |
| | |
| Chapter 1: INTRODUCTION | [1-12] |
| 1.1. Polyphenols in fruits and vegetables | 3 |
| 1.2. Effect of processing on polyphenols and antioxidant properties | 4 |
| 1.3. Utilisation of fruit by-products | 5 |
| 1.4. Fruits and vegetables of Assam, India | 6 |
| 1.5. Objectives | 8 |
| Bibliography | 8 |
| | |
| Chapter 2: REVIEW OF LITERATURE | [13-45] |
| 2.1. Polyphenols as important phytochemicals in plants | 13 |
| 2.1.2. Phenolic acids | 14 |
| 2.1.2. Flavonoids | 14 |
| 2.1.3. Biosynthesis of polyphenol | 15 |
| 2.1.4. Functional role of polyphenols (phenolic acids and flavonoids) | 18 |
| 2.2. Polyphenols in fruits and vegetables | 20 |
| 2.3. Polyphenol bioavailability and metabolism | 22 |
| | |
| 2.4. Effect of processing of fruits and vegetables on the polyphenol content | 23 |
| 2.5. Processing effects on the polyphenol content of fruits and | 24 |

| | | |
|---|---|---------|
| | vegetables | |
| 2.6. | Fruit by-products, their phytochemical content and utilization | 26 |
| | Bibliography | 30 |
| Chapter 3: TO DETERMINE THE PHYTOCHEMICAL CONTENT AND ANTIOXIDANT CAPACITIES IN SELECTED FRESH FRUITS AND COOKED VEGETABLES OF ASSAM. | | [46-84] |
| 3.1. | Introduction | 46 |
| 3.2. | Material and methods | 48 |
| 3.2.1. | Materials | 48 |
| 3.2.2. | Fruit samples | 48 |
| 3.2.3. | Vegetable samples | 48 |
| 3.2.4. | Cooking treatments | 53 |
| 3.2.5. | Sample extraction | 53 |
| 3.2.6. | Phytochemical content and antioxidant activities | 53 |
| 3.2.6.1. | Determination of total phenolic content | 53 |
| 3.2.6.2. | Determination of total flavonoid content | 54 |
| 3.2.6.3. | Determination of ferric reducing antioxidant property (FRAP) | 54 |
| 3.2.6.4. | Determination of DPPH radical scavenging activity | 54 |
| 3.2.6.5. | Determination of metal chelating capacity | 56 |
| 3.2.7. | RP-HPLC study of the polyphenols | 56 |
| 3.2.8. | Statistical analysis | 57 |
| 3.3. | Results and discussion | 57 |
| 3.3.1. | Phytochemical and antioxidant properties of thirteen fruit samples | 57 |
| 3.3.2. | RP-HPLC analysis of the phenolic acids in the thirteen fruit samples | 59 |
| 3.3.3. | Changes in the phytochemical content after processing in the vegetable samples | 65 |
| 3.3.3.1. | Changes in total phenolic content (TPC) and total flavonoid content (TFC) | 65 |
| 3.3.4. | Changes in the antioxidant activities after processing in the vegetable samples | 68 |
| 3.3.4.1. | Changes in ferric reducing antioxidant potential (FRAP) | 68 |
| 3.3.4.2. | Changes in DPPH radical scavenging activity | 69 |
| 3.3.4.3. | Changes in metal chelation capacity (MCC) | 71 |

| | | |
|---|--|----------|
| 3.3.5. | Changes in the phenolic acids composition in the processed four selected vegetables | 72 |
| 3.4. | Conclusion | 79 |
| | Bibliography | 80 |
| Chapter 4: EFFECT OF SPRAY DRYING OF FOUR FRUIT JUICES ON PHYSICOCHEMICAL, PHYTOCHEMICAL AND ANTIOXIDANT PROPERTIES | | [85-104] |
| 4.1. | Introduction | 85 |
| 4.2. | Materials and methods | 86 |
| 4.2.1. | Materials | 86 |
| 4.2.2. | Fruit juice preparation | 86 |
| 4.2.3. | pH, total solid content and viscosity of the feed samples | 87 |
| 4.2.4. | Spray drying of the juice-maltodextrin mixture | 87 |
| 4.2.5. | Yield of powder | 87 |
| 4.2.6. | Physicochemical properties of the spray dried powder | 88 |
| 4.2.6.1. | Moisture content | 88 |
| 4.2.6.2. | Bulk density, tapped density, Hausner's ratio and Carr index | 88 |
| 4.2.6.3. | Colour properties of the fresh feed and reconstituted samples | 89 |
| 4.2.6.4. | pH and titratable acidity of the powdered samples | 89 |
| 4.2.6.5. | Solubility | 90 |
| 4.2.6.6. | Hygroscopicity | 90 |
| 4.2.7. | Surface morphology study of the spray dried powder by scanning electron microscopy (SEM) | 90 |
| 4.2.8. | Particle size distribution of the powdered samples | 90 |
| 4.2.9. | Phytochemical content and antioxidant activities of the fresh juice and dried powder | 91 |
| 4.2.9.1. | Sample extraction | 91 |
| 4.2.9.2. | Determination of total phenolic content | 91 |
| 4.2.9.3. | Determination of total flavonoid content | 91 |
| 4.2.9.4. | Determination of ferric reducing antioxidant property (FRAP) | 92 |
| 4.2.9.5. | Determination of DPPH activity | 92 |
| 4.2.10. | Statistical analysis | 92 |
| 4.3. | Results and discussion | 93 |
| 4.3.1. | Total solid content, pH and viscosity of the feed sample | 93 |

| | | |
|--|---|-----------|
| 4.3.2. | Particle size distribution, cohesiveness and flow ability of the powdered sample | 94 |
| 4.3.3. | Physicochemical parameters of the spray dried fruit juice powders | 96 |
| 4.3.4. | Surface morphology study of the spray dried juice powder | 97 |
| 4.3.5. | Colour of the spray dried fruit juice powders | 97 |
| 4.3.6. | Phytochemical content and activity of spray dried fruit juice | 99 |
| 4. 4. | Conclusion | 101 |
| | Bibliography | 101 |
| Chapter 5: EFFECT OF DIFFERENT PASTEURISATION METHODS ON PHYTOCHEMICAL AND ANTIOXIDANT PROPERTY OF FIVE FRUIT JUICES | | [105-131] |
| 5.1 | Introduction | 101 |
| 5.2 | Materials and methods | 106 |
| 5.2.1. | Materials | 106 |
| 5.2.2. | Fruit juice preparation | 107 |
| 5.2.3. | Total plate count of the treated samples | 107 |
| 5.2.4. | Colour | 107 |
| 5.2.5. | Changes in the phytochemical and antioxidant activity of the processed juice samples | 108 |
| 5.2.5.1. | Determination of total phenolic content | 108 |
| 5.2.5.2. | Determination of total flavonoid content | 108 |
| 5.2.5.3. | Determination of ferric reducing antioxidant property (FRAP) | 109 |
| 5.2.5.4. | Determination of DPPH radical scavenging activity | 109 |
| 5.2.6. | HPLC study of the polyphenols | 109 |
| 5.2.7. | Statistical analysis | 110 |
| 5.3. | Results and discussion | 110 |
| 5.3.1. | Microbial load in the processed juice samples | 110 |
| 5.3.2. | Colour values of the juice samples and overall change in colour (ΔE) after processing | 111 |
| 5.3.3. | Phytochemical and antioxidant changes | 113 |
| 5.3.4. | HPLC determination of the phenolic acids and ascorbic acid content in the processed juice samples | 119 |
| 5.4. | Conclusions | 127 |
| | Bibliography | 127 |

| | |
|--|-----------|
| Chapter6: OPTIMISATION OF PHENOLIC EXTRACTION FROM CARAMBOLA POMACE BY RESPONSE SURFACE METHODOLOGY AND ITS MICROENCAPSULATION BY SPRAY AND FREEZE DRYING METHODS | [132-155] |
| 6.1. Introduction | 132 |
| 6.2. Materials and methods | 134 |
| 6.2.1. Chemicals and reagents | 134 |
| 6.2.2. Materials | 134 |
| 6.2.3. Extraction of polyphenols from the pomace | 134 |
| 6.2.4. Experimental design for optimisation using Central composite rotatable design (CCRD) by response surface methodology (RSM) | 134 |
| 6.2.4.1. Total phenolic content (TPC) | 135 |
| 6.2.4.2. Ferric reducing antioxidant property (FRAP) | 135 |
| 6.2.4.3. DPPH radical scavenging activity | 135 |
| 6.2.5. RP-HPLC study of the polyphenol extract | 136 |
| 6.2.6. Encapsulation process | 137 |
| 6.2.6.1. Moisture content | 137 |
| 6.2.6.2. Bulk density | 138 |
| 6.2.6.3. Solubility | 138 |
| 6.2.6.4. Colour | 138 |
| 6.2.6.5. Hygroscopicity | 138 |
| 6.2.7. Total phenolic content analysis (TPC), surface phenolic content (SPC) and encapsulating efficiency | 139 |
| 6.2.8. <i>In vitro</i> simulated gastric and intestinal digestion release study of the encapsulates | 139 |
| 6.2.9. Surface morphology study by scanning electron microscope (SEM) | 140 |
| 6.2.10. Statistical analysis | 140 |
| 6.3. Results and discussion | 140 |
| 6.3.1. Optimisation and fitting of the model | 140 |
| 6.3.1.1. Effect of temperature and ethanol concentration on TPC | 140 |
| 6.3.1.2. Effect of temperature and ethanol concentration on FRAP | 141 |
| 6.3.1.3. Effect of temperature and ethanol concentration on DPPH radical scavenging activity | 143 |

| | | |
|--|---|-----------|
| 6.3.1.4. | Verification of the predictive model | 144 |
| 6.3.2. | RP-HPLC study of the crude polyphenols extract | 144 |
| 6.3.3. | Microencapsulation of the polyphenols extract | 145 |
| 6.3.3.1. | Physical properties of the encapsulates | 145 |
| 6.3.3.2. | Encapsulating efficiency, TPC and SPC | 147 |
| 6.3.3.3. | <i>In vitro</i> gastric and intestinal release of the microencapsulate under simulated conditions | 148 |
| 6.3.3.4. | Surface morphology study by Scanning electron microscope (SEM) | 149 |
| 6.4. | Conclusion | 151 |
| | Bibliography | 151 |
| Chapter 7: <i>IN VITRO</i> PHYSICOCHEMICAL, PHYTOCHEMICAL AND FUNCTIONAL PROPERTIES OF FIBRE RICH FRACTIONS DERIVED FROM BY-PRODUCTS OF SIX DIFFERENT FRUITS | | [156-176] |
| 7.1. | Introduction | 156 |
| 7.2. | Materials and methods | 157 |
| 7.2.1. | Chemicals and reagents | 157 |
| 7.2.2. | Proximate composition | 157 |
| 7.2.2.1. | Moisture content | 157 |
| 7.2.2.2. | Crude protein content | 158 |
| 7.2.2.3. | Crude lipid content | 158 |
| 7.2.2.4. | Ash content | 158 |
| 7.2.3. | Preparation of the Fibre Rich Fraction | 159 |
| 7.2.4. | Colour and dietary fibre content | 159 |
| 7.2.4.1. | Colour determination | 159 |
| 7.2.4.2. | Determination of the total dietary fibre (TDF), insoluble dietary fibre (IDF) and soluble dietary fibre (SDF) content of the prepared fibre samples | 159 |
| 7.2.5. | Physicochemical Properties | 161 |
| 7.2.5.1. | Moisture content | 161 |
| 7.2.5.2. | Bulk density | 161 |
| 7.2.5.3. | Water holding capacity | 161 |
| 7.2.5.4. | Swelling capacity | 161 |
| 7.2.5.5. | <i>Oil holding capacity</i> | 162 |

| | | |
|---|---|-----------|
| 7.2.6. | Phytochemical content and antioxidant activity | 162 |
| 7.2.6.1. | Sample extraction | 162 |
| 7.2.6.2. | Determination of total phenolic content | 162 |
| 7.2.6.3. | Determination of total flavonoid content | 162 |
| 7.2.6.4. | Determination of ferric reducing antioxidant property (FRAP) | 163 |
| 7.2.6.5. | Determination of DPPH Activity | 163 |
| 7.2.6.6. | Determination of Metal Chelation Capacity | 163 |
| 7.2.6.7. | RP-HPLC study of the polyphenols | 164 |
| 7.2.7. | Functional properties | 164 |
| 7.2.7.1. | Glucose adsorption capacity and amylase activity inhibition rate | 164 |
| 7.2.7.2. | Glucose diffusion rate and glucose diffusion reduction index (GDRI) | 165 |
| 7.2.8. | Statistical Analysis | 165 |
| 7.3. | Results and discussion | 165 |
| 7.3.1. | Proximate composition | 165 |
| 7.3.2. | Yield and total dietary fibre | 166 |
| 7.3.3. | Colour | 167 |
| 7.3.4. | Physicochemical properties | 167 |
| 7.3.5. | Phytochemical content and antioxidant activity | 168 |
| 7.3.5.1. | Phenolic acids compositions of the selected fibres determined by RP-HPLC | 169 |
| 7.3.6. | Functional properties | 172 |
| 7.3.6.1. | Glucose adsorption and amylase activity inhibition rate | 172 |
| 7.3.6.2. | Glucose diffusion rate (GDR) and glucose diffusion reduction index (GDRI) | 173 |
| 7.4. | Conclusion | 174 |
| | Bibliography | 174 |
| Chapter 8: DEVELOPMENT OF A CARAMBOLA POMACE FIBRE FORTIFIED MIX FRUIT BEVERAGE POWDER AND ITS CHARACTERIZATION | | [177-199] |
| 8.1. | Introduction | 177 |
| 8.2. | Materials and methods | 178 |

| | | |
|------------|--|-----|
| 8.2.1. | Materials | 179 |
| 8.2.2. | Fruit juice mix preparation for spray drying | 179 |
| 8.2.3. | Viscosity and pH of the mix fruit juice fibre-maltodextrin feed sample at 10°B | 179 |
| 8.2.4. | Spray drying of the mix fruit juice fibre-maltodextrin | 179 |
| 8.2.5. | Experimental design for optimization using central composite rotatable design (CCRD) by response surface methodology (RSM) | 179 |
| 8.2.6. | Determination of the response variables | 180 |
| 8.2.6.1. | Moisture content | 180 |
| 8.2.6.2. | Yield of powder | 180 |
| 8.2.6.3. | Solubility | 181 |
| 8.2.6.4. | Bulk density | 181 |
| 8.2.6.5. | Hygroscopicity | 181 |
| 8.2.7. | Characterization of the obtained spray dried fibre fortified mix fruit beverage powder | 181 |
| 8.2.7.1. | Determination of proximate content. | 181 |
| 8.2.7.1.1. | Moisture content | 181 |
| 8.2.7.1.2. | Crude protein content | 181 |
| 8.2.7.1.3. | Crude lipid content | 182 |
| 8.2.7.1.4. | Ash content | 182 |
| 8.2.7.1.5. | Total reducing sugar | 183 |
| 8.2.7.1.6. | Total dietary fibre | 183 |
| 8.2.7.2. | Color comparison of the reconstituted beverage with the feed solution before drying | 184 |
| 8.2.7.3. | Particle size distribution | 185 |
| 8.2.7.4. | Bulk density (BD), tapped density (TD), Hausner's ratio (HR) and Carr Index (CI) | 185 |
| 8.2.7.5. | Dissolution test and pH of the beverage powder | 186 |
| 8.2.7.6. | Water activity | 186 |
| 8.2.7.7. | Surface morphology of the spray dried powder by scanning electron microscopy (SEM) | 186 |
| 8.2.7.8. | Sensory analysis using 9-point Hedonic scale | 186 |
| 8.2.7.9. | Statistical analysis | 187 |
| 8.3. | Results and discussion | 187 |

| | | |
|---|--|-----------|
| 8.3.1. | Optimization and fitting of the model | 187 |
| 8.3.2. | Effect of independent variables on responses in the spray dried beverage powder | 187 |
| 8.3.2.1. | Moisture content | 187 |
| 8.3.2.2. | Yield | 188 |
| 8.3.2.3. | Solubility | 188 |
| 8.3.2.4. | Bulk density | 190 |
| 8.3.2.5. | Hygroscopicity | 190 |
| 8.3.3. | Verification of the predictive model | 192 |
| 8.3.4. | pH, total solid content and viscosity of the feed sample | 192 |
| 8.3.5. | Proximate composition of the beverage powder and dissolution test, pH and water activity (a_w) of the powder | 192 |
| 8.3.6. | Bulk density, tapped density and flow property of the powder | 193 |
| 8.3.7. | Change in color <i>L a b</i> values when the reconstituted beverage was compared to the fresh feed sample | 193 |
| 8.3.8. | Particle size distribution and Span | 194 |
| 8.3.9. | Surface morphology of the beverage powder | 194 |
| 8.3.10. | Sensory analysis results of the reconstituted beverage powder | 195 |
| 8.4. | Conclusion | 195 |
| | Bibliography | 196 |
| Chapter 9: EFFECT OF THE CARAMBOLA POMACE FORTIFIED MIX FRUIT BEVERAGE POWDER ON THE SERUM CHOLESTEROL AND GLUCOSE LEVELS AND ON THE FUNCTIONS OF TWO LIVER ENZYMES IN SPRAGUE DAWLEY RAT | | [200-214] |
| 9.1. | Introduction | 200 |
| 9.2. | Materials and Method | 201 |
| 9.2.1. | Materials | 202 |
| 9.2.2. | Test diet | |
| 9.2.3. | Animal experiment conditions | 202 |
| 9.2.4. | Dosage determination | 203 |
| 9.2.5. | Induction of high cholesterol condition and administration of test diet | 203 |
| 9.2.6. | Determination of the serum biochemistry | 204 |

| | |
|--|-----------|
| 9.2.7. Changes in body weight of the two test group | 204 |
| 9.2.8. Histopathology of the heart and aorta tissue | 205 |
| 9.2.9. Identification of short chain fatty acids in the caecal matter of the two test group by GCMS | 205 |
| 9.2.10. Statistical analysis | 206 |
| 9.3. Results and discussion | 206 |
| 9.3.1. Serum biochemistry | 206 |
| 9.3.2. Changes in body weight | 208 |
| 9.3.3. Histopathology | 208 |
| 9.3.4. SCFA by GCMS of the caecal matter | 208 |
| 9.4. Conclusions | 212 |
| Bibliography | 212 |
| | |
| Chapter 10: CONCLUSION AND FUTURE SCOPE OF WORK | [215-223] |
| 10.1. Conclusion | 215 |
| 10.1.1. Chapter 3 | 215 |
| 10.1.2. Chapter 4 | 217 |
| 10.1.3. Chapter 5 | 217 |
| 10.1.4. Chapter 6 | 219 |
| 10.1.5. Chapter 7 | 219 |
| 10.1.6. Chapter 8 | 220 |
| 10.1.7. Chapter 9 | 221 |
| 10.2. Future scope of work | 222 |
| List of publications | 223 |