

LIST OF TABLES

Chapter	Table no.	Table caption	Page no.
1	1.1	Physico-chemical properties of 16 USEPA priority PAHs.	6
2	2.1	Soil PAHs concentrations (ng g^{-1}) reported from elsewhere.	11
	2.2	Concentration ranges of PAHs (ng ml^{-1}) reported from different groundwater sources worldwide.	12
	2.3	Carcinogenic classification of 16 USEPA priority PAHs as per IARC, DHHS and US-EPA.	15
	2.4	Values of PAHs diagnostic ratios for a particular source category.	18-19
3	3.1(a)	Description of sampling sites around Borholla oilfield.	4
	3.1(b)	Description of sampling sites around Nambar oilfield.	5
	3.2	Retention time, Maximum % RSD, LOD value and External Standard Concentration of 16 PAH compounds.	11
	3.3	Recovery ranges of PAHs for water and soil samples.	17
	3.4	LOD values for heavy metals.	18
4	4.1(a)	Mean PAHs concentrations (ng g^{-1}) with minimum, maximum, median and geometric mean in soils around Borholla oilfield during pre-monsoon, monsoon and post-monsoon seasons of 2011.	2
	4.1(b)	Mean PAHs concentrations with minimum, maximum, median and geometric mean (ng g^{-1}), in soils around Borholla oilfield during pre-monsoon, monsoon and post-monsoon seasons of 2013.	3
	4.2	Mean PAHs concentrations with minimum, maximum, median and geometric mean (ng g^{-1}) in soils around Nambar oilfield during pre-monsoon, monsoon and post-monsoon seasons.	7
	4.3 (a)	Site-wise mean concentrations of PAHs (ng g^{-1}) with minimum, maximum, median and geometric mean values in soils around Borholla oilfield during pre-monsoon, monsoon and post-monsoon seasons of 2011.	15
	4.3(b)	Site-wise mean concentrations of PAHs (ng g^{-1}) with minimum, maximum, median and geometric mean values in soils around Borholla oilfield during pre-monsoon, monsoon and post-monsoon seasons of 2013.	16

Chapter	Table no.	Table caption	Page no.
4	4.4	Site-wise mean concentrations of PAHs with minimum, maximum, median and geometric mean (ng g^{-1}) in soils around Nambar oilfield during pre-monsoon, monsoon and post-monsoon seasons.	19
	4.5	Soil PAHs concentrations reported from elsewhere.	20
	4.6	Percentage contributions of two-, three-, four-, five- and six- ring PAHs around Borholla oilfield.	23
	4.7	Percentage contributions of two-, three-, four-, five- and six- ring PAHs around Nambar oilfield.	24
	4.8	Mean BaPeq concentrations (ng g^{-1}) in soils around Borholla oilfield during pre-monsoon, monsoon and post-monsoon seasons in the study period.	28
	4.9	Site-wise mean BaPeq concentrations (ng g^{-1}) of PAHs around Borholla oilfield during the study period.	29
	4.10	Mean BaPeq concentrations (ng g^{-1}) in soils around Nambar oilfield during pre-monsoon, monsoon and post-monsoon seasons during the study period.	30
	4.11	Site-wise mean BaPeq concentrations (ng g^{-1}) of PAHs around Nambar oilfield.	31
	4.12	Seasonal mean diagnostic ratios of PAHs along with their ranges in soils around Borholla oilfield.	33
	4.13	Seasonal mean diagnostic ratios of PAHs along with their ranges in soils around Nambar oilfield.	36
	4.14	PCA–MLR of PAHs: identification of sources and their respective contribution to soil PAHs around Borholla oilfield. The significant factor loadings are in bold.	41
	4.15	PCA–MLR of PAHs: identification of sources and their respective contribution to soil PAHs around Nambar oilfield. The significant factor loadings are in bold.	43
	4.16	Concentrations (mg g^{-1}) of organic carbon (OC) and black carbon (BC) in soils around Borholla and Nambar oilfields.	44
	4.17	Correlations of PAHs with organic carbon (OC) and black carbon (BC) during different seasons in soils	45

	around Borholla oilfield.	
4.18	Correlation of PAHs with organic carbon (OC) and black carbon (BC) during 3 sampling seasons in soil around Nambar oilfield.	46
4.19(a)	Minimum, maximum, mean, median and geometric mean concentrations (mg kg^{-1}) of metals in soils around Borholla oilfield during three seasons of 2011.	53
4.19(b)	Minimum, maximum, mean median and geometric mean concentrations (mg kg^{-1}) of metals in soils around Borholla oilfield during three seasons of 2013.	53
4.20	Minimum, maximum, mean, median and geometric mean concentrations (mg kg^{-1}) of metals in soils around Nambar oilfield during the sampling period.	55
4.21	Comparative account of heavy metal concentrations (mg kg^{-1}) of the present study with studies from elsewhere.	57
4.22	Pollution indices and corresponding degrees of contamination levels. Pollution Index (PI), Ecological risk factor (Er^i) and Risk Index (RI).	58
4.23	Pollution Index (PI) and Nemerow Integrated Pollution Index (NIPI) in soils around Borholla oilfield.	59
4.24	Potential ecological risk index of a single element (Eri) and potential ecological Risk Index (RI) in soils around Borholla oilfield.	62
4.25	Pollution Index (PI), Nemerow Integrated Pollution Index (NIPI) in soils around Nambar oilfield.	63
4.26	Potential ecological risk index of a single element (Er^i) and potential ecological Risk Index (RI) in soils around Nambar oilfield.	65
4.27	Correlation between PAHs and heavy metals in three different seasons in soil around Borholla oilfield.	68
4.28	Correlation between PAHs and heavy metals in three different seasons in soil around Nambar oilfield.	69
4.29 (a)	Minimum, maximum, and mean concentrations (mg kg^{-1}) of major ions in soil around Borholla oilfield during three seasons of 2011.	70
4.29 (b)	Minimum, maximum, and mean concentrations (mg kg^{-1}) of major ions in soil around Borholla oilfield during three seasons of 2013.	71
4.30	Minimum, maximum, and mean concentrations (mg kg^{-1}) of major ions in Nambar during the sample	71

	period.	
4.31	Correlation between PAHs and major ions in three different seasons in soil around Borholla oilfield.	73
4.32	Correlation between PAHs and major ions in three different seasons in soil around Nambar oilfield.	74
4.33	Results of Principle Component Analysis-Multiple Linear Regression (PCA-MLR) in surface soil around Borholla oilfield.	76
4.34	Results of Principle Component Analysis-Multiple Linear Regression (PCA-MLR) in surface soil around Nambar oilfield.	77
4.35(a)	Mean PAHs concentrations with minimum and maximum, median and geometric mean (ng ml^{-1}) in groundwater around Borholla oilfield in 3 different sampling seasons during 2011.	81
4.35(b)	Mean PAHs concentrations with minimum and maximum, median and geometric mean (ng ml^{-1}) in groundwater around Borholla oilfield in 3 different sampling seasons during 2013.	82
4.36	Mean PAHs concentrations with minimum and maximum, median and geometric mean (ng ml^{-1}) in groundwater around Nambar oilfield during 3 different sampling seasons.	85
4.37(a)	Site-wise mean concentrations of PAHs (ng ml^{-1}) with minimum and maximum, median and geometric mean values in groundwater around Borholla oilfield during pre-monsoon, monsoon and post-monsoon seasons of 2011.	89
4.37(b)	Site-wise mean concentrations of PAHs (ng ml^{-1}) with minimum and maximum, median and geometric mean values in groundwater around Borholla oilfield during pre-monsoon, monsoon and post-monsoon seasons of 2013.	90
4.38	Site-wise mean concentrations of PAHs (ng ml^{-1}) with minimum and maximum values, median and geometric mean in groundwater around Nambar oilfield during pre-monsoon, monsoon and post-monsoon seasons.	92
4.39	Concentration ranges of PAHs (ng ml^{-1}) reported from different ground water sources worldwide.	94

4.40	Percentage contributions of two-, three-, four-, five- and six -ring PAHs in groundwater around Borholla oilfield.	97
4.41	Percentage contribution of two-, three-, four-, five- and six- ring PAHs in groundwater around Nambar oilfield.	98
4.42	Mean BaP _{eq} concentrations (ng ml ⁻¹) in groundwater around Borholla oilfield during pre-monsoon, monsoon and post-monsoon seasons of both 2011 and 2013.	101
4.43	Site-wise mean BaP _{eq} concentrations (ng ml ⁻¹) of PAHs in groundwater during the whole sampling period around Borholla oilfield.	102
4.44	Mean BaP _{eq} concentrations (ng ml ⁻¹) in groundwater around Nambar oilfield during pre-monsoon, monsoon and post-monsoon seasons.	103
4.45	Site-wise mean BaP _{eq} concentrations (ng ml ⁻¹) of PAHs in groundwater during the sampling period around Nambar oilfield.	104
4.46	Seasonal mean diagnostic ratio of PAHs in groundwater around Borholla oilfield.	106
4.47	Seasonal mean diagnostic ratio of PAHs in groundwater around Nambar oilfield.	108
4.48(a)	Minimum, maximum, mean, median and geometric mean concentrations (mg lit ⁻¹) of metals in groundwater around Borholla oilfield during 2011.	112
4.48(b)	Minimum, maximum, mean, median and geometric mean concentrations (mg lit ⁻¹) of metals in groundwater around Borholla oilfield during 2013.	112
4.49	Minimum, maximum, mean, median and geometric mean concentrations (mg lit ⁻¹) of metals in groundwater around Nambar oilfield during three seasons.	113
4.50	Comparative account of heavy metal concentrations (mg lit ⁻¹) in groundwater of present study with studies from elsewhere.	114
4.51	Heavy metal Pollution Index (HPI) for groundwater around Borholla oilfield.	116
4.52	Heavy metal Pollution Index (HPI) for groundwater around Nambar oilfield.	117
4.53	Mean concentrations (mg lit ⁻¹) of total inorganic carbon (TIC), total organic carbon (TOC) and total carbon (TC) along with their maximum and minimum	118

	value in groundwater around Borholla and Nambar oilfield.	
4.54	Correlation of PAHs with TIC, TOC and TC in groundwater around Borholla oilfield.	119
4.55	Correlation of PAHs with TIC, TOC and TC in groundwater around Nambar oilfield.	120
4.56	Correlation between PAHs and heavy metals during three different seasons in groundwater around Borholla oilfield.	122
4.57	Correlation between PAHs and heavy metals during three different seasons in groundwater around Nambar oilfield.	123
4.58(a)	Minimum, maximum and mean concentrations (mg lit ⁻¹) of major ions in groundwater around Borholla oilfield during 2011.	124
4.58(b)	Minimum, maximum and mean concentrations (mg lit ⁻¹) major ions in groundwater around Borholla oilfield during 2013.	125
4.59	Minimum, maximum, and mean concentrations (mg lit ⁻¹) of major ions in groundwater around Nambar oilfield during the sampling period.	125
4.60	Correlation between PAHs and major ions during three different seasons in groundwater around Borholla oilfield.	127
4.61	Correlation between PAHs and major ions during three different seasons in groundwater around Nambar oilfield.	128
4.62	Averages of signature of PAHs (PAH _i / ΣPAHs) in surface soil and groundwater around Borholla oilfield.	129
4.63	Correlation of seasonal average source signature of PAHs of surface soil and groundwater around Borholla oilfield.	130
4.64	Averages of signature of PAHs (PAH _i / ΣPAHs) in surface soil and groundwater around Nambar oilfields.	130
4.65	Correlation of seasonal average source signature of PAHs of surface soil and groundwater around Nambar oilfields.	131

LIST OF FIGURES

Chapter	Figure no.	Figure caption	Page no.
1	1.1	Molecular structures of 16 USEPA PAHs.	3
	1.2	The distribution and transfer of PAHs in various environmental matrices (Modified USEPA).	8
3	3.1	The study area (map prepared by integrating different thematic layers in GIS environment). Sampling locations in Borholla (B1-B15) and in Nambar (N1-N21).	2
	3.2	Chromatogram of external standard containing 16 PAHs mixture along with their retention time.	9
	3.3	A schematic illustration of sample preparation and analysis for PAHs and Heavy metals.	13
4	4.1	Seasonal trends of concentrations (ng g^{-1}) of Σ PAHs, Σ LMWPAHs and Σ HMWPAHs around Borholla oilfield during the years 2011 and 2013.	5
	4.2	Seasonal concentration trends (ng g^{-1}) of individual PAHs around Borholla oilfield during the study period.	6
	4.3	Seasonal concentrations trends (ng g^{-1}) of Σ PAHs, Σ LMWPAHs and Σ HMWPAHs around Nambar oilfield.	8
	4.4	Seasonal concentration trends (ng g^{-1}) of individual PAHs around Nambar oilfield during the study period.	9
	4.5	One twenty hour backward trajectories arriving the study area on (a) 27 th April 2011 (pre-monsoon), (c) 25 th July 2011(monsoon) and (e) 19 th November 2011(post-monsoon); (b), (d) and (f) are respective vertical temperature profiles.	11-12
	4.6	One twenty hour backward trajectories arriving the study area on (a) 13 th April 2013 (pre-monsoon), (c) 25 th July 2013 (monsoon) and (e) 14 th November 2013 (post-monsoon); (b), (d) and (f) are respective vertical temperature profiles.	12-13

Chapter	Figure no.	Figure caption	Page no.
4	4.7	Spatial variations of Σ PAHs concentrations (ng g^{-1}) of soils around Borholla oilfield in three different sampling seasons during (a) 2011 and (b) 2013.	17
	4.8	Spatial variations of Σ PAHs concentrations (ng g^{-1}) in soils around Nambar oilfield during different seasons.	18
	4.9	The percentage compositions of two-, three-, four-, five- and six- ring PAHs at different sites during different seasons of 2011 and 2013 around Borholla oilfield.	21
	4.10	Percentage compositions of two-, three-, four-, five- and six- ring PAHs in soils around Borholla oilfield during (a) 2011, (b) 2013 and (c) mean two years of sampling.	22
	4.11	Percentage compositions of two-, three-, four-, five- and six- ring PAHs at different sampling sites around Nambar oilfield in three different seasons and mean percentage of each season.	24
	4.12	Abundance of carcinogenic PAHs and non-carcinogenic PAHs in soils around Borholla oilfield during the study period.	26
	4.13	Abundance of carcinogenic PAHs and non-carcinogenic PAHs in soils around Nambar oilfield during the study period.	26
	4.14	Isomer pair ratio of soils around Borholla oilfield: (a) Ant/(Ant+Phe) vs. Flu/(Flu + Pyr), (b) BaA/(BaA + Chr) vs IcdP/(IcdP + BgP) and (c) Phe/(Phe+Ant) vs Fla/(Fla + Pyr).	34
	4.15	Isomer pair ratio of soil around Nambar oilfield: (a) Ant/ (Ant+Phe) vs. Flu/ (Flu + Pyr), (b) BaA/ (BaA + Chr) vs IcdP/ (IcdP + BgP) and (c) Phe/ (Phe+Ant) vs Fla/ (Fla + Pyr).	37
	4.16	Linear regression analyses between PAHs ratios and tracer species/ratios in soil around Borholla oilfield.	38
	4.17	Linear regression analyses between PAHs ratios and tracer species/ratios in soil around Nambar oilfield.	39
	4.18	Linear regression of Σ PAHs, Σ LMWPAHs and Σ HMWPAHs on OC around Borholla oilfield during the sampling period.	48

4.19	Linear regression of Σ PAHs, Σ LMWPAHs and Σ HMWPAHs on BC around Borholla oilfield during the sampling period.	49
4.20	Linear regression of Σ PAHs, Σ LMWPAHs and Σ HMWPAHs on OC around Nambar oilfield during the sampling period	50
4.21	Linear regression of Σ PAHs, Σ LMWPAHs and Σ HMWPAHs on BC around Nambar oilfield during the sampling period.	51
4.22	Pollution Load Index (PLI) in surface soil in different seasons around Borholla oilfield.	61
4.23	Pollution Load Index (PLI) in surface soil in different seasons around Nambar oilfield.	64
4.24	Seasonal trends of concentrations (ng ml^{-1}) of Σ PAHs, Σ LMWPAHs and Σ HMWPAHs in groundwater around Borholla oilfield during the years 2011 and 2013.	83
4.25	Seasonal concentration (ng ml^{-1}) trends of individual PAHs in groundwater around Borholla oilfield during the study period.	84
4.26	Seasonal trends of concentrations (ng ml^{-1}) of Σ PAHs, Σ LMWPAHs and Σ HMWPAHs in groundwater around Nambar oilfield during the sampling period.	86
4.27	Seasonal concentration (ng ml^{-1}) trends of individual PAHs in groundwater around Nambar oilfield during the study period.	87
4.28	Spatial variations of Σ PAHs concentrations (ng ml^{-1}) of groundwater around Borholla oilfield in three different sampling seasons during (a) 2011 and (b) 2013.	91
4.29	Spatial variations of Σ PAHs concentrations (ng ml^{-1}) of groundwater around Nambar oilfield in three different sampling seasons.	93
4.30	The percentage compositions of two-, three-, four-, five- and six- ring PAHs at different sites during different seasons of 2011 and 2013 around Borholla oilfield.	95
4.31	Percentage compositions of two-, three-, four-, five- and six- ring PAHs around Borholla oilfield during (a) 2011 (b) 2013 and (c) mean two year of sampling.	96
4.32	Percentage compositions of two-, three-, four-, five- and six- ring PAHs at different sampling sites around	98

	Nambar oilfield in three different seasons and mean percentage of each season.	
4.33	Abundance of carcinogenic and non-carcinogenic PAHs in groundwater around Borholla oilfield during the study period.	99
4.34	Abundance of carcinogenic and non-carcinogenic PAHs in groundwater around Nambar oilfield during the study period.	100
4.35	Isomer pair ratio of groundwater around Borholla oilfield: (a) Ant/(Ant + Phe) vs Flu/(Flu + Pyr) and (b) Phe/ (Phe+Ant) vs Fla/(Fla + Pyr)	107
4.36	Isomer pair ratio of groundwater around Borholla oilfield: (a) Ant/(Ant+Phe) vs. Flu/(Flu + Pyr), (b) Phe/(Phe+Ant) vs Fla/(Fla + Pyr)	110