

# List

## OF FIGURES

---

### Chapter 1

1.1	Post-harvest treatment of Phytoremediator plants	20
1.2	General Morphology of the Diatom frustule	22
1.3	Diatom, <i>Navicula</i> sp	23
1.4	Scanning electron microscopy of four different diatom species representing centric and pennate symmetry	25

### Chapter 3

3.1	<i>Eichhornia crassipes</i> (water hyacinth) plant	96
3.2	Percent Pb removal by <i>E. crassipes</i> at 15, 20 and 30 ppm Pb at 2 <sup>nd</sup> day	98
3.3	Photosynthetic rate in <i>E. crassipes</i> at different concentrations of Pb for 10 days of exposure	99
3.4	Total chlorophyll content of <i>E. crassipes</i> at different pH with 30.0 mg/L initial conc. as observed after 5 <sup>th</sup> and 10 <sup>th</sup> day of exposure	100
3.5	(a) SEM image of <i>E. crassipes</i> leaves after 2nd days in control; (b) The EDX spectrum of cross-section of leaf of control of <i>E. crassipes</i>	101
3.5	(c) SEM images of longitudinal sections of <i>E. crassipes</i> leaf treated with Pb (30 mg/L); (d) EDX taken from the leaf of Pb exposed plants	102
3.6	(a, b) SEM-EDX images of <i>Eichhornia crassipes</i> shoot (petiole) biomass without absorption of Pb <sup>2+</sup> ions	102
3.6	(c, d) SEM-EDX images of <i>Eichhornia crassipes</i> petiole biomass with absorption of Pb <sup>2+</sup> ions	103
3.7	(a, b) The SEM micrographs showed changes of the vascular cells of the root samples exposed to 30mg/L Pb treatment; (b). EDX confirmed the presence of Pb	103

3.7	(c, d) SEM micrograph of control root cross section of <i>E. crassipes</i>	104
3.8	Pb accumulation in vascular bundles of <i>E. crassipes</i>	104
3.9	Scanning electron micrograph and EDX spectra of Pb distribution in root of <i>E. crassipes</i> . Pb internalization was confirmed by SEM (a); studies in combination with EDX (b)	104
3.10	SEM-EDX spectra of Pb distribution in epidermis of root of <i>E. crassipes</i>	105
3.11	SEM micrograph of <i>E. crassipes</i> root (a) control; (b) Pb (II) loaded.	105
3.12	EDX micrograph of <i>E. crassipes</i> root (c) control; (d) Pb (II) loaded Magnification: 3000X	106
 <b>Chapter 4</b>		
4.1	Some of the morphological changes in <i>T. natans</i> on 3 <sup>rd</sup> , 5 <sup>th</sup> and 7 <sup>th</sup> day intervals	120-121
4.2	Graphical representation of effect of arsenic on total chlorophyll content	122
4.2(A)	Graphical representation of amount of proline synthesis under stress condition	122
4.3	(A) <i>T. natans</i> in control condition; (B) <i>T. natans</i> after arsenic treatment; (C) SEM micrograph of leaves of control <i>T. natans</i> ; (D) Leaves of arsenic treated <i>T. natans</i> ; (E) EDX of leaves of control <i>T. natans</i> ; (F) EDX of leaves of arsenic treated <i>T. natans</i>	126-127
4.3	(G) Arsenic treated shoot of <i>T. natans</i> ; (H) Control shoot of <i>T. natans</i> ; (I) EDX of shoot of control <i>T. natans</i> ; (J) EDX of shoot of arsenic treated <i>T. natans</i>	127-128
4.3	(K) Internal distributions of electron dense deposition of elements in the vascular bundles of <i>T. natans</i> shoot (As treated)	128
4.3	(L) Arsenic treated root of <i>T. natans</i> ; (M) Control root of <i>T. natans</i> ; (N) EDX of root of control <i>T. natans</i> ; (O) EDX of	129

## List of figures

	root of arsenic treated <i>T. natans</i>	
4.4.1	FTIR spectra of control leaf of <i>T. natans</i>	132
4.4.2	FTIR spectra of arsenic treated leaf	132
4.4.3	FTIR spectra of control shoot of <i>T. natans</i>	132
4.4.4	FTIR spectra of arsenic treated shoot	133
4.4.5	FTIR spectra of control root of <i>T. natans</i>	133
4.4.6	FTIR spectra of arsenic treated root	133
 <b>Chapter 5</b>		
5.1	<i>M. hastata</i> in natural condition	151
5.2	Graphical representation of total chlorophyll content in <i>M. hastata</i>	156
5.3	Phytoaccumulation of Cd by <i>M. hastata</i>	157
5.4	SEM- EDX micrographs of leaf surface of <i>M. hastata</i> (A–F) (A) leaf epidermis of plant grown in 15mg/L Cd solution (1400X); (B) leaf epidermis of control plant (650X); (C) magnified view of stomata showing diameter in control leaf (1500X); (D) showing closing of stomata (1500X); (E) EDX spectra of control plant leaf; (E) magnified view of plate; (F) EDX spectra of Cd treated plant, showing the presence of Cd ions along with other ions	159-160
5.5	SEM- EDX micrographs (A) Shoot of <i>M. hastata</i> exposed to 15 mg/L Cd for 10 days ; (B) the control shoot ; (C) EDX spectra of control shoot; (D) EDX spectra of Cd treated shoot	161
5.6	(A, B) SEM micrographs Control root; (C, D) Cd treated <i>M. hastata</i> root, showing some surface shrinking after metal binding; (E) Energy-dispersive X- ray spectra of control root; (F) EDX spectra of treated root confirmed the presence of Cd	162
5.7	TEM Micrograph of chloroplast in the leaf cells of <i>M. hastata</i> (A) Control leaf showing numerous chloroplasts; (B) showing one chloroplast; (C) Cd treated with 15 mg/L for 10 days; (D) to show the effect of Cd on thylakoid inside	163

	the chloroplast	
5.8	TEM Micrograph of the root mesophyll cell mitochondrion (A) control <i>M. hastata</i> plant; (B) micrograph of Cd treated mitochondria; (C) showing electron dense deposition within the mitochondria	164-165
5.9	TEM of <i>M. hastata</i> root mesophyll cells nucleus (A) control; (B) 15/L Cd treated	165
5.10	TEM micrograph of <i>M. hastata</i> root cortical cell (A) control cell showing no metal deposition along with the cell wall; (B) 15 mg/L of Cd treated showing Cd <sup>2+</sup> deposition along the surface and in the cell wall	166
5.11	TEM micrograph of vacuoles of root cells (A) control showing no deposition within the vacuole; (B) Metal accumulation within the vacuole	167
5.12	FTIR spectra of control leaf biomass of <i>M. hastata</i>	171
5.13	FTIR spectra of Cd treated leaf biomass of <i>M. hastata</i>	171
5.14	FTIR spectra of control shoot biomass of <i>M. hastata</i>	171
5.15	FTIR spectra of Cd treated shoot of <i>M. hastata</i>	172
5.16	FTIR spectra of control root biomass of <i>M. hastata</i>	172
5.17	FTIR spectra of Cd treated root biomass of <i>M. hastata</i>	172
<b>Chapter 6</b>		
6.1	<i>E. crassipes</i> and <i>T. natans</i> plants in control condition	200
6.2	Graphical representation of total chlorophyll content in <i>T. natans</i>	203
6.3	Graphical representation of total chlorophyll content in <i>E. crassipes</i>	204
6.4	SEM-EDX of fresh shoot of <i>T. natans</i> control (i, a); and exposed to Sb (ii, b) for 10 <sup>th</sup> days	211-212
6.4	SEM- EDX of control (d, iii); and treated (Sb) (c, iv) root of <i>T. natans</i> plant	212-213
6.4	SEM- EDX of control (e, v); and treated (Sb) (f, vi) shoot of <i>E. crassipes</i> plant	213-214

List of figures

6.4	SEM- EDX of control (g, vi) and Sb treated root (h, viii) of <i>E. crassipes</i>	215
6.4	Sb accumulation by the root hair of <i>E. crassipes</i> ; (h) Root hair of control plant; (i) Root hair of Sb treated plant	216
6.5	TEM micrograph shows a young mesophyll cell of control plant of <i>T. natans</i> showing numerous chloroplasts (A); A chloroplast with well-organized inner membranes with both grana and stroma in a cell of a young control leaf of <i>T. natans</i> (B); Chloroplast with a well-developed and organized thylakoid system in a cell of a mature control leaf (B); Damaged disintegrating chloroplasts showing the presence of irregular, rounded shape, numerous starch grains (C); disturbance of the orientation of the grana (D)	218
6.6	TEM micrograph shows a bean - shaped mesophyll chloroplast of control plant <i>E. crassipes</i> . The grana are orderly arranged and interconnected with stroma thylakoids (A); A mesophyll chloroplast of <i>E. crassipes</i> plant treated with 7.47 mg/L Sb solution showing a large starch grain and disturbed grana structure (B)	218
6.7	Transmission electron micrographs of root cells of <i>E. crassipes</i> showing numerous mitochondrion in the cortical cells of control plant (A); Ruptured mitochondria and reduced number of cristae of mitochondria in the cortical cells exposed at 7.47 mg/L SbCl <sub>3</sub> solution (B)	220
6.8	TEM micrographs of transverse section of roots of <i>T. natans</i> ; (A) Control plant showing the vacuoles; (B) SbCl <sub>3</sub> treated plant root showing the deposition within the vacuole	220
6.9	TEM micrographs of root cells of <i>E. crassipes</i> ; In the control plant root cell no deposition within the vacuole (A); black deposits were found in the vacuole of SbCl <sub>3</sub> treated plant root cells (B)	221
6.10	TEM micrograph of root cells of <i>T. natans</i> showing mitochondria, vacuole and Golgi appear well preserved (A); After application of Sb, the disorganization of microtubules	221

	in Golgi body appeared (B)	
6.11	TEM micrographs of root cells of <i>T. natans</i> of control plant (A); The cell wall treated with 7.47mg L <sup>-1</sup> Sb, showing numerous black deposits in the wall of the cell (B)	222

## Chapter 7

7.1	Diatoms in the environment (Source: Author)	239
7.2	SEM image of a pinnate diatom of <i>Navicula</i> species (Source: author)	240
7.3	Growth curve of the <i>Navicula</i> sp.in WC media	251
7.4	Growth Curve of different Subcultures of diatom <i>Navicula</i> sp.	251
7.5	Variation of chlorophyll with As concentration	252
7.6	Living diatoms seen in light microscope. <i>Navicula</i> sp.; Microscope magnification was set on 400 times. Diatom growth in Control media (A) and As containing media (B)	252
7.7	As accumulation in <i>Navicula</i> species	254
7.8	SEM micrograph of control group of <i>Navicula</i> colony (A); An X-ray emission spectrum from an unexposed diatom sample showed no As peaks (B)	255
7.8	SEM micrograph of As treated of <i>Navicula</i> colony (C, D); An X-ray emission spectrum from an As exposed diatom sample showed As peaks (E)	255-256
7.9	SEM images showing the pore sizes of control media (A); (B) Pore sizes of diatom <i>Navicula</i> within Arsenic Medium	257
7.10	Scanning Electron Microscopic view and EDX of a diatom, <i>Navicula</i> sp. It showed nanometer sized in control (A, D); (B, II) in As media	258-259
7.11	XRD pattern of the diatom <i>Navicula</i> sp.	259
7.12	FTIR spectra of diatom <i>Navicula</i> sp. treated with As solution	260
7.13	FTIR spectra of control diatom group	261
7.14	Zeta potential profile of control (A); As treated diatom(B)	263