

List of Figure

2.1	A cross sectional view of a myelinated axon	11
2.2	Nerve conduction study with Medicaid EMG machine Model No EMG-2000.	16
2.3	Nerve biopsy of (a) Patient's id: 015 and (b) Patient's id: 058, collected from Pathology Lab, GNRC Hospital, Guwahati, Assam, India.	16
2.4	Skin biopsy of (a) Patient's id: 023 and (b) Patient's id: 027, collected from Pathology Lab, GNRC Hospital, Guwahati, Assam, India	17
2.5	An equivalent electric circuit of H-H model.	24
2.6	An electric circuit of a single myelinated axon.	28
2.7	An electric circuit of a single demyelinated axon.	29
2.8	NCV versus demyelination factor (γ).	30
2.9	An electric circuit of a coupled model of axon.	31
2.10	NCV versus demyelination factor (γ) in a coupled model of axons.	32
3.1	(a) A bundle of myelinated nerve fibers, (b) SEM image of a bundle of axons in a myelinated nerve and (c) schematic representation of a bundle of axons.	48
3.2	An equivalent electric circuit of a myelinated nerve fiber	49
3.3	(a) Estimation of NCV in terms of alignment factor and (b) variance of NCV with increasing alignment of axons in the bundle.	55
3.4	<i>Duttaphrynus melanostictus</i>	56
3.5	AD instrument with nerve chamber used to record nerve signals.	57
3.6	Electric neuro signal of a normal sciatic nerve of toad	58
3.7	SEM image of a myelinated nerve fiber.	59
4.1	(a) A bundle of axons in a demyelinated nerve fiber with degraded myelin sheath, (b) image representing a bundle of axons in a demyelinated nerve obtained from scanning electron microscope (SEM) and (c) schematic representation of bundle of	65

	axons corresponding to SEM image.	
4.2	Electric circuit model of a demyelinated nerve.	66
4.3	Estimation of NCV in terms of demyelination factor with alignment parameter A .	71
4.4	Electric neuro signals of (a) control, and nerves treated with (b) $0.1\mu\text{g/ml}$, (c) $1.0\mu\text{g/ml}$ and (d) $10\mu\text{g/ml}$ of crude venom concentrations respectively	74
4.5	SEM images of (a) control, and nerves treated with (b) $0.1\mu\text{g/ml}$, (c) $1.0\mu\text{g/ml}$ and (d) $10\mu\text{g/ml}$ of crude venom concentrations respectively	75
4.6	A graphical representation of NCV versus demyelinating factor with an inset of degree of demyelination versus crude venom concentration.	77
4.7	Effect of purified Nk-PLA ₂ with different concentration on sciatic nerves of toad and their nerve conduction signals (proximal and distal action potential) with corresponding SEM images of their sciatic nerves.	78
4.8	Effect of purified 3Ftx with different concentration on nerve conduction signals.	79
4.9	Mechanism of channel blocking in cell membrane	81
5.1	(a) A myelinated nerve (nerve I) being coupled with a demyelinated nerve (nerve II) and (b) image of nerve I and nerve II to be coupled for the formation of coupled nerve model obtained from scanning electron microscope (SEM).	87
5.2	An electric circuit model of a couple nerve fibers.	88
5.3	Estimation of NCV in the coupled nerve for aligned ($A=1$) and misaligned ($A=1/2$) axons in the model.	93
5.4	A coupled nerve in a nerve chamber.	93
5.5	Electro neuro signals in demyelinated and coupled nerve.	97
5.6	Theoretical estimation of NCV in demyelinated and coupled nerve in terms of alignment factor at (a) $A=1/2$ and (b) $A=1$.	99
5.7	Comparison of AP in demyelinated and coupled nerve.	100
5.8	Blocking mechanism and coupling effect in coupled nerve.	101