

LIST OF FIGURES

Chapter	Figure	Title	Page No.
1	1.1	Schematic diagram for structural representation of layered double hydroxide (LDH)	1-4
	1.2	Various synthetic methodologies of LDH	1-5
	1.3	Schematic diagram of co-precipitation method for preparation of LDH	1-6
	1.4	Various properties of LDHs	1-9
	1.5	Various applications of LDH	1-10
	1.6	Various methods for treatment of organic pollutants	1-13
	1.7	Maximum adsorption capacities (q_m , mg/g) for DB, RY, AR, DR and BB dyes at 298 K using LDHs-SDS	1-15
	1.8	Dye removal (%) of methyl blue from aqueous solution over LDH and LDH-carbon dots adsorbents	1-16
	1.9	Schematic diagram of the proposed mechanism for photocatalytic degradation of organic pollutants	1-19
	1.10	Degradation (%) of rhodamine B (RB) on photodegradation by Zn/M-NO ₃ -LDHs	1-21
	1.11	Photocatalytic efficiencies of Ni/Ti LDH and various commercially available catalysts for degradation of methylene blue in aqueous solution under visible light	1-22
	1.12	Degradation (%) of phenol over ZnTi-LDH and derived mixed oxides calcined at different temperatures	1-23
	1.13	Schematic diagram for nitro-aldol condensation (Henry) reaction	1-23
	1.14	Schematic diagram for general mechanism of nitro-aldol condensation (Henry) reaction	1-24
	1.15	Schematic representation for various applications of Henry reaction product, 2-nitro-alkanol	1-25

	1.16	Difference between conventional and microwave heating	1-28
3	3.1	The molecular structure of various organic dye pollutants used in the present investigation	3-1
	3A.1	(a) PXRD patterns and (b) High angle 2θ value, (c) Relative Intensity and (d) Crystallinity (%) at some 2θ angles of all the LDH samples	3-4
	3A.2	Relationship of nickel content with unit cell parameters in LDH samples	3-4
	3A.3	FTIR spectra of all the LDH samples	3-6
	3A.4	(a) TGA graphs, (b) Micro-Raman spectra, (c) UV-vis DRS spectra and (d) Band gap energy calculated from UV-vis DRS of all LDH samples	3-7
	3A.5	SEM images of (a) LDH1, (b) LDH2, (c) LDH3 and (d) LDH4	3-9
	3A.6	TEM images at different resolution (a), (b) & (c) and SAED patterns (d) of LDH3	3-9
	3A.7	N ₂ adsorption-desorption isotherms and pore size distribution (inset) of (a) LDH1, (b) LDH2, (c) LDH3 and (d) LDH4	3-10
	3A.8	Effect of (a) contact time, (b) adsorbent dosage, (c) initial dye concentration and (d) solution pH on adsorption of MO onto LDH adsorbents (Optimized adsorption conditions: MO concentration = 20 mg/L, adsorbent dose = 5 mg, equilibrium time = 90 min, pH = 6 and $T = 25\text{ }^{\circ}\text{C}$)	3-12
	3A.9	UV-vis spectra for adsorption of MO onto (a) LDH1, (b) LDH2, (c) LDH3 and (d) LDH4 at various time intervals	3-14
	3A.10	Pseudo-first order kinetic plots for adsorption of MO onto (a) LDH1, (b) LDH2, (c) LDH3 and (d) LDH4	3-15

3A.11	Pseudo-second order kinetic plots for adsorption of MO onto (a) LDH1, (b) LDH2, (c) LDH3 and (d) LDH4	3-16
3A.12	Intraparticle diffusion plot for adsorption of MO onto all the LDH samples	3-16
3A.13	Langmuir isotherm plots for the adsorption of MO onto (a) LDH1, (b) LDH2, (c) LDH3 and (d) LDH4	3-18
3A.14	Freundlich isotherm plots for the adsorption of MO onto (a) LDH1, (b) LDH2, (c) LDH3 and (d) LDH4	3-19
3A.15	D–R isotherm plots for adsorption of MO onto (a) LDH1, (b) LDH2, (c) LDH3 and (d) LDH4	3-19
3A.16	Schematic diagram for efficient removal of MO over LDH3 adsorbent	3-20
3A.17	(a) UV-vis spectra of MO before and after adsorption onto LDH3 and inset is photographs of MO before (left) and after adsorption (right), (b) PXRD patterns of LDH3 before and after adsorption of MO	3-21
3A.18	SEM images of LDH3 (a) before and (b) after adsorption of MO	3-21
3A.19	(a) FTIR spectra of LDH3 after adsorption of MO and inset is enlarged view of the spectra in the wavenumber range of 1000–1300 cm^{-1} and (b) FTIR spectra of pure MO and LDH3	3-22
3A.20	% Dye uptake of various anionic dye pollutants over LDH3 (Optimized conditions: MO concentration = 20 mg/L, adsorbent dose = 5 mg, equilibrium time = 90 min, pH = 6 and $T = 25\text{ }^{\circ}\text{C}$)	3-23
3A.21	Reusability of LDH3 for adsorption of MO	3-24
3B.1	(a) PXRD patterns and (b) FTIR spectra of LDH samples	3-27
3B.2	(a) TGA graphs and (b) UV-vis DRS spectra of LDH	3-28

	samples	
3B.3	SEM images of LDH samples	3-29
3B.4	(a–c) TEM images at different resolution and (d) SAED patterns of CuMgAl ₄ LDH	3-29
3B.5	N ₂ adsorption-desorption isotherms of LDH samples	3-30
3B.6	Dye removal (%) of MO over various LDHs ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-31
3B.7	Dye removal (%) of different anionic and cationic dyes over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-32
3B.8	UV-vis spectra of the anionic dyes after adsorption over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-33
3B.9	Influence of (a) contact time (min) and (b) adsorbent amount (mg) on dye removal (%) of MO, BTB, EBT and CR over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, $T = 25$ °C, pH = 7)	3-34
3B.10	Influence of pH on dye removal (%) of MO, BTB, EBT and CR over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C)	3-35
3B.11	Influence of initial dye concentrations on dye removal (%) of MO, BTB, EBT and CR over CuMgAl LDH ($V_{\text{solution}} = 20$ mL, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-36
3B.12	Pseudo-first order kinetic plots for adsorption of MO, BTB, EBT and CR over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-37
3B.13	Pseudo-second order kinetic plots for adsorption of MO, BTB, EBT and CR over CuMgAl ₄ LDH (V_{solution}	3-38

	= 20 mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	
3B.14	Intraparticle diffusion kinetic plot for adsorption of MO, BTB, EBT and CR over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-39
3B.15	Langmuir isotherm plots for adsorption of MO, BTB, EBT and CR over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-40
3B.16	Freundlich isotherm plots for adsorption of MO, BTB, EBT and CR over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-41
3B.17	FTIR spectra for adsorption of MO, BTB, EBT and CR over CuMgAl ₄ LDH	3-43
3B.18	Multicyclic adsorption of MO, BTB, EBT and CR over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7)	3-43
3B.19	Selective adsorption of dye from a mixture of anionic and cationic dyes over CuMgAl ₄ LDH ($V_{\text{solution}} = 20$ mL, $C_o = 20$ mg/L, adsorbent amount = 5 mg, $T = 25$ °C, pH = 7 for MO and 11 for MB)	3-44
3C.1	Powder X-ray diffraction patterns of all the samples	3-47
3C.2	(a) FTIR spectra and (b) TGA curves of all the samples	3-48
3C.3	(a) SEM image, (b) TEM image, (c) lattice fringes of (003) plane and (d) SAED pattern of NiAl-S ₁ LDH	3-49
3C.4	Dye removal (%) of CR on various LDHs with time (10 mg/L CR, 10 mg adsorbent, $T = 25$ °C)	3-50
3C.5	(a) UV-vis spectra of CR on adsorption with different time (10 mg/L CR, 10 mg adsorbent), (b) effect of time (10 mg adsorbent), (c) effect of adsorbent amount	3-51

	(5–20 mg adsorbent) and (d) effect of initial dye concentrations (10–60 mg/L CR, 10 mg adsorbent) on adsorption of CR onto NiAl-S ₁ LDH at $T = 25\text{ }^{\circ}\text{C}$	
3C.6	Dye removal (%) of CR on NiAl-S ₁ LDH at various pH ranges	3-53
3C.7	(a) Adsorption isotherm, (b) Langmuir, (c) Freundlich and (d) D–R isotherm models for adsorption of CR on NiAl-S ₁ LDH (10 mg adsorbent, $T = 25\text{ }^{\circ}\text{C}$)	3-54
3C.8	(a) Pseudo-first order, (b) pseudo-second order, (c) intraparticle diffusion and (d) van't Hoff plot for adsorption of CR on NiAl-S ₁ LDH (10 mg adsorbent)	3-56
3C.9	XRD patterns of NiAl-S ₁ LDH before and after adsorption of CR	3-58
3C.10	SEM images of NiAl-S ₁ LDH before and after adsorption of CR	3-59
3C.11	FTIR spectra of CR, bare NiAl-S ₁ LDH and CR adsorbed NiAl-S ₁ LDH	3-59
3C.12	Dye removal (%) of CR with number of cycles on NiAl-S ₁ LDH (10 mg/L CR, 10 mg adsorbent, $T = 25\text{ }^{\circ}\text{C}$)	3-61
4	4.1 Powder XRD patterns of ZnFe LDH	4-4
	4.2 FTIR spectrum of ZnFe LDH	4-5
	4.3 TGA thermogram of ZnFe LDH	4-6
	4.4 SEM images at (a) lower and (b) higher resolutions, (c) TEM image and (d) SAED patterns of ZnFe LDH	4-7
	4.5 (a) UV-vis diffuse reflectance spectra and (b) corresponding band gap energy of ZnFe LDH	4-7
	4.6 (a) N ₂ adsorption-desorption isotherms, and (b) pore size distribution (PSD) curve of ZnFe LDH	4-9
4A.1	UV-visible spectra for photocatalytic degradation of phenol under (a) visible and (b) UV light irradiation;	4-11

	(c) Photocatalytic degradation of phenol as a function of irradiation time under different light irradiation; (d) Photocatalytic degradation of various phenolic compounds as a function of irradiation time under visible light irradiation (Condition: $C_o = 0.5$ mM, $V = 20$ mL, catalyst amount = 10 mg, pH = 7)	
4A.2	UV-visible spectra for photocatalytic degradation of (a) 2CP, (b) 3CP and (c) 4CP; (d) Degradation (%) of 2CP, 3CP and 4CP with irradiation time over ZnFe LDH under visible light irradiation (Condition: $C_o = 0.5$ mM, $V = 20$ mL, catalyst amount = 10 mg, pH = 7)	4-12
4A.3	Effect of amount of ZnFe LDH on photocatalytic degradation of phenol, 2CP, 3CP and 4CP under visible light irradiation (Condition: $C_o = 0.5$ mM, $V = 20$ mL, pH = 7)	4-13
4A.4	Effect of initial concentration of phenol, 2CP, 3CP and 4CP on photocatalytic activity of ZnFe LDH for degradation under visible light irradiation (Condition: $V = 20$ mL, catalyst amount = 10 mg, pH = 7)	4-14
4A.5	Effect of pH on photocatalytic activity of ZnFe LDH for degradation of phenol and its compounds under visible light irradiation (Condition: $C_o = 0.5$ mM, $V = 20$ mL, catalyst amount = 10 mg)	4-15
4A.6	Pseudo-first order kinetic plots for photocatalytic degradation of phenol, 2CP, 3CP and 4CP over ZnFe-LDH under visible light irradiation (Condition: $C_o = 0.5$ mM, $V = 20$ mL, catalyst amount = 10 mg, pH = 7)	4-17
4A.7	Schematic diagram of the proposed mechanism for photocatalytic degradation of phenol over ZnFe-LDH	4-18
4A.8	Proposed degradation pathway for photocatalytic degradation of phenol initiated by \cdot OH radical	4-20

4A.9	Optimized structures of all species involved in the Phenol + $\cdot\text{OH}$ reaction at B3LYP/6-31+G(d,p) level of theory	4-21
4A.10	Potential energy surface (PES) diagram for the degradation of Phenol + $\cdot\text{OH}$ reaction at B3LYP/6-31+G(d,p)	4-24
4A.11	Recyclability of ZnFe LDH for photocatalytic degradation of phenol under visible light irradiation	4-25
4B.1	Photolysis, adsorption and photocatalysis of MB and RhB over ZnFe LDH under UV light irradiation (Conditions: $C_o = 10$ mg/L, $V_{\text{solution}} = 50$ mL, catalyst amount = 5 mg for MB and 10 mg for RhB, pH = 7)	4-28
4B.2	UV-visible spectra for photocatalytic degradation of MB and RhB over ZnFe LDH under UV light irradiation ($C_o = 10$ mg/L, $V_{\text{solution}} = 50$ mL, catalyst amount = 5 mg for MB and 10 mg for RhB, pH = 7)	4-28
4B.3	Photolysis, adsorption and photocatalysis of MB and RhB over ZnFe LDH under visible light irradiation (Conditions: $C_o = 10$ mg/L, $V_{\text{solution}} = 50$ mL, catalyst amount = 5 mg for MB and 10 mg for RhB, pH = 7)	4-29
4B.4	UV-visible spectra for photocatalytic degradation of MB and RhB over ZnFe LDH under visible light irradiation ($C_o = 10$ mg/L, $V_{\text{solution}} = 50$ mL, catalyst amount = 5 mg for MB and 10 mg for RhB, pH = 7)	4-30
4B.5	Effect of catalyst amount on photocatalytic degradation of MB and RhB with irradiation time under UV light irradiation (Conditions: $C_o = 10$ mg/L, $V_{\text{solution}} = 50$ mL, catalyst amount = 5 mg for MB and 10 mg for RhB, pH = 7)	4-31
4B.6	Effect of initial dye concentrations on photocatalytic degradation of MB and RhB with irradiation time	4-32

	under UV light irradiation (Conditions: $V_{\text{solution}} = 50$ mL, catalyst amount = 5 mg for MB and 10 mg for RhB, pH = 7)	
4B.7	Effect of pH on photocatalytic degradation of MB and RhB with irradiation time under UV light irradiation (Conditions: $C_0 = 10$ mg/L, $V_{\text{solution}} = 50$ mL, catalyst amount = 5 mg for MB and 10 mg for RhB)	4-33
4B.8	Kinetic plots of $\ln(C_0/C)$ as a function of time for photodegradation of MB and RhB under UV light irradiation	4-33
4B.9	Recyclability test for photocatalytic degradation of MB and RhB over ZnFe LDH. (Conditions: $C_0 = 10$ mg/L, $V_{\text{solution}} = 50$ mL, catalyst amount = 5 mg for MB and 10 mg for RhB, pH = 7)	4-36
4B.10	XRD patterns of fresh and reused ZnFe LDH photocatalyst	4-37
5	5A.1 X-ray powder diffraction patterns of a) MgAl, NiMgAl and CoMgAl LDHs and; (b) NiMgAl (O) and CoMgAl (O) mixed oxides	5-4
	5A.2 FTIR spectra of (a) MgAl, NiMgAl and CoMgAl LDHs and; (b) NiMgAl (O) and CoMgAl (O) mixed oxides	5-6
	5A.3 TGA curves of MgAl, NiMgAl and CoMgAl LDHs	5-6
	5A.4 SEM images of (a) MgAl, (b) NiMgAl and (c) CoMgAl LDHs	5-7
	5A.5 (a) N_2 adsorption-desorption isotherms and (b) pore size distribution curves of MgAl (O), NiMgAl (O) and CoMgAl (O) mixed oxides	5-8
	5A.6 N_2 adsorption-desorption isotherms of NiMgAl (O) mixed oxide at different calcination temperatures	5-13
	5A.7 Turnover frequency (TOF) of MgAl (O), NiMgAl (O)	5-14

	and CoMgAl (O) mixed oxides	
5B.1	Powder XRD patterns of (a) LDHs and (b) LDH derived mixed oxides (o – NiO, * - MgO and ^ - MgAl ₂ O ₄); and FTIR spectra of (c) LDHs and (d) LDH derived mixed oxides	5-18
5B.2	Thermogravimetric analysis curves of LDHs	5-19
5B.3	SEM images of LDHs and inset shows for LDH derived mixed oxides	5-20
5B.4	N ₂ adsorption-desorption isotherms of LDH derived mixed oxides	5-21
5B.5	Recyclability test of the catalyst LDH3 (O) (Reaction conditions: 1 mmol 4-nitrobenzaldehyde, 10 mmol nitromethane, 10 mg catalyst, 40% MW power (280 Watt), Temperature 50 °C)	5-27

LIST OF TABLES

Chapter	Table	Title	Page No.
1	1.1	Advantages of heterogeneous catalysts over homogeneous catalysts	1-2
	1.2	Various adsorption parameters for removal of reactive brilliant red X-3B using three different LDH adsorbents	1-16
	1.3	Adsorption efficiencies of various adsorbent in terms of degradation (%) and maximum adsorption capacity, q_m (mg/g) for removal of EBT	1-17
	1.4	Degradation (%) and K_{app} values for photodegradation of MV and MG over various anion intercalated ZnFe LDH photocatalysts	1-20
3	3A.1	Elemental analyses, unit cell parameters and average crystallite sizes of the LDH samples	3-5
	3A.2	Band gap energy and textural properties of LDH samples	3-8
	3A.3	Pseudo-first order and pseudo-second order kinetic parameters for adsorption of MO onto various LDHs	3-15
	3A.4	Langmuir, Freundlich and Dubinin–Radushkevich (D–R) isotherm parameters of MO onto various LDHs	3-17
	3A.5	Comparison of adsorption of MO onto LDH3 with earlier reports	3-23
	3B.1	Chemical composition, unit cell parameters and textural properties of various LDH samples	3-27
	3B.2	Pseudo-first order and second order kinetic parameters for adsorption of different dyes over CuMgAl ₄ LDH	3-38
	3B.3	Langmuir, Freundlich and Dubinin–Radushkevich (D–R) isotherm parameters for adsorption of different dyes over CuMgAl ₄ LDH	3-41

	3C.1	Unit cell parameters and average crystallite sizes of all the samples	3-47
	3C.2	Langmuir, Freundlich and D-R adsorption isotherm parameters for adsorption of CR on NiAl-S ₁ LDH	3-54
	3C.3	Adsorption kinetic parameters for adsorptive removal of CR on NiAl-S ₁ LDH	3-56
	3C.4	Thermodynamic parameters for adsorptive removal of CR over NiAl-S ₁ LDH	3-57
	3C.5	Comparison of adsorption efficiency of NiAl-S ₁ LDH for removal of CR with various reported adsorbents	3-60
4	4.1	The list of various organic pollutants used in the present investigation	4-1–4-2
	4.2	Lattice parameters, crystallite size, band gap energy, BET surface area, pore volume and pore diameter of ZnFe LDH	4-4
	4A.1	% Degradation and pseudo first order kinetic parameters for photocatalytic degradation of phenol, 2CP, 3CP and 4CP over ZnFe LDH under different light irradiations	4-16
	4A.2	Reaction enthalpy ($\Delta_r H^0$) and Gibbs free energy ($\Delta_r G^0$) for the photocatalytic degradation of phenol	4-23
	4A.3	Total energy (Hartree) and relative energies (kcal/mol) of all species involved in the reaction calculated at B3LYP/6-31+G (d,p) levels of theory	4-23
	4B.1	Degradation (%), K_{app} and linear regression value for photodegradation of MB and RhB over ZnFe LDH under UV light irradiation	4-29
	4B.2	Comparison of photocatalytic activity of ZnFe LDH with reported catalysts	4-35
5	5A.1	Chemical analysis, crystallographic data and crystallite size of mixed oxides	5-5

5A.2	Textural properties of LDH derived mixed oxides	5-8
5A.3	Nitroalkylation of 4-nitrobenzaldehyde with nitromethane over various catalysts at room temperature	5-9
5A.4	Effect of temperature on nitro-aldol condensation reaction of 4-nitrobenzaldehyde and nitromethane catalyzed by NiMgAl mixed oxide	5-10
5A.5	Effect of catalyst amount (NiMgAl mixed oxide) on nitro-aldol condensation reaction of 4-nitrobenzaldehyde and nitromethane	5-11
5A.6	Nitro-aldol condensation reaction of nitromethane with different aldehydes using NiMgAl (O) mixed oxide	5-11
5A.7	Effect of calcination temperature on nitro-aldol condensation reaction of 4-nitrobenzaldehyde and nitromethane catalyzed by NiMgAl (O) mixed oxide	5-13
5A.8	Recyclability of NiMgAl (O) mixed oxide	5-15
5B.1	Chemical analyses of the precursor LDHs	5-20
5B.2	Textural properties of various mixed oxides derived from LDHs	5-22
5B.3	Nitro-aldol condensation of 4-nitrobenzaldehyde with nitromethane over various mixed oxides derived from LDHs	5-23
5B.4	Effect of catalyst amount on conversion (%) of nitro-aldol condensation reaction over LDH3 (O)	5-24
5B.5	Effect of MW power (%) on conversion (%) of nitro-aldol condensation reaction over LDH3 (O)	5-25
5B.6	Nitro-aldol condensation reaction of nitromethane with different aldehydes over LDH3 (O)	5-25–5-26