

List of Figures

Chapter	Figure No.	Figure Caption	Page No.
<i>Chapter 1</i>	1.1	Schematic representation of cluster placed in between atom, molecule and bulk material	1-5
	1.2	Examples of various types of clusters	1-6
	1.3	Variation of BE with cluster size for Pd _n (n=1-7) clusters	1-8
	1.4	Variation of IPs of the most stable isomers with N=n+1	1-9
	1.5	Cluster size (N) dependence to the extent of reaction of Au _N ⁻ and O ₂ . Separate plots correspond to different O ₂ concentrations	1-9
	1-6	Dependence of melting temperature (T) on the size of Au nanoparticles	1-10
	1-7	Fourth Century AD <i>Lycurgus cup</i>	1-11
<i>Chapter 2</i>	2.1	Points on potential energy surface	2-4
<i>Chapter 3</i>	3.1.1	Structures of neutral Au _{n+2} and Au _n Be ₂ (n=1-5) clusters	3-6
	3.1.2	Structures of cationic Au _{n+2} and Au _n Be ₂ (n=1-5) clusters	3-7
	3.1.3	Structures of anionic Au _{n+2} and Au _n Be ₂ (n=1-5) clusters	3-8
	3.1.4	Size dependence of the (A) averaged binding energies Eb(n), (B) fragmentation energies ΔE(n) and (C) second-order difference of energies Δ ₂ E(n) and for the lowest energy structures of neutral Au _{n+2} and Au _n Be ₂ (n=1–5) clusters	3-10
	3.1.5	Size dependence of the (A) averaged binding energies Eb(n), (B) fragmentation energies ΔE(n) and (C)	3-11

	second-order difference of energies $\Delta_2E(n)$ and for the lowest energy structures of cationic Au_{n+2} and Au_nBe_2 ($n=1-5$) clusters	
3.1.6	Size dependence of the (A) averaged binding energies $E_b(n)$, (B) fragmentation energies $\Delta E(n)$ and (C) second-order difference of energies $\Delta_2E(n)$ and for the lowest energy structures of anionic Au_{n+2} and Au_nBe_2 ($n=1-5$) clusters	3-12
3.1.7	Comparison of stability of neutral and charged Au_nBe_2 clusters	3-13
3.1.8	Shifting of BCP in Au-Au and Au-Be bonds. The red dots indicate BCP	3-15
3.1.9	Trajectory field in some of the Au_n and Au_nBe_2 clusters. Gold and magnesium atoms are represented by yellow and green spheres respectively. Bond paths and basin paths are indicated by dark green and blue lines, while the interatomic surfaces are indicated by dark blue lines. Red and green dots indicate bond critical points and ring critical points, respectively	3-16
3.1.10	HOMO and LUMO isosurfaces of Au_3 and $AuBe_2$ clusters	3-17
3.2.1	Optimized structures of neutral Au_{n+2} and Au_nMg_2 ($n = 1-5$) clusters	3-21
3.2.2	Optimized structures of cationic Au_{n+2} and Au_nMg_2 ($n = 1-5$) clusters	3-22
3.2.3	Optimized structures of anionic Au_{n+2} and Au_nMg_2 ($n = 1-5$) clusters	3-23
3.2.4	Variation of binding energies with respect to cluster size for (A) neutral, (B) cationic and (C) anionic clusters in bare and magnesium doped gold clusters	3-26
3.2.5	Comparison of binding energy per atom among charge	3-27

	and neutral clusters of Au_nMg_2 clusters	
3.2.6	Variation of fragmentation energies with respect to cluster size for (A) neutral, (B) cationic and (C) anionic clusters in bare and magnesium doped gold clusters	3-28
3.2.7	Variation of second order difference of energies with respect to cluster size for (A) neutral, (B) cationic and (C) anionic clusters in bare and magnesium doped gold clusters	3-29
3.2.8	Variation of VEA values with respect to cluster size for pure and Mg doped Au clusters	3-32
3.2.9	Variation of VIP values with respect to cluster size for pure and Mg doped Au clusters	3-33
3.2.10	Shifting of BCP in Au-Au and Au-Mg bonds. The red dots indicate BCP	3-36
3.2.11	Trajectory field in some of the Au_n and Au_nMg_2 clusters. Gold and magnesium atoms are represented by yellow and green spheres respectively	3-37
3.2.12	HOMO and LUMO isosurfaces of Au_3 and AuMg_2 clusters	3-38
3.3.1	Optimized structures of neutral Au_{n+2} and Au_nAl_2 ($n = 1-5$) clusters	3-43
3.3.2	Optimized structures of cationic Au_{n+2} and Au_nAl_2 ($n = 1-5$) clusters	3-44
3.3.3	Optimized structures of anionic Au_{n+2} and Au_nAl_2 ($n = 1-5$) clusters	3-45
3.3.4	Variation of binding energies with respect to cluster size for (a) neutral, (b) cationic and (c) anionic clusters in bare and Al doped gold clusters	3-47
3.3.5	Comparison of binding energy per atom among charge and neutral clusters of Au_nAl_2 clusters	3-48

	3.3.6	Variation of fragmentation energies with respect to cluster size for (a) neutral, (b) cationic and (c) anionic clusters in bare and aluminium doped gold clusters	3-49
	3.3.7	Variation of second order difference of energies with respect to cluster size for (a) neutral, (b) cationic and (c) anionic clusters in bare and aluminium doped gold clusters	3-51
	3.3.8	Variation of VEA values with respect to cluster size for pure and Al doped Au clusters	3-52
	3.3.9	Variation of VIP values with respect to cluster size for pure and Al doped Au clusters	3-54
	3.3.10	Shifting of BCP in Au-Au and Au-Al bonds. The red dots indicate BCP	3-56
	3.3.11	Trajectory field in some of the Au_n and Au_nAl_2 clusters. Gold and aluminium atoms are represented by yellow and grey spheres respectively	3-58
	3.3.12	HOMO and LUMO isosurfaces of Au_3 and $AuAl_2$ clusters	3-59
Chapter 4	4.1.1	Optimized geometries of reactants, intermediates, transition states and products involved in the oxidative pathways of NO at B3LYP level	4-8
	4.1.2	The HOMO and LUMO isosurfaces for Au_2^- dimer, O_2 , and NO molecules	4-10
	4.1.3	Potential energy profile of the NO oxidation by the Au_2^- dimer. The relative energies (in $kcal\ mol^{-1}$) were calculated with the ZPE corrections at the B3LYP/LANL2DZ level	4-12
	4.2.1	Optimized geometries for complexes between O_2 and NO molecules with Au-Ag- dimers in two different sites. The numbers in parentheses are calculated natural charges of atoms (in e)	4-18

	4.2.2	HOMO and LUMO isosurfaces for Au-Ag ⁻ dimer, O ₂ and NO molecules	4-21
	4.2.3	Optimized geometries of reactants, intermediates, transition states and products involved in the oxidative pathways of NO by Au-Ag ⁻ dimer along Ag and Au sites	4-23
	4.2.4	Relative energy profile for the NO oxidation by Au-Ag ⁻ dimer along Ag site	4-24
	4.2.5	Relative energy profile for the NO oxidation by Au-Ag ⁻ dimer along Au site	4-26
Chapter 5	5.1.1	Optimized geometries of reactants and transition states at M06-2X/6-31+G(d,p) level. Bond lengths are in Angstroms	5-7,5-8
	5.1.2	Optimized geometries of reactant complexes, product complexes and products at M06-2X/6-31+G(d,p) level.	5-9
	5.1.3	Potential energy profile for the CF ₃ CF ₂ CH ₂ OCH ₃ + OH reactions. Relative energies (in kcal mol ⁻¹) at M06-2X/6-31+G(d,p) level	5-12
	5.2.1	Optimized geometries of reactants and transition states at M06-2X/6-31+G(d,p) level of theory. Bond lengths are in Angstroms	5-25
	5.2.2	Optimized geometries of reactant complexes, product complexes and products at M06-2X/6-31+G(d,p) level of theory	5-26
	5.2.3	Schematic potential energy diagram for i-C ₃ F ₇ OCH ₃ + OH reactions. Relative energies (in kcal mol ⁻¹) with ZPE at M06-2X/6-311++G(d,p) level. The values given in parentheses are calculated at M06-2X/6-31+G(d,p) level of theory	5-29
	5.2.4	Rate constants for hydrogen abstraction reactions of i-C ₃ F ₇ OCH ₃ with OH radicals and total rate constant	5-33

(k_{OH}) for $i\text{-C}_3\text{F}_7\text{OCH}_3 + \text{OH}$ reactions

- 5.2.5 Rate constants for hydrogen abstraction reactions of $i\text{-C}_3\text{F}_7\text{OC(O)H} + \text{OH/Cl}$ reactions with experimental data 5-37