LIST OF FIGURES

Figure 1.1: Formation of fibrin from fibrinogen
Figure 1.2: Structure of Nattokinase
Figure 1.3: Structure of streptokinase
Figure 1.4: Structure of Urokinase
Figure 1.5: Structure of tissue plasminogen activator (t-PA)
Figure 1.6: Schematic representation of plasmin formation through <i>in vivo</i> processing
Figure 1.7: Schematic representation of fibrinolysis by plasmin
Figure 3.1: Intravenous administration of purified fibrin(ogen)olytic/ fibrinolytic enzyme at a dose of 10 mg/kg in tail vein rats
Figure 3.2: Echocardiogram of control as well as treated rats after 14 days of intravenous administration of purified fibrin(ogen)olytic/ fibrinolytic enzyme/ PBS (control) at a dose of 10 mg/kg in tail vein rats
Figure 3.3: Measurement of the PenH value (respiratory system) by whole body plethysmograph after 14 days of intravenous administration of purified fibrin(ogen)olytic/ fibrinolytic enzyme or PBS (control) at a dose of 10 mg/kg in tail vein rat
Figure 3.4: Collection of blood sample from control and treated rats by mechanical disruption of retro orbital sinus
Figure 3.5: Collection of blood sample from control and treated rats for hematological (blue) and serum profile analysis (red)
Figure 3.6: BIOPAC system was used to capture rat before intravenous administration of purified enzymes in tail vein
Figure 3.7: Intravenous administration of purified enzyme in carrageenan induced rat tail vein
Figure 4.1: Zone of hydrolysis around the FF01 bacterial colonies on fibrin agar medium

Figure 4.2: Pure culture	morphology and	gram staining of	bacterial strains	FF01.9	3

- Figure 4.7: MALDI-TOF-MS spectrum of Bacethrombase. 102

- Figure 4.11: Effect of Bacethrombase on extracellular membrane proteins 108
- Figure 4.12: Effect of temperature on activity of Bacethrombase...... 109
- Figure 4.13: Effect of temperature on Bacethrombase activity 109
- Figure 4.14: Effect of substrate (fibrinogen) concentration on protease activity of Bacethrombase at 2.5 µM to 58.5 µM.
- Figure 4.15: Plasminogen activation property of Bacethrombase 112
- Figure 4.17A: A SDS-PAGE analysis of fibrinogen degradation by Bacethrombase

Figure 4.17B: SDS-PAGE analysis of fibrin degradation by Bacethrombase...... 115

- Figure 4.20: Cell cytotoxicity of Bacethrombase on Hela and HT29 cells 119
- Figure 4.21A: A Dose-dependent platelet deaggregation by Bacethrombase 120
- Figure 4.21B: Effect of Bacethrombase on collagen-induced platelet aggregation 121
- Figure 4.22: Effect of pre-incubation time (0 30 min) of Bacethrombase (0.1 μ M) / platelet poor plasma (PPP) at 37 °C on Ca²⁺cloting time of PPP 123

- Figure 4.27: Effect of i.v. injection of Bacethrombase on ECG of rats...... 129

Figure 4.29: Determination of carrageenan-induced thrombus formation in rat tail
Figure 4.30: (a-f) Dose-dependent thrombolytic effect of Bacethrombase, streptokinase and plasmin
Figure 4.31: Effect of different concentrations of Bacethrombase on activated thromboplastin time (APTT) and prothrombin time (PT) of PPP 136
Figure 4.32: Assessment of thrombin time (TT) at various dosages of Bacethrombase
Figure 4.33: The <i>in vivo</i> dose-dependent defibrinogenating activity of Bacethrombase
Figure 4.34: Dose dependent <i>in vivo</i> anticoagulant action of Bacethrombase 138
Figure 5.1: Zone of hydrolysis around the FF02B bacterial colonies on fibrin plate agar medium
Figure 5.2: Pure culture of colony morphology, Gram staining of bacterial strain FF02B
Figure 5.3: Phylogenetic relationships of strain FF02B and other closely related <i>Brevibacillus</i> species based on partial 16S rDNA sequences
Figure 5.4: Phylogenetic relationships of strain FF02B and other closely related <i>Brevibacillus</i> species based on 16S-23S ISR homologous sequences 153
Figure 5.5A: Purification of crude Brevithrombolase from <i>Brevibacillus brevis</i> strain FF02B by Hydrophobic interaction chromatography
Figure 5.5B: Purification of Brevithrombolase from <i>Brevibacillus brevis</i> strain FF02B by Gel-filtration chromatography
Figure 5.6: Determination of purity and molecular mass of Brevithrombolase by 12.5% SDS-PAGE analysis under reduced and non-reduced conditions.
Figure 5.7: MALDI-TOF-MS spectrum of Brevithrombolase
Figure 5.8: Determination of secondary structure of native Brevithrombolase by circular dichroism (CD) analysis

Figure 5.9: Determination of Km and Vmax of Brevithrombolase by Lineweaver-
Burk double reciprocal plot
Figure 5.10: Effect of temperature on Brevithrombolase activity. Values are mean of \pm SD of three replicates
Figure 5.11: Effect of pH on Brevithrombolase activity. Values are mean of ± SD of three replicates
Figure 5.12: Plasminogen activation property of Brevithrombolase 162
Figure 5.13: 15% SDS-PAGE (reducing conditions) analysis of native and deglycosylated Brevithrombolase
Figure 5.14: A comparison of fibrin degradation between 1.0 µM of (A) Brevithrombolase, and (B) plasmin:
Figure 5.15: A comparison of fibrinolytic activity between plasmin and Brevithrombolase under identical conditions (37 °C, pH 7.4) 166
Figure 5.16: Comparison of RP-UPLC elution profile of fibrin degradation products, generated by plasmin / Brevithrombolase, under identical experimental conditions
Figure 5.17: Cell cytotoxicity of Brevithrombolase on Hela and HT29 cells 168
Figure 5.18: A comparison of the dose-dependent anticoagulant activity among Brevithrombolase, warfarin, heparin and plasmin under identical experimental conditions
Figure 5.19: Effect of pre-incubation (0–40 min) of Brevithrombolase/platelet poor plasma (PPP) on Ca ²⁺ cloting time of PPP
Figure 5.20: Effect of Brevithrombolase in FXa inhibition as analyzed by 12.5% SDS-PAGE
Figure 5.21: Dose-dependent inhibition of amidolytic activity of thrombin by Brevithrombolase
Figure 5.22: Effect of Brevithrombolase (0.05 -1 µM) on fibrinogen clotting time of thrombin
Figure 5.24: Kinetics of thrombin degradation by Brevithrombolase at 37 °C 174

Figure 5.25: Effect of <i>i.v</i> injection of Brevithrombolase (10mg/kg) on ECG of
treated rat 177
Figure 5.26(A-H): Light microscopic observation of the effect of Brevithrombolase
(at a dose of 10 mg/kg) on different organs of treated rats 179
Figure 5.27: The in vivo effect of Brevithrombolase on tail bleeding and blood
clotting time in Wistar strain albino rats
Figure 5.28: Dose-dependent thrombolytic effect of Brevithrombolase 180
Figure 5.29: Dose dependent activated partial thromboplastin time (APTT) and
prothrombin time (PT) of Brevithrombolase 182
Figure 5.30: Dose-dependent thrombin time (TT) of Brevithrombolase-treated rats.
Figure 5.31: The in vivo dose-dependent fibrin(ogen)olytic activity of
Brevithrombolase
Figure 5.32: Dose-dependent <i>in vivo</i> anticoagulant activity of Brevithrombolase185