

Bibliography

- [1] Fry, B.G., & Wu, W. Assembling an Arsenal: Origin and Evolution of the Snake Venom Proteome Inferred from Phylogenetic Analysis of Toxin Sequences, *Mol Biol Evol.* **21** (5), 870–883, 2004.
- [2] Mackessy, S.P., & Baxter, L.M. Bioweapons synthesis and storage: The venom gland of front-fanged snakes, *Zool Anz.* **245** (1&2), 147–159, 2006.
- [3] Mackessy, S.P. Biochemistry and Pharmacology of Colubrid Snake Venoms, *J. Toxicol.* **21** (1&2), 43–83, 2002.
- [4] Sajevic, T. et al. Haemostatically active proteins in snake venoms, *Toxicon* **57** (5), 627–645, 2011.
- [5] Birrell, G.W. et al. The diversity of bioactive proteins in Australian snake venoms, *Mol Cell Proteomics.* **6** (6), 973–986, 2007.
- [6] Berling, I., & Isbister, G.K. Hematologic Effects and Complications of Snake Envenoming, *Transfus Med Rev* **29** (2), 82–89, 2015.
- [7] Ferreira, S.H. et al. Isolation of bradykinin-potentiating peptides from *Bothrops jararaca* venom, *Biochemistry* **9** (13), 2583–2593, 1970.
- [8] Cushman, D.W., & Ondetti, M.A. History of the design of captopril and related inhibitors of angiotensin converting enzyme, *Hypertension* **17** (4), 589–592, 1991.
- [9] BL, D., & PR., S. Snake venom derived molecules in tumor angiogenesis and its application in cancer therapy; an overview, *Curr Top Med Chem.* **15** (7), 649–657, 2015.
- [10] Wisner, A. et al. Snake venom proteinases as tools in hemostasis studies: structure-function relationship of a plasminogen activator purified from *Trimeresurus stejnegeri* venom, *Haemostasis* **31** (3-6), 133–140, 2001.
- [11] Andrews, R.K. et al. The use of snake venom toxins as tools to study platelet receptors for collagen and von Willebrand factor, *Haemostasis* **31** (3-6), 155–172, 2001.
- [12] Marsh, N.A. Diagnostic uses of snake venom, *Haemostasis* **31** (3-6), 211–217, 2001.
- [13] Volkers, N. Venom anyone? Researchers tweak “bad” compounds into good medicines, *J Natl Cancer Inst.* **91** (8), 667–668, 1999.
- [14] Kasturiratne, A. et al. The global burden of snakebite: A literature analysis and modelling based on regional estimates of envenoming and deaths, *PLoS Med.* **5** (11), e1604, 2008.

- [15] Khandelwal, G. et al. *Naja kaouthia*: Two cases of Asiatic cobra envenomations, *J Emerg Med* **32** (2), 171–174, 2007.
- [16] Wüster, W. Taxonomic changes and toxinology: Systematic revisions of the asiatic cobras (*Naja naja* species complex), *Toxicon* **34** (4), 399–406, 1996.
- [17] Moore, C.B. America's mythical snakes, *The Scientific monthly* **68** (1), 52–58, 1949.
- [18] Jakes, L. *Mythic Creatures—the Snake*, 2011.
- [19] Rieppel, O., & Kearney, M. The origin of snakes: Limits of a scientific debate, *Biologist* **48** (3), 110–114, 2001.
- [20] Jonniaux, P., & Kumazawa, Y. Molecular phylogenetic and dating analyses using mitochondrial DNA sequences of eyelid geckos (Squamata: Eublepharidae), *Gene* **407** (1-2), 105–115, 2008.
- [21] Siler, C.D., & Brown, R.M. Evidence for repeated acquisition and loss of complex body-form characters in an insular clade of southeast Asian semi-fossorial skinks, *Evolution* **65** (9), 2641–2663, 2011.
- [22] Caldwell, M.W. et al. The oldest known snakes from the Middle Jurassic-Lower Cretaceous provide insights on snake evolution, *Nature commun.* **6**, 5996, 2015.
- [23] Rauhut, O.W.M. et al. A New Rhynchocephalian from the Late Jurassic of Germany with a Dentition That Is Unique amongst Tetrapods, *PLoS ONE* **7** (10), 46893–46908, 2012.
- [24] Evans, S.E. At the feet of the dinosaurs: the early history and radiation of lizards, *Biol Rev Camb Philos Soc.* **78** (4), 513–551, 2003.
- [25] Spinner, M. et al. Snake velvet black: hierarchical micro- and nanostructure enhances dark colouration in *Bitis rhinoceros*, *Sci Rep.* **3**, 1833–1846, 2013.
- [26] Spinner, M. et al. Non-contaminating camouflage: Multifunctional skin microornamentation in the West African Gaboon viper (*Bitis rhinoceros*), *PLoS ONE* **9** (3), 91087–91099, 2014.
- [27] Alibardi, L. Differentiation of snake epidermis, with emphasis on the shedding layer, *J Morphol.* **264** (2), 178–190, 2005.
- [28] Close, M., & Cundall, D. Snake lower jaw skin: Extension and recovery of a hyperextensible keratinized integument, *J Exp Zool A Ecol Genet Physiol.* **321** (2), 78–97, 2014.
- [29] Chen, Q. et al. Reduced performance of prey targeting in pit vipers with contralaterally occluded infrared and visual senses, *PLoS ONE* **7** (5), 34980–34989, 2012.

- [30] Bolívar-G, W. et al. Discovery of a novel accessory structure of the pitviper infrared receptor organ (Serpentes: Viperidae), *PLoS ONE* **9** (3), 90609–90622, 2014.
- [31] Vilela, B. et al. Body Size, Extinction Risk and Knowledge Bias in New World Snakes, *PLoS ONE* **9** (11), 113418–113429, 2014.
- [32] Rial, R. V. et al. Evolution of wakefulness, sleep and hibernation: From reptiles to mammals, *Neurosci Biobehav Rev.* **34** (8), 1144–1160, 2010.
- [33] Sivan, J. et al. Temporal activity and dietary selection in two coexisting desert snakes, the Saharan sand viper (*Cerastes vipera*) and the crowned leafnose (*Lytorhynchus diadema*), *Zoology* **116** (2), 113–117, 2013.
- [34] Rugiero, L. et al. Climate-change-related shifts in annual phenology of a temperate snake during the last 20 years, *Acta Oecol.* **51**, 42–48, 2013.
- [35] Huang, S. et al. What are the closest relatives of the hot-spring snakes (Colubridae, Thermophis), the relict species endemic to the Tibetan Plateau?, *Mol Phylogenet Evol.* **51** (3), 438–446, 2009.
- [36] Lourdais, O. et al. Cold climate specialization: Adaptive covariation between metabolic rate and thermoregulation in pregnant vipers, *Physiol Behav.* **119**, 149–155, 2013.
- [37] Tan, T.L. et al. Bitten by the “flying” tree snake, *Chrysopelea paradisi*, *J Emerg Med.* **42** (4), 420–423, 2012.
- [38] Boyd, J.J. et al. Venomous snakebite in mountainous terrain: prevention and management, *Wilderness Environ Med.* **18** (3), 190–202, 2007.
- [39] Lane, A., & Shine, R. Phylogenetic relationships within laticaudine sea snakes (Elapidae), *Mol Phylogenet Evol.* **59** (3), 567–577, 2011.
- [40] Brooks, S.E. et al. Vulnerability of Cambodian water snakes: Initial assessment of the impact of hunting at Tonle Sap Lake, *Biol Conserv.* **139**, 401–414, 2007.
- [41] Ismail, M., & Memish, Z. a. Venomous snakes of Saudi Arabia and the Middle East: A keynote for travellers, *Int J Antimicrob Agents.* **21** (2), 164–169, 2003.
- [42] Uetz, P., & Jirí, H. Reptile database, 2014.
- [43] Steve Csurhes and Paul Fisher. *Pest animal risk assesment, Cobra (all species)*, 2010.
- [44] Wüster, W., & Thorpe, R.S. Asiatic cobras: systematics and snakebite., *Experientia* **47** (2), 205–209, 1991.

- [45] Ahmed, F.A. et al. *Amphibians and Reptiles of North East India; A photographic guide*, Guwahati, India, 2009.
- [46] Sinha, K. Snake bites kill 46K in India yearly, *The Times of India* 2011.
- [47] Mebs, D. *Venomous and Poisonous Animals: A Handbook for Biologists, Toxicologists and Toxinologists, Physicians and Pharmacists*, CRC Press, Boca Raton, FL, 2002.
- [48] Drabeck, D.H. et al. Why the honey badger don't care: Convergent evolution of venom-targeted nicotinic acetylcholine receptors in mammals that survive venomous snake bites, *Toxicon* **99**, 68–72, 2015.
- [49] Fry, B.G. et al. Evolution of an arsenal: structural and functional diversification of the venom system in the advanced snakes (*Caenophidia*), *Mol Cell Proteomics*. **7** (2), 215–46, 2008.
- [50] Jouiaei, M. et al. Evolution of an ancient venom: recognition of a novel family of cnidarian toxins and the common evolutionary origin of sodium and potassium neurotoxins in sea anemone, *Mol Biol Evol.* **32** (6), 1598–1608, 2015.
- [51] Fry, B.G. et al. Isolation of a neurotoxin (alpha-colubritoxin) from a nonvenomous colubrid: evidence for early origin of venom in snakes, *J Mol Evol.* **57** (4), 446–52, 2003.
- [52] Thomas, F. et al. Kelps feature systemic defense responses: insights into the evolution of innate immunity in multicellular eukaryotes, *New Phytol.* **204** (3), 567–576, 2014.
- [53] Weinstein, S.A. et al. *Handbook of Venoms and Toxins of Reptiles*, CRC Press, New york, (US states), 2010.
- [54] Kardong, K. v., & Lavln-Murcio, P.A. Venom delivery of snakes as high-pressure and low-pressure systems, *Copeia* **1993** (3), 644–50, 1993.
- [55] Kochva, E. Atractaspis (Serpentes, Atractaspididae) the burrowing asp; a multidisciplinary minireview, *Bull. Nat. Hist. Mus. Lond.* **68** (28), 91–99, 2002.
- [56] Young, B.A. et al. The buccal buckle: the functional morphology of venom spitting in cobras, *J Exp Biol.* **207** (20), 3483–3494, 2004.
- [57] Rosenberg, H. 1. Histology, histochemistry, and emptying mechanism of the venom glands of some elapid snakes, *J. Morphol.* **123** (2), 133–56, 1967.
- [58] Young, B.A. et al. Functional subdivision of the venom gland musculature and the regulation of venom expulsion in rattlesnakes, *J Morphol.* **246** (3), 249–259, 2000.

- [59] Kuch, U. et al. Snake fangs from the Lower Miocene of Germany: Evolutionary stability of perfect weapons, *Naturwissenschaften* **93** (2), 84–87, 2006.
- [60] Casewell, N.R. et al. Dynamic evolution of venom proteins in squamate reptiles, *Nat Commun.* **3**, 1066, 2012.
- [61] Fry, B.G. et al. The toxicogenomic multiverse: convergent recruitment of proteins into animal venoms, *Ann. rev. genom. human gen.* **10**, 483–511, 2009.
- [62] Liu, S. et al. “Anatomical” View of the Protein Composition and Protein Characteristics for Gloydius shedaoensis Snake Venom via Proteomics Approach, *Anat Rec (Hoboken)*. **294** (2), 273–282, 2011.
- [63] Hider, R.C. et al. *Separation and purification of toxins from snake venoms*. RC, Pergamon Press, New york, (US states), 1991.
- [64] Calvete, J.J. Snake venomomics: From the inventory of toxins to biology, *Toxicon* **75**, 44–62, 2013.
- [65] Koh, D.C.I. et al. Snake venom components and their applications in biomedicine, *Cell. Mol. Life Sci.* **63** (24), 3030–3041, 2006.
- [66] Vejayan, J. et al. Comparative analysis of the venom proteome of four important Malaysian snake species, *J. Venom. Anim. Toxins incl. Trop. Dis.* **20** (1), 6, 2014.
- [67] Laustsen, A.H. et al. Snake venomomics of monocled cobra (*Naja kaouthia*) and investigation of human IgG response against venom toxins, *Toxicon* **99**, 23–35, 2015.
- [68] Tan, K.Y. et al. Venomomics, lethality and neutralization of *Naja kaouthia* (monocled cobra) venoms from three different geographical regions of Southeast Asia, *J Proteomics.* **120** (2), 105–125, 2015.
- [69] Pradniwat, P., & Rojnuckarin, P. The structure-function relationship of thrombin-like enzymes from the green pit viper (*Trimeresurus albolabris*), *Toxicon* **100**, 53–59, 2015.
- [70] Rajesh, R. V. et al. High aryl acylamidase activity associated with cobra venom acetylcholinesterase: Biological significance, *Biochimie* **91** (11-12), 1450–1456, 2009.
- [71] Nanjaraj Urs, A.N. et al. Local and systemic toxicity of *Echis carinatus* venom: neutralization by *Cassia auriculata* L. leaf methanol extract, *J Nat Med.* **69** (1), 111–122, 2015.
- [72] Costa, T.R. et al. Snake venom L-amino acid oxidases: an overview on their antitumor effects, *J Venom Anim Toxins Incl Trop Dis.* **20**, 23, 2014.

- [73] Bordon, K.C.F. et al. Bordonein-L, a new L-amino acid oxidase from *Crotalus durissus terrificus* snake venom: isolation, preliminary characterization and enzyme stability, *J Venom Anim Toxins Incl Trop Dis.* **21**, 26, 2015.
- [74] Kini, R.M. Excitement ahead: Structure, function and mechanism of snake venom phospholipase A2 enzymes, *Toxicon* **42** (8), 827–840, 2003.
- [75] Jridi, I. et al. Hemilipin, a novel *Hemiscorpius lepturus* venom heterodimeric phospholipase A2, which inhibits angiogenesis in vitro and in vivo, *Toxicon* **105**, 34–44, 2015.
- [76] Herrera, C. et al. Tissue localization and extracellular matrix degradation by PI, PII and PIII snake venom metalloproteinases: clues on the mechanisms of venom-induced hemorrhage, *PLoS Negl Trop Dis.* **9** (4), e0003731, 2015.
- [77] Oyama, E., & Takahashi, H. Purification and characterization of two high molecular mass snake venom metalloproteinases (P-III SVMPs), named SV-PAD-2 and HR-Ele-1, from the venom of *Protobothrops elegans* (Sakishimababu), *Toxicon* **103**, 30–8, 2015.
- [78] Osipov, A. V. et al. Cobra Venom Contains a Protein of the CRISP Family, *Russian J Bioorganic Chem.* **27**, 198–199, 2001.
- [79] Yamazaki, Y. et al. Wide distribution of cysteine-rich secretory proteins in snake venoms: Isolation and cloning of novel snake venom cysteine-rich secretory proteins, *Arch. Biochem. Biophys.* **412** (1), 133–141, 2003.
- [80] Mukherjee, A.K., & Mackessy, S.P. Pharmacological properties and pathophysiological significance of a Kunitz-type protease inhibitor (Rusvikunin-II) and its protein complex (Rusvikunin complex) purified from *Daboia russelii russelii* venom, *Toxicon* **89**, 55–66, 2014.
- [81] Trummal, K. et al. Molecular diversity of snake venom nerve growth factors, *Toxicon* **58** (4), 363–368, 2011.
- [82] Sartim, M.A. et al. Galatrox is a C-type lectin in *Bothrops atrox* snake venom that selectively binds LacNAc-terminated glycans and can induce acute inflammation, *Glycobiology* **24** (11), 1010–1021, 2014.
- [83] Arruda Macêdo, J.K. et al. Disintegrins from snake venoms and their applications in cancer research and therapy, *Curr Protein Pept Sci.* **16** (6), 532–548, 2015.
- [84] Jackson, T.N.W. et al. Venom down under: Dynamic evolution of Australian elapid snake toxins, *Toxins* **5** (12), 2621–2655, 2013.
- [85] Kini, R.M., & Doley, R. Structure, function and evolution of three-finger toxins: Mini proteins with multiple targets, *Toxicon* **56** (6), 855–867, 2010.

- [86] Nirthanan, S. et al. Non-conventional toxins from Elapid venoms, *Toxicon* **41** (4), 397–407, 2003.
- [87] Mukherjee, A.K. et al. Two acidic, anticoagulant PLA2 isoenzymes purified from the venom of monocled cobra *Naja kaouthia* exhibit different potency to inhibit thrombin and factor Xa via phospholipids independent, non-enzymatic mechanism, *PLoS ONE* **9** (8), e101334, 2014.
- [88] Doley, R., & Mukherjee, A.K. Purification and characterization of an anticoagulant phospholipase A2 from Indian monocled cobra (*Naja kaouthia*) venom, *Toxicon* **41** (1), 81–91, 2003.
- [89] Kumar, J.R. et al. Biochemical and pharmacological characterization of three toxic phospholipase A2s from *Daboia russelii* snake venom, *Comp Biochem Physiol C Toxicol Pharmacol* **168** (1), 28–38, 2015.
- [90] Doley, R., & Kini, R.M. Protein complexes in snake venom, *Cell. Mol. Life Sci.* **66** (17), 2851–2871, 2009.
- [91] Schaloske, R.H., & Dennis, E. a. The phospholipase A2 superfamily and its group numbering system, *Biochim Biophys Acta.* **1761** (11), 1246–1259, 2006.
- [92] Gutiérrez, J.M., & Lomonte, B. Phospholipases A2: Unveiling the secrets of a functionally versatile group of snake venom toxins, *Toxicon* **62**, 27–39, 2013.
- [93] Debashree, S. et al. Mechanism of in vivo anticoagulant and haemolytic activity by a neutral phospholipase A(2) purified from *Daboia russelii russelii* venom: correlation with clinical manifestations in Russell's Viper envenomed patients, *Toxicon* **15** (76), 291–300, 2013.
- [94] Chaisakul, J. et al. Hypotensive and vascular relaxant effects of phospholipase A2 toxins from Papuan taipan (*Oxyuranus scutellatus*) venom, *Euro J Pharmacol.* **723**, 227–233, 2014.
- [95] Kini, M. Structure-Function Relationships of Phospholipases, *J Biol Chem* **262** (30), 14402–14407, 1987.
- [96] Doley, R. et al. Differential hydrolysis of erythrocyte and mitochondrial membrane phospholipids by two phospholipase A2 isoenzymes (NK-PLA2-I and NK-PLA2-II) from the venom of the Indian monocled cobra *Naja kaouthia*., *Archives of biochemistry and biophysics* **425** (1), 1–13, 2004.
- [97] Mukherjee, A.K. Correlation between the phospholipids domains of the target cell membrane and the extent of *Naja kaouthia* PLA(2)-induced membrane damage: evidence of distinct catalytic and cytotoxic sites in PLA(2) molecules, *Biochim Biophys Acta.* **1770** (2), 187–195, 2007.
- [98] Izidoro, L.F.M. et al. Snake venom L-amino acid oxidases: Trends in pharmacology and biochemistry, *Biomed Res Int.* 2014.

- [99] Sakurai, Y. et al. Inhibition of human platelet aggregation by L-amino acid oxidase purified from *Naja naja kaouthia* venom, *Toxicon* **39** (12), 1827–1833, 2001.
- [100] Tan, N.H., & Swaminathan, S. Purification and properties of the L-amino acid oxidase from monocellate cobra (*Naja naja kaouthia*) venom, *Int J Biochem.* **24** (6), 967–973, 1992.
- [101] Markland, F.S., & Swenson, S. Snake venom metalloproteinases, *Toxicon* **62**, 3–18, 2013.
- [102] Jiang, Y. et al. Venom gland transcriptomes of two elapid snakes (*Bungarus multicinctus* and *Naja atra*) and evolution of toxin genes, *BMC Genomics* **12** (1), 2011.
- [103] Matsui, T. et al. Snake venom proteases affecting hemostasis and thrombosis, *Biochim Biophys Acta.* **1477** (1-2), 146–156, 2000.
- [104] Ferrer, V.P. et al. A Novel Hyaluronidase from Brown Spider (*Loxosceles intermedia*) Venom (Dietrich's Hyaluronidase): From Cloning to Functional Characterization, *PLoS Negl Trop Dis.* **7** (5), 2013.
- [105] Harrison, R.A. et al. Identification of cDNAs encoding viper venom hyaluronidases: Cross-generic sequence conservation of full-length and unusually short variant transcripts, *Gene* **392** (1-2), 22–33, 2007.
- [106] Kemparaju, K., & Girish, K.S. Snake venom hyaluronidase: A therapeutic target, *Cell Biochemi. Funct.* **24** (1), 7–12, 2006.
- [107] Rusmili, M.R.A. et al. Proteomic characterization and comparison of Malaysian *Bungarus candidus* and *Bungarus fasciatus* venoms, *J Proteomics.* **10**, 122, 2014.
- [108] Tan, C.H. et al. Venom-gland transcriptome and venom proteome of the Malaysian king cobra (*Ophiophagus hannah*), *BMC Genomics.* **16** (1), 687, 2015.
- [109] Robaire, B., & Kato, G. Acetylcholinesterase and alpha-bungarotoxin: a study of their possible interaction, *FEBS lett.* **46** (1), 218–223, 1974.
- [110] Bourne, Y. et al. Crystal structure of snake venom acetylcholinesterase in complex with inhibitory antibody fragment Fab410 bound at the peripheral site: evidence for open and closed states of a back door channel, *J Biol Chem.* **290** (3), 1522–35, 2015.
- [111] Sunagar, K. et al. Molecular evolution of vertebrate neurotrophins: Co-option of the highly conserved nerve growth factor gene into the advanced snake venom arsenal, *PLoS ONE* **8** (11), e81827, 2013.

- [112] Wijeyewickrema, L.C. et al. Nerve growth factor inhibits metalloproteinase-disintegrins and blocks ectodomain shedding of platelet glycoprotein VI, *J Biol Chem.* **285** (16), 11793–11799, 2010.
- [113] Osipov, A. V. et al. Cobra venom contains a pool of cysteine-rich secretory proteins, *Biochem Biophys Res Commun.* **328** (1), 177–182, 2005.
- [114] Wang, Y.L. et al. Cobra CRISP functions as an inflammatory modulator via a novel Zn²⁺ and heparan sulfate-dependent transcriptional regulation of endothelial cell adhesion molecules, *J Biol Chem.* **285** (48), 37872–37883, 2010.
- [115] Lecht, S. et al. Anti-angiogenic activities of snake venom CRISP isolated from *Echis carinatus*, *Biochim Biophys Acta.* **1850** (6), 1169–1179, 2015.
- [116] Fukuhara, H. et al. New binding face of C-type lectin-like domains, *Structure* **22** (12), 1694–6, 2014.
- [117] Klein, R.C. et al. A C-type lectin from *Bothrops jararacussu* venom disrupts Staphylococcal biofilms, *PLoS ONE* **10** (3), e0120514, 2015.
- [118] Mukherjee, A.K. et al. A new C-type lectin (RVsnaclec) purified from venom of *Daboia russelii russelii* shows anticoagulant activity via inhibition of FXa and concentration-dependent differential response to platelets in a Ca²⁺-independent manner, *Thromb Res.* **134** (5), 1150–6, 2014.
- [119] García-Fernández, R. et al. Three-dimensional Structure of a Kunitz-type Inhibitor in Complex with an Elastase-like Enzyme, *J Biol Chem.* **290** (22), 14154–65, 2015.
- [120] Calvete, J.J. et al. Snake venom disintegrins: novel dimeric disintegrins and structural diversification by disulphide bond engineering, *Biochem J.* **372** (Pt 3), 725–734, 2003.
- [121] Okuda, D. et al. A new gene structure of the disintegrin family: A subunit of dimeric disintegrin has a short coding region, *Biochemistry* **41** (48), 14248–14254, 2002.
- [122] Moiseeva, N. et al. Structure of acostatin, a dimeric disintegrin from Southern copperhead (*Agkistrodon contortrix contortrix*), at 1.7 Å resolution, *Acta Crystallogr D Biol Crystallogr.* **64** (Pt 4), 466–470, 2008.
- [123] Ruoslahti, E., & Pierschbacher, M.D. Arg-Gly-Asp: a versatile cell recognition signal, *Cell* **44** (4), 517–518, 1986.
- [124] McLane, M.A. et al. New insights on disintegrin-receptor interactions: eristostatin and melanoma cells, *Haemostasis* **31** (3-6), 177–182, 2001.
- [125] McLane, M.A. et al. Disintegrins in health and disease, *Front Biosci.* **13**, 6617–6637, 2008.

- [126] Da Silva, M. et al. Anti-platelet effect of cumanastatin 1, a disintegrin isolated from venom of South American *Crotalus rattlesnake*, *Thromb Res.* **123** (5), 731–739, 2009.
- [127] Sánchez, E.E. et al. Colombistatin: A disintegrin isolated from the venom of the South American snake (*bothrops colombiensis*) that effectively inhibits platelet aggregation and SK-Mel-28 cell adhesion, *Arch Toxicol.* **83** (3), 271–279, 2009.
- [128] Ito, M. et al. Complete amino acid sequence of kaouthiagin, a novel cobra venom metalloproteinase with two disintegrin-like sequences, *Biochemistry* **40** (14), 4503–11, 2001.
- [129] McGivern, J.J. et al. RNA-seq and high-definition mass spectrometry reveal the complex and divergent venoms of two rear-fanged colubrid snakes, *BMC Genomics* **15**, 1061, 2014.
- [130] Pawlak, J., & Kini, R.M. Unique gene organization of colubrid three-finger toxins : Complete cDNA and gene sequences of denmotoxin , a bird-specific toxin from colubrid snake *Boiga dendrophila* (Mangrove Catsnake), *Biochimie* **90** (6), 868–877, 2008.
- [131] Doley, R. et al. Accelerated exchange of exon segments in Viperid three-finger toxin genes (*Sistrurus catenatus edwardsii*; Desert Massasauga), *BMC Evol Biol.* **8**, 196, 2008.
- [132] Girish, V.M. et al. Identification and Structural Characterization of a New Three-Finger Toxin Hemachatoxin from *Hemachatus haemachatus* Venom, *PLoS ONE* **7** (10), e48112, 2012.
- [133] Chang, C.C., & Lee, C.Y. Isolation of neurotoxin from the venom of *Bungarus Multicinctus* and their modes of neuromuscular blocking action, *Arch. Int. Pharmacodyn. Ther.* **144**, 241–257, 1963.
- [134] Heyborne, W.H., & Mackessy, S.P. Identification and characterization of a taxon-specific three-finger toxin from the venom of the Green Vinesnake (*Oxybelis fulgidus*; Family *Colubridae*), *Biochimie* **95** (10), 1923–1932, 2013.
- [135] Nirthanan, S. et al. Neuromuscular effects of candoxin , a novel toxin from the venom of the Malayan krait (*Bungarus candidus*), *Br. J. Pharmacol.* **139** (4), 832–844, 2003.
- [136] Pawlak, J. et al. Denmotoxin, a three-finger toxin from the colubrid snake *Boiga dendrophila* (mangrove catsnake) with bird-specific activity, *J Biol Chem.* **281** (39), 29030–29041, 2006.
- [137] Rajagopalan, N. et al. Beta-cardiotoxin: a new three-finger toxin from *Ophiophagus hannah* (king cobra) venom with beta-blocker activity, *FASEB J.* **21** (13), 3685–3695, 2007.

- [138] Banerjee, Y. et al. Biophysical Characterization of Anticoagulant Hemextin AB Complex from the Venom of Snake *Hemachatus haemachatus*, *Biophys J.* **93** (11), 3963–3976, 2007.
- [139] Roy, A. et al. Structural and functional characterization of a novel homodimeric three-finger neurotoxin from the venom of *Ophiophagus hannah* (King cobra), *J Biol Chem.* **285** (11), 8302–8315, 2010.
- [140] Pawlak, J. et al. Irditoxin, a novel covalently linked heterodimeric three-finger toxin with high taxon-specific neurotoxicity, *FASEB J.* **23** (2), 534–545, 2009.
- [141] Chanda, C. et al. Anti-platelet activity of a three-finger toxin (3FTx) from Indian monocled cobra (*Naja kaouthia*) venom, *Biochem Biophys Res Commun.* **441** (3), 550–554, 2013.
- [142] Tsetlin, V. Snake venom alpha-neurotoxins and other “three-finger” proteins, *FEBS J.* **264** (2), 281–286, 1999.
- [143] Dryer, S.E., & Chiappinelli, V.A. Kappa-bungarotoxin: an intracellular study demonstrating blockade of neuronal nicotinic receptors by a snake neurotoxin, *Brain res.* **289** (1-2), 317–321, 1983.
- [144] Jolkkonen, M. et al. Muscarinic toxins from the black mamba *Dendroaspis polylepis*, *Eur J Biochem* **585** (2), 579–585, 1995.
- [145] Bourne, Y. et al. Acetylcholinesterase Inhibition by Fasciculin: Crystal Structure of the Complex, *Cell* **83** (3), 503–512, 1995.
- [146] Kini, R.M. et al. Flanking proline residues identify the L-type Ca²⁺ channel binding site of calciseptine and FS2, *Biochemistry* **37** (25), 9058–9063, 1998.
- [147] Nguyen, T.T.N. et al. Cardiotoxin-I: An Unexpectedly Potent Insulinotropic Agent, *ChemBioChem* **13**, 1805–1812, 2012.
- [148] Wu, P.L. et al. Non-cytotoxic cobra cardiotoxin A5 binds to $\alpha v\beta 3$ integrin and inhibits bone resorption: Identification of cardiotoxins as non-rgd integrin-binding proteins of the Ly-6 family, *J Biol Chem.* **281** (12), 7937–7945, 2006.
- [149] Cheng, C.H. et al. Dynamics and functional differences between dendroaspin and rhodostomin: Insights into protein scaffolds in integrin recognition, *Protein Sci.* **21** (12), 1872–1884, 2012.
- [150] Banerjee, Y. et al. Hemextin AB Complex – A Snake Venom Anticoagulant Protein Complex That Inhibits Factor VIIa Activity, *Pathophysiol Haemost Thromb.* **543** (4-5), 184–187, 2005.
- [151] Koivula, K. et al. The three-finger toxin MTalpha is a selective alpha(2B)-adrenoceptor antagonist, *Toxicon* **56** (3), 440–447, 2010.

- [152] Quinton, L. et al. Isolation and pharmacological characterization of AdTx1, a natural peptide displaying specific insurmountable antagonism of the α 1A-adrenoceptor, *Br J Pharmacol.* **159** (2), 316–325, 2010.
- [153] Doley, R. et al. Role of accelerated segment switch in exons to alter targeting (ASSET) in the molecular evolution of snake venom proteins, *BMC Evol Biol.* **9**, 146, 2009.
- [154] McDowell, R.S. et al. Mambin, a potent glycoprotein IIb-IIIa antagonist and platelet aggregation inhibitor structurally related to the short neurotoxins, *Biochemistry* **31**, 4766–4772, 1992.
- [155] Wang, C.I. et al. Isolation and structural and pharmacological characterization of alpha-elapitoxin-Dpp2d, an amidated three finger toxin from black mamba venom, *Biochemistry* **53** (23), 3758–3766, 2014.
- [156] Fujimi, T.J. et al. Molecular evolution and diversification of snake toxin genes, revealed by analysis of intron sequences, *Gene* **313**, 111–8, 2003.
- [157] Kini, R.M. Evolution of three-finger toxins - A versatile mini protein scaffold, *Acta Chim Slov.* **58** (4), 693–701, 2011.
- [158] Kini, R.M. Molecular moulds with multiple missions: Functional sites in three-finger toxins, *Clin Exp Pharmacol Physiol.* **29**, 815–822, 2002.
- [159] Mori, N., & Tu, A.T. Isolation and primary structure of the major toxin from sea snake, *Acalyptophis peronii*, venom, *Arch Biochem Biophys.* **260** (1), 10–17, 1988.
- [160] Dewan, J.C. et al. Crystal structure of kappa-bungarotoxin at 2.3-Å resolution, *Biochemistry* **33** (44), 13147–13154, 1994.
- [161] Stewart, C.S. et al. Isolation, characterization and pentamerization of alpha-cobrotoxin specific single-domain antibodies from a naïve phage display library: preliminary findings for antivenom development, *Toxicon* **49** (5), 699–709, 2007.
- [162] Osipov, A. V. et al. The first representative of glycosylated three-fingered toxins: Cytotoxin from the *Naja kaouthia* cobra venom, *Eur J Biochem.* **271** (10), 2018–2027, 2004.
- [163] Pahari, S. et al. The venom gland transcriptome of the Desert Massasauga rattlesnake (*Sistrurus catenatus edwardsii*): towards an understanding of venom composition among advanced snakes (Superfamily Colubroidea), *BMC Mol Biol.* **8**, 115, 2007.
- [164] Utkin, Y.N. et al. “Weak Toxin” from *Naja kaouthia* is a Nontoxic Antagonist of α 7 and Muscle-type Nicotinic Acetylcholine Receptors, *J Biol Chem.* **276** (19), 15810–15815, 2001.

- [165] Kasheverov, I.E. et al. A model for short alpha-neurotoxin bound to nicotinic acetylcholine receptor from *Torpedo californica*: comparison with long-chain alpha-neurotoxins and alpha-conotoxins, **29**, 398–411, 2005.
- [166] Scherf, T. et al. Three-dimensional solution structure of the complex of alpha-bungarotoxin with a library-derived peptide, *Proc Natl Acad Sci* **94** (12), 6059–6064, 1997.
- [167] Osipov, A. V. et al. Dimeric alpha-Cobratoin X-ray structure: Localization of intermolecular disulfides and possible mode of binding to nicotinic acetylcholine receptors, *J Biol Chem.* **287** (9), 6725–6734, 2012.
- [168] Endo, T., & Tamiya, N. *Snake toxin*, Pergamon Press, New york, (US states), 1991.
- [169] Trémeau, O. et al. Genetic engineering of snake toxins: The functional site of erabutoxin A, as delineated by site-directed mutagenesis, includes variant residues, *J Biol Chem.* **270** (16), 9362–9369, 1995.
- [170] Antil, S. et al. Variability among the sites by which curaremimetic toxins bind to *Torpedo* acetylcholine receptor, as revealed by identification of the functional residues of α -cobratoin, *J Biol Chem.* **274** (49), 34851–34858, 1999.
- [171] Fiordalisi, J.J. et al. Site-directed mutagenesis of kappa-bungarotoxin: implications for neuronal receptor specificity, *Biochemistry* **33** (13), 3872–7, 1994.
- [172] Grant, G.A., & Chiappinelli, V.A. kappa-Bungarotoxin: complete amino acid sequence of a neuronal nicotinic receptor probe, *Biochemistry* **24** (6), 1532–7, 1985.
- [173] Sue, S. et al. Structures of Heparin-Derived Disaccharide Bound to Cobra Cardiotoxins: Context-Dependent Conformational Change of Heparin upon Binding to the Rigid Core of the Three-Fingered Toxin, *Biochemistry* **40** (36), 10436–10446, 2001.
- [174] Konshina, A.G. et al. Snake cytotoxins bind to membranes via interactions with phosphatidylserine head groups of lipids, *PLoS ONE* **6** (4), e19064, 2011.
- [175] Bilwes, A. et al. X-ray structure at 1.55 Å of toxin gamma, a cardiotoxin from *Naja nigricollis* venom. Crystal packing reveals a model for insertion into membranes, *J Mol Biol.* **239** (1), 122–136, 1994.
- [176] Kini, R.M. et al. Non-enzymatic inhibitors of coagulation and platelet aggregation from *Naja nigricollis* venom are cardiotoxins, *Biochem Biophys Res Commun.* **150** (3), 1012–1016, 1988.

- [177] Kini, R.M., & Evans, H.J. Mechanism of platelet effects of cardiotoxins from *Naja nigricollis* crawshawii (spitting cobra) snake venom, *Thromb Res.* **52** (3), 185–95, 1988.
- [178] Kini, R.M., & Evans, H.J. A common cytolytic region in myotoxins, hemolysins, cardiotoxins and antibacterial peptides, *Int J Pept Protein Res.* **34** (4), 277–86, 1989.
- [179] Kini, R.M., & Evans, H.J. Role of cationic residues in cytolytic activity: modification of lysine residues in the cardiotoxin from *Naja nigricollis* venom and correlation between cytolytic and antiplatelet activity, *Biochemistry* **28** (23), 9209–15, 1989.
- [180] Kini, R.M., & Banerjee, Y. Dissection approach: a simple strategy for the identification of the step of action of anticoagulant agents in the blood coagulation cascade, *J Thromb Haemost.* **3** (1), 170–171, 2004.
- [181] S, S. et al. The inhibition of clotting complexes of the extrinsic coagulation cascade by the phospholipase A2 isoenzymes from *Naja nigricollis* venom, *Thromb Res* **55**, 481–491, 1989.
- [182] Le Du, M.H. et al. Structure of fasciculin 2 from green mamba snake venom: evidence for unusual loop flexibility, *Acta Crystallogr D Biol Crystallogr.* **52** (Pt 1), 87–92, 1996.
- [183] Marchot, P. The fasciculin-acetylcholinesterase interaction, *J Soc Biol.* **193** (6), 505–508, 1999.
- [184] Adem, A. et al. Toxins from the venom of the green mamba *Dendroaspis angusticeps* that inhibit the binding of quinuclidinyl benzilate to muscarinic acetylcholine receptors, *Biochim Biophys Acta.* **968** (3), 340–5, 1988.
- [185] Karlsson, E. et al. Snake toxins with high selectivity for subtypes of muscarinic acetylcholine receptors, *Biochimie* **82** (9-10), 793–806, 2000.
- [186] Carsi, J.M. et al. m2-toxin: A selective ligand for M2 muscarinic receptors, *Mol Pharmacol.* **56** (5), 933–7, 1999.
- [187] Ségalas, I. et al. Solution structure of a green mamba toxin that activates muscarinic acetylcholine receptors, as studied by nuclear magnetic resonance and molecular modeling, *Biochemistry* **34** (4), 1248–60, 1995.
- [188] Bradley, K.N. Muscarinic toxins from the green mamba, *Pharmacol Ther.* **85** (2), 87–109, 2000.
- [189] Näreoja, K. et al. Adrenoceptor activity of muscarinic toxins identified from mamba venoms, *Br J Pharmacol.* **164** (2b), 538–550, 2011.
- [190] Servent, D., & Menez, A. *Handbook of Neurotoxicology*, Massaro, E. Human Press, Humana, Totowa, NJ, 2001.

- [191] Joubert, F.J., & Taljaard, N. Snake venoms. The amino acid sequences of two Melanoleuca-type toxins, *Hoppe Seylers Z Physiol Che.* **361** (3), 425–436, 1980.
- [192] Utkin, Y.N. et al. First tryptophan-containing weak neurotoxin from cobra venom, *Toxicon* **39** (7), 921–927, 2001.
- [193] Chang, L. et al. Structure-function studies on Taiwan cobra long neurotoxin homolog, *Biochim Biophys Acta.* **1480** (1-2), 293–301, 2000.
- [194] Poh, S.L. et al. A synthetic weak neurotoxin binds with low affinity to Torpedo and chicken $\alpha 7$ nicotinic acetylcholine receptors, *Eur J Biochem.* **269** (17), 4247–4256, 2002.
- [195] Aird, S.D. et al. Primary structure of γ -bungarotoxin, a new postsynaptic neurotoxin from venom of *Bungarus multicinctus*, *Toxicon* **37**, 609–625, 1999.
- [196] Torres, A.M. et al. NMR structure of bucardin, a neurotoxin from the venom of the Malayan krait (*Bungarus candidus*), *Biochem J.* **360** (Pt 3), 539–548, 2001.
- [197] Antil-delbeke, S. et al. Protein structure and folding: molecular Determinants by Which a Long Chain Toxin from Snake Venom Interacts with the Neuronal $\alpha 7$ -Nicotinic Acetylcholine Receptor, *J Biol Chem.* **275** (38), 29594–29601, 2000.
- [198] Chiappinelli, V.A. et al. Binding of native kappa-neurotoxins and site-directed mutants to nicotinic acetylcholine receptors, *Toxicon* **34** (11-12), 1243–56, 1996.
- [199] Paterson, D., & Nordberg, A. Neuronal nicotinic receptors in the human brain, *Prog Neurobiol.* **61** (1), 75–111, 2000.
- [200] Lindstrom, J. Nicotinic acetylcholine receptors in health and disease, *Mol Neurobiol.* **15** (2), 193–222, 1997.
- [201] Wu, J., & Lukas, R.J. Naturally-expressed nicotinic acetylcholine receptor subtypes, *Biochem Pharmacol.* **82** (8), 800–807, 2011.
- [202] De Welle, J.R. et al. Calciseptine, a peptide isolated from black mamba venom, is a specific blocker of the L-type calcium channel, *Proc Natl Acad Sci* **88** (6), 2437–40, 1991.
- [203] Yasuda, O. et al. FS2. a mamba venom toxin, is a specific blocker of the L-type calcium channels, *Artery* **21**, 287–302, 1994.
- [204] Kini, R.M., & Evans, H.J. Prediction of potential protein-protein interaction sites from amino acid sequence. Identification of a fibrin polymerization site, *FEBS Lett.* **385** (1-2), 81–6, 1996.

- [205] Watanabe, T.X. et al. Smooth muscle relaxing and hypotensive activities of synthetic calciseptine and the homologous snake venom peptide FS2, *Jpn J Pharmacol.* **68** (3), 305–13, 1995.
- [206] Gatineau, E. et al. Role of tyrosine and tryptophan residues in the structure-activity relationships of a cardiotoxin from *Naja nigricollis* venom, *Biochemistry* **26** (25), 8046–8055, 1987.
- [207] Waldo, A.L. et al. Effect of d-sotalol on mortality in patients with left ventricular dysfunction after recent and remote myocardial infarction. The SWORD Investigators. Survival With Oral d-Sotalol, *Lancet* **348** (9019), 7–12, 1996.
- [208] Joubert, F.J., & Taljaard, N. Some properties and the complete primary structures of two reduced and S-carboxymethylated polypeptides (S5C1 and S5C10) from *Dendroaspis jamesoni kaimosae* (Jameson's mamba) venom, *Biochim Biophys Acta* **579** (1), 228–33, 1979.
- [209] Williams, J.A. Disintegrins: RGD-containing proteins which inhibit cell/matrix interactions (adhesion) and cell/cell interactions (aggregation) via the integrin receptors, *Pathol Biol (Paris)*. **40** (8), 813–21, 1992.
- [210] Wattam, B. et al. Arg-Tyr-Asp (RYD) and Arg-Cys-Asp (RCD) motifs in dendroaspin promote selective inhibition of beta1 and beta3 integrins, *Biochem J.* **356** (1), 11–17, 2001.
- [211] Lu, X. et al. The effect of the single substitution of arginine within the RGD tripeptide motif of a modified neurotoxin dendroaspin on its activity of platelet aggregation and cell adhesion, *Cell Commun Adhes.* **13** (3), 171–183, 2006.
- [212] Schleifer, K.J. Comparative molecular modelling study of the calcium channel blockers nifedipine and black mamba toxin FS2, *J Comput Aided Mol Des.* **11** (5), 491–501, 1997.
- [213] Shiu, J.-H. et al. Solution structure of gamma-bungarotoxin: the functional significance of amino acid residues flanking the RGD motif in integrin binding, *Proteins* **57** (4), 839–49, 2004.
- [214] Sutcliffe, M.J. et al. Three-dimensional structure of the RGD-containing neurotoxin homologue dendroaspin, *Nat Struct Biol.* **1** (11), 802–7, 1994.
- [215] Pu, X.C. et al. A novel analgesic toxin (Hannalgesin) from the venom of king cobra (*Ophiophagus hannah*), *Toxicon* **33** (11), 1425–1431, 1995.
- [216] Nei, M., & Rooney, A.P. Concerted and birth-and-death evolution of multigene families, *Annu rev genet* **39**, 121–52, 2005.
- [217] Fry, B.G. et al. Molecular evolution and phylogeny of elapid snake venom three-finger toxins, *J Mol Evol.* **57** (1), 110–129, 2003.

- [218] Daltry, J.C. et al. Diet and snake venom evolution, *Nature* **379**, 537–540, 1996.
- [219] Sekhar, C.C., & Chakrabarty, D. Fibrinogenolytic toxin from Indian monocled cobra (*Naja kaouthia*) venom, *J Biosci.* **36** (2), 355–361, 2011.
- [220] Mukherjee, A.K. Phospholipase A2-interacting weak neurotoxins from venom of monocled cobra *Naja kaouthia* display cell-specific cytotoxicity, *Toxicon* **51**, 1538–1543, 2008.
- [221] Debnath, A. et al. A lethal cardiotoxic-cytotoxic protein from the Indian monocellate cobra (*Naja kaouthia*) venom, *Toxicon* **56** (4), 569–579, 2010.
- [222] Mackessy, S.P. Venom ontogeny in the Pacific rattlesnakes *Crotalus viridis helleri* and *C. v. oreganus*, *Copeia* 92–101, 1988.
- [223] Lowry, O.H. et al. Protein measurement with the Folin phenol reagent, *J Biol Chem.* **193** (1), 265–275, 1951.
- [224] Laemmli, U.K. Cleavage of structural proteins during the assembly of the head of bacteriophage T4, *Nature* **227** (5259), 680–685, 1970.
- [225] Hendrickson, H.S. et al. Chiral synthesis of a dithiolester analog of phosphatidylcholine as a substrate for the assay of phospholipase A2, *J Lipid Res.* **24** (11), 1532–1537, 1983.
- [226] Reynolds, L.J. et al. Analysis of human synovial fluid phospholipase A2 on short chain phosphatidylcholine-mixed micelles: development of a spectrophotometric assay suitable for a microtiterplate reader, *Anal Biochem.* **204** (1), 190–197, 1992.
- [227] Joubert, F.J., & Taljaard, N. *Naja haje haje* (Egyptian cobra) venom. Some properties and the complete primary structure of three toxins (CM-2, CM-11 and CM-12), *Eur J Biochem.* **90** (2), 359–367, 1978.
- [228] Meier, J., & Theakston, R.D. Approximate LD50 determinations of snake venoms using eight to ten experimental animals, *Toxicon* **24** (4), 395–401, 1986.
- [229] Yamakawa K, Nozaki M, and H.Z. *Animal, Plant, and Microbial Toxins*, Springer New York, 1976.
- [230] Vishwanath, B.S., & Gowda, T. V. Interaction of aristolochic acid with *Vipera russelli* phospholipase A2: its effect on enzymatic and pathological activities, *Toxicon* **25** (9), 929–937, 1987.
- [231] Vishwanath, B.S. et al. Purification and partial biochemical characterization of an edema inducing phospholipase A2 from *Vipera russelli* (Russell's viper) snake venom, *Toxicon* **26** (8), 713–720, 1988.

- [232] Kondo, H. et al. Studies on the quantitative method for determination of hemorrhagic activity of Habu snake venom, *Jpn J Med Sci Biol.* **13**, 43–52, 1960.
- [233] Parasuraman, S. et al. Blood sample collection in small laboratory animals, *J Pharmacol Pharmacother.* **1** (2), 87–93, 2010.
- [234] McQueen, M.J. True Arrhenius relationships of human lactate dehydrogenase, *J. Clin. Chem. Clin. Biochem* **13** (1), 17–19, 1975.
- [235] Witt, I., & Trendelenburg, C. Joint study to establish reference values for clinical chemical parameters in childhood (author's transl), *J. Clin. Chem. Clin. Biochem* **20** (4), 235–242, 1982.
- [236] Hohorst, H.J. et al. Studies on the creatine kinase equilibrium in muscle and the significance of ATP and ADP levels, *Biochem Biophys Res Commun.* **7**, 142–146, 1962.
- [237] Turner, D.C. et al. A protein that binds specifically to the M-line of skeletal muscle is identified as the muscle form of creatine kinase, *Proc Natl Acad Sci* **70** (3), 702–705, 1973.
- [238] Kontinen, A., & Somer, H. Specificity of serum creatine kinase isoenzymes in diagnosis of acute myocardial infarction, *Br Med J.* **1** (5850), 386–389, 1973.
- [239] Ferlitsch, A. et al. The diagnostic value of CK-isoenzymes in suspected acute myocardial infarction (author's transl), *Dtsch Med Wochenschr.* **101** (9), 312–317, 1976.
- [240] Somer, H. et al. Creatine kinase isoenzymes in neuromuscular diseases, *J Neurol Sci.* **29** (2-4), 129–136, 1976.
- [241] Mukherjee, A.K., & Maity, C.R. Biochemical composition, lethality and pathophysiology of venom from two cobras - *Naja naja* and *N. kaouthia*, *Comp Biochem Physiol.* **131** (2), 125–132, 2002.
- [242] Ouyang, C., & Teng, C.M. Fibrinogenolytic enzymes of *Trimeresurus mucrosquamatus* venom, *Biochim Biophys Acta.* **420** (2), 298–308, 1976.
- [243] Eagle, H. Studies on blood coagulation: II. The formation of fibrin from thrombin and fibrinogen, *J Gen Physiol.* **18** (4), 547–555, 1935.
- [244] Henschen, A. et al. Covalent structure of fibrinogen, *Ann N Y Acad Sci.* **408**, 28–43, 1983.
- [245] Mosesson, M.W. Fibrinogen and fibrin structure and functions, *J Thromb Haemost.* **3** (8), 1894–1904, 2005.

- [246] Iwanaga, S. Structure and function of fibrinogen, *Tanpakushitsu kakusan koso*. **14** (3), 175–195, 1969.
- [247] Lang, T. et al. Multi-centre investigation on reference ranges for ROTEM thromboelastometry, *Blood Coagul Fibrinolysis*. **16** (4), 301–310, 2005.
- [248] Niemi, T.T., & Kuitunen, A.H. Artificial colloids impair haemostasis. An in vitro study using thromboelastometry coagulation analysis, *Acta Anaesthesiol Scand*. **49** (3), 373–378, 2005.
- [249] Steer, P.L., & Krantz, H.B. Thromboelastography and Sonoclot analysis in the healthy parturient, *J Clin Anesth*. **5** (5), 419–424, 1993.
- [250] LaForce, W.R. et al. Evaluation of the SonoClot Analyzer for the measurement of platelet function in whole blood, *Ann Clin Lab Sci*. **22** (1), 30–33, 1992.
- [251] Gutiérrez, J.M. et al. Impact of regional variation in *Bothrops asper* snake venom on the design of antivenoms: Integrating antivenomics and neutralization approaches, *J Proteome Res*. **9** (1), 564–577, 2010.
- [252] Katsuki, R. et al. Tramadol, but not its major metabolite (mono-O-demethyl tramadol) depresses compound action potentials in frog sciatic nerves, *Br J Pharmacol*. **149** (3), 319–327, 2006.
- [253] Edman, P. A method for the determination of amino acid sequence in peptides, *Arch Biochem*. **22** (3), 475, 1949.
- [254] Mukherjee, A.K. Non-covalent interaction of phospholipase A(2) (PLA(2)) and kaouthiotoxin (KTX) from venom of *Naja kaouthia* exhibits marked synergism to potentiate their cytotoxicity on target cells, *J venom res*. **1**, 37–42, 2010.
- [255] Yamaguchi, Y. et al. Characterization, amino acid sequence and evolution of edema-inducing, basic phospholipase A2 from *Trimeresurus flavoviridis* venom, *Toxicon* **39** (7), 1069–1076, 2001.
- [256] Reali, M. et al. Neurotoxic and myotoxic actions of *Naja naja kaouthia* venom on skeletal muscle in vitro, *Toxicon* **41**, 657–665, 2003.
- [257] Persson, B. et al. Elapid venom toxins: multiple recruitments of ancient scaffolds, *Eur J Biochem*. **234** (1-2), 225–234, 1999.
- [258] Nanjaraj Urs, A.N. et al. Progressive Hemorrhage and Myotoxicity Induced by *Echis carinatus* Venom in Murine Model: Neutralization by Inhibitor Cocktail of N,N,N',N'-Tetrakis (2-Pyridylmethyl) Ethane-1,2-Diamine and Silymarin, *PLoS ONE* **10** (8), e0135843, 2015.
- [259] Teixeira, C.F.P. et al. Inflammatory effects of snake venom myotoxic phospholipases A2, *Toxicon* **42** (8), 947–962, 2003.

- [260] Uma, B., & Veerabasappa Gowda, T. Molecular mechanism of lung hemorrhage induction by VRV-PL-VIIIa from Russell's viper (*Vipera russelli*) venom, *Toxicon* **38** (8), 1129–1147, 2000.
- [261] Saravia, P. et al. Geographic and ontogenic variability in the venom of the neotropical rattlesnake *Crotalus durissus*: Pathophysiological and therapeutic implications, *Rev Biol Trop.* **50** (1), 337–346, 2002.
- [262] Chippaux, J.P. et al. Snake venom variability: Methods of study, results and interpretation, *Toxicon* **29** (11), 1279–1303, 1991.
- [263] Leong, P.K. et al. Cross neutralisation of Southeast Asian cobra and krait venoms by Indian polyvalent antivenoms, *Trans R Soc Trop Med Hyg.* **106** (12), 731–737, 2012.
- [264] Sells, P.G. Animal experimentation in snake venom research and in vitro alternatives, *Toxicon* **42** (2), 115–133, 2003.
- [265] Kuruppu, S. et al. Neurotoxins from Australo-Papuan elapids: a biochemical and pharmacological perspective, *Crit Rev Toxicol.* **38** (1), 73–86, 2008.
- [266] Hodgson, W.C., & Wickramaratna, J.C. In vitro neuromuscular activity of snake venoms, *Clin Exp Pharmacol Physiol.* **29** (9), 807–814, 2002.
- [267] Chen, K.C. et al. The mechanism of cytotoxicity by *Naja naja atra* cardiotoxin 3 is physically distant from its membrane-damaging effect, *Toxicon* **50** (6), 816–824, 2007.
- [268] Symer, D.E. et al. A requirement for membrane-associated phospholipase A2 in platelet cytotoxicity activated by receptors for immunoglobulin G and complement, *J Exp Med.* **177** (4), 937–947, 1993.
- [269] Vulfius, C.A. et al. Inhibition of nicotinic acetylcholine receptors, a novel facet in the pleiotropic activities of snake venom phospholipases A2, *PLoS ONE* **9** (12), e115428, 2014.
- [270] Chiou, Y.-L. et al. Modulated mechanism of phosphatidylserine on the catalytic activity of *Naja naja atra* phospholipase A2 and *Notechis scutatus* scutatus notexin, *Toxicon* **92**, 113–22, 2014.
- [271] Kini, R.M. Structure-function relationships and mechanism of anticoagulant phospholipase A2 enzymes from snake venoms, *Toxicon* **45** (8), 1147–1161, 2005.
- [272] Dutta, S. et al. Anticoagulant mechanism and platelet deaggregation property of a non-cytotoxic, acidic phospholipase A2 purified from Indian cobra (*Naja naja*) venom: Inhibition of anticoagulant activity by low molecular weight heparin, *Biochimie* **110**, 93–106, 2015.

- [273] Wijeyewickrema, L.C. et al. Fractionation of snake venom metalloproteinases by metal ion affinity: A purified cobra metalloproteinase, Nk, from *Naja kaouthia* binds Ni²⁺-agarose, *Toxicon* **50** (8), 1064–1072, 2007.
- [274] Dufton, M.J., & Hider, R.C. Structure and pharmacology of elapid cytotoxins, *Pharmacol Ther.* **36** (1), 1–40, 1988.
- [275] Kumar, T.K. et al. A case study of cardiotoxin III from the Taiwan cobra (*Naja naja atra*). Solution structure and other physical properties, *Adv Exp Med Biol.* **391**, 115–129, 1996.
- [276] Feofanov, A. V et al. Cancer cell injury by cytotoxins from cobra venom is mediated through lysosomal damage, *Biochem J.* **390** (Pt 1), 11–18, 2005.
- [277] Debnath, A. et al. Venom of Indian monocellate cobra and *Russell's viper* show anticancer activity in experimental models, *J Ethnopharmacol.* **111** (3), 681–684, 2007.
- [278] Gutiérrez, J.M., & Rucavado, A. Snake venom metalloproteinases: their role in the pathogenesis of local tissue damage, *Biochimie* **82** (9-10), 841–850, 2000.
- [279] Hamako, J. et al. Purification and characterization of kaouthiagin, a von Willebrand factor-binding and -cleaving metalloproteinase from *Naja kaouthia* cobra venom, *Thromb Haemost.* **80** (3), 499–505, 1998.
- [280] Chen, L.W. et al. Bactericidal effect of *Naja nigricollis* toxin γ is related to its membrane-damaging activity, *Peptides* **32** (8), 1755–1763, 2011.
- [281] Chen, L.W. et al. Membrane-damaging activity of Taiwan cobra cardiotoxin 3 is responsible for its bactericidal activity, *Toxicon* **58** (1), 46–53, 2011.
- [282] Xu, C. et al. A bactericidal homodimeric phospholipases A₂ from *Bungarus fasciatus* venom, *Peptides* **28** (5), 969–973, 2007.
- [283] Mordvintsev, D.Y. et al. Behavioural effects in mice and intoxication symptomatology of weak neurotoxin from cobra *Naja kaouthia*, *Basic Clin Pharmacol Toxicol.* **100** (4), 273–278, 2007.
- [284] Behrooz Fathi, H. et al. The facilitatory actions of snake venom phospholipase A₂ neurotoxins at the neuromuscular junction are not mediated through voltage-gated K⁺ channels, *Toxicon* **39**, 1871–1882, 2001.
- [285] Shashidharamurthy, R. et al. Variations in biochemical and pharmacological properties of Indian cobra (*Naja naja naja*) venom due to geographical distribution, *Mol Cell Biochem.* **229** (1-2), 93–101, 2002.
- [286] Shashidharamurthy, R., & Kemparaju, K. Region-specific neutralization of Indian cobra (*Naja naja*) venom by polyclonal antibody raised against the eastern regional venom: A comparative study of the venoms from three

- different geographical distributions, *Int Immunopharmacol.* **7** (1), 61–69, 2007.
- [287] Calvete, J.J. et al. Snake venomomics. Strategy and applications, *J Mass Spectrom.* **42** (11), 1405–1414, 2007.
- [288] Vonk, F.J. et al. Snake venom: From fieldwork to the clinic: Recent insights into snake biology, together with new technology allowing high-throughput screening of venom, bring new hope for drug discovery, *BioEssays* **33** (4), 269–79, 2011.
- [289] Warrell, D.A. Snake bite, *The Lancet* **375**, 77–88, 2010.
- [290] Gutiérrez, J.M. et al. Confronting the neglected problem of snake bite envenoming: The need for a global partnership, *PLoS Med.* **3** (6), 0727–0731, 2006.
- [291] Uetz., P., & Hosek, J. The Reptile Database, 2015.
- [292] Wallach, V. et al. In praise of subgenera: taxonomic status of cobras of the genus *Naja Laurenti* (Serpentes: Elapidae), *Zootaxa* **2236**, 26–36, 2009.
- [293] Kulkeaw, K. et al. Proteome and immunome of the venom of the Thai cobra, *Naja kaouthia*, *Toxicon* **49** (7), 1026–1041, 2007.
- [294] Earl, S.T.H. et al. Post-translational modification accounts for the presence of varied forms of nerve growth factor in Australian elapid snake venoms, *Proteomics* **6** (24), 6554–6565, 2006.
- [295] Malih, I. et al. Proteomic analysis of moroccan cobra *naja haje legionis* venom using tandem mass spectrometry, *J Proteomics.* **96**, 240–252, 2014.
- [296] Chem, J.J.B. et al. Characterization of Elapidae Snake Venom Components Using Optimized Reverse-Phase High-Performance Liquid Chromatographic Conditions and Screening Assays for Neurotoxin and Phospholipase A₂ Activities, *Biochemistry* **25** (22), 7235–7243, 1986.
- [297] Takacs, Z. et al. Cobra (*Naja spp* .) Nicotinic Acetylcholine Receptor Exhibits Resistance to Erabu Sea Snake (*Laticauda semifasciata*) Short-Chain alpha Neurotoxin, *J Mol Evol.* **58** (5), 516–526, 2004.
- [298] Tsetlin, V.I. Three-finger snake neurotoxins and Ly6 proteins targeting nicotinic acetylcholine receptors: pharmacological tools and endogenous modulators, *Trends Pharmacol Sci.* **36** (2), 109–23, 2015.
- [299] Utkin, Y.N. Three-finger toxins, a deadly weapon of elapid venom - Milestones of discovery, *Toxicon* **62**, 50–55, 2013.
- [300] Batuwangala, T. et al. Structure of the snake-venom toxin convulxin, *Acta Crystallogr D Biol Crystallogr.* **60** (Pt 1), 46–53, 2004.

- [301] Blanchet, G. et al. Polypharmacology profiles and phylogenetic analysis of three-finger toxins from mamba venom: Case of aminergic toxins, *Biochimie* **103**, 109–117, 2014.
- [302] Fruchart-gaillard, C. et al. How Three-Finger-Fold Toxins Interact With Various Cholinergic Receptors, *J Mol Neurosci.* **30** (1), 7–8, 2006.
- [303] Gasanov, S.E. et al. Snake Venom Cytotoxins, Phospholipase A₂s, and Zn²⁺-dependent Metalloproteinases: Mechanisms of Action and Pharmacological Relevance, *J Clin Toxicol.* **4** (1), 1000181, 2014.
- [304] Qian, Y.C. et al. cDNA sequence analysis and expression of four long neurotoxin homologues from *Naja naja atra*, *Biochim Biophys Acta.* **1443** (1-2), 233–238, 1998.
- [305] Dufton, M.J., & Hider, R.C. Conformational properties of the neurotoxins and cytotoxins isolated from Elapid snake venoms, *CRC Crit Rev Biochem.* **14** (2), 113–171, 1983.
- [306] Endo, T., & Tamiya, N. Current view on the structure-function relationship of postsynaptic neurotoxins from snake venoms, *Pharmacol Ther.* **34** (3), 403–451, 1987.
- [307] Pillet, L. et al. Genetic engineering of snake toxins: Role of invariant residues in the structural and functional properties of a curaremimetic toxin, as probed by site-directed mutagenesis, *J Biol Chem.* **268** (2), 909–916, 1993.
- [308] Nirathanan, S., & Gwee, M.C.E. Critical Review Three-Finger -Neurotoxins and the Nicotinic Acetylcholine Receptor, Forty Years On, *J. Pharmacol. Sci.* **17** (1), 1–17, 2004.
- [309] Dubovskii, P. V et al. Cobra cardiotoxins: membrane interactions and pharmacological potential, *Curr Med Chem.* **21** (3), 270–287, 2014.
- [310] Armugam, A., & Jeyaseelan, K. Cytotoxic potency of cardiotoxin from *Naja sputatrix*: development of a new cytolytic assay, *Biochem. J.* **43**, 35–43, 2002.
- [311] Chu, N.-S. Contribution of a snake venom toxin to myasthenia gravis: the discovery of alpha-bungarotoxin in Taiwan, *J Hist Neurosci.* **14** (2), 138–148, 2005.
- [312] Rogers, C.M., & Brown, E.R. Differential sensitivity to calciseptine of L-type Ca²⁺ currents in a “lower” vertebrate (*Scyliorhinus canicula*), a protochordate (*Branchiostoma lanceolatum*) and an invertebrate (*Alloteuthis subulata*), *Exp Physiol.* **86** (6), 689–694, 2001.
- [313] Tsetlin, V.I. et al. Conformational studies of neurotoxin II from *Naja naja oxiana*. Selective N-acylation, circular dichroism and nuclear-magnetic-resonance study of acylation products, *Eur J Biochem.* **94** (2), 337–346, 1979.

- [314] Roly, Z.Y. et al. A comparative in silico characterization of functional and physicochemical properties of 3FTx (three finger toxin) proteins from four venomous snakes, *Bioinformation* **10** (5), 281–287, 2014.
- [315] Blanchet, G. et al. Polypharmacology profiles and phylogenetic analysis of three-finger toxins from mamba venom: Case of aminergic toxins, *Biochimie* **103**, 109–117, 2014.
- [316] Ñ, K.J. et al. Structurally conserved K⁻-neurotoxin genes encode functionally diverse proteins in the venom of *Naja sputatrix* 1, **553**, 333–341, 2003.
- [317] Chang, L. et al. Purification and characterization of a neurotoxin from the venom of *Ophiophagus hannah* (king cobra), *Biochem Biophys Res Commun.* **294** (3), 574–578, 2002.
- [318] Achyuthan, K.E., & Ramachandran, L.K. Cardiotoxin of the Indian cobra (*Naja naja*) is a pyrophosphatase, **3** (2), 149–155, 1981.
- [319] Krarup, C. Compound sensory action potential in normal and pathological human nerves, *Muscle Nerve.* **29** (4), 465–483, 2004.
- [320] Katoh, E. et al. Structural basis for the biological activity of dendrotoxin-I, a potent potassium channel blocker, *Biopolymers* **54** (1), 44–57, 2000.
- [321] Mukherjee, A.K. Correlation between the phospholipids domains of the target cell membrane and the extent of *Naja kaouthia* PLA₂-induced membrane damage: Evidence of distinct catalytic and cytotoxic sites in PLA₂ molecules, *Biochim Biophys Acta.* **1770** (2), 187–195, 2007.
- [322] Fletcher, J.E. et al. Similarities and differences in mechanisms of cardiotoxins, melittin and other myotoxins, *Toxicon* **34** (11-12), 1301–1311, 1996.
- [323] Vincent, J.P. et al. Properties of association of cardiotoxin with lipid vesicles and natural membranes. A fluorescence study, *FEBS Lett.* **85** (1), 103–108, 1978.
- [324] Aripov, T.F. et al. Central Asian cobra venom cytotoxins-induced aggregation, permeability and fusion of liposomes, *Gen Physiol Biophys.* **8** (5), 459–473, 1989.
- [325] Chien, K.Y. et al. Fusion of sphingomyelin vesicles induced by proteins from Taiwan cobra (*Naja naja atra*) venom. Interactions of zwitterionic phospholipids with cardiotoxin analogues, *J Biol Chem.* **266** (5), 3252–9, 1991.
- [326] Carbone, M.A., & Macdonald, P.M. Cardiotoxin II segregates phosphatidylglycerol from mixtures with phosphatidylcholine: (31)P and (2)H NMR spectroscopic evidence, *Biochemistry* **35** (11), 3368–78, 1996.

- [327] Aripov, T.F. et al. The influence of cytotoxins from Central Asian cobra venom and melittin from bee venom on the thermodynamic properties of phospholipid bilayer, *Gen Physiol Biophys.* **6** (4), 343–57, 1987.
- [328] Lee, S.C. et al. Endocytotic routes of cobra cardiotoxins depend on spatial distribution of positively charged and hydrophobic domains to target distinct types of sulfated glycoconjugates on cell surface, *J Biol Chem.* **289** (29), 20170–20181, 2014.
- [329] Chien, K.Y. et al. Two distinct types of cardiotoxin as revealed by the structure and activity relationship of their interaction with zwitterionic phospholipid dispersions, *J Biol Chem.* **269** (20), 14473–83, 1994.
- [330] Konshina, A.G. et al. Interaction of cardiotoxin A5 with membrane: Role of conformational heterogeneity and hydrophobic properties, *Bioorg Khim.* **29** (6), 523–533, 2003.
- [331] Lu, X. et al. Preferential antagonism of the interactions of the integrin alpha IIb beta 3 with immobilized glycoprotein ligands by snake-venom RGD (Arg-Gly-Asp) proteins. Evidence supporting a functional role for the amino acid residues flanking the tripeptide RGD in , *Biochem J.* **304** (Pt 3), 929–936, 1994.
- [332] Shiu, J.H. et al. Solution structure of gamma-bungarotoxin: The functional significance of amino acid residues flanking the RGD motif in integrin binding, *Proteins* **57** (4), 839–849, 2004.
- [333] Ferraro, E. et al. Molecular control of neuromuscular junction development, *J Cachexia Sarcopenia Muscle* **3**, 13–23, 2012.
- [334] Ogay, A.Y. et al. Weak neurotoxin from *Naja kaouthia* cobra venom affects haemodynamic regulation by acting on acetylcholine receptors, *Toxicon* **45** (1), 93–99, 2005.
- [335] Chang, L. Sen et al. Structure-function studies on Taiwan cobra long neurotoxin homolog, *Biochim Biophys Acta.* **1480** (1-2), 293–301, 2000.
- [336] Chang, L.S. et al. Characterization and gene organization of Taiwan banded krait (*Bungarus multicinctus*) γ -bungarotoxin, *J Protein Chem.* **21** (4), 223–229, 2002.
- [337] Mordvintsev, D.Y. et al. Weak toxin WTX from *Naja kaouthia* cobra venom interacts with both nicotinic and muscarinic acetylcholine receptors, *FEBS J.* **276** (18), 5065–5075, 2009.
- [338] Goldstein, S.A.N. et al. The charybdotoxin receptor of a Shaker K⁺ channel: Peptide and channel residues mediating molecular recognition, *Neuron* **12**, 1377–1388, 1994.

- [339] Miller, C. et al. Charybdotoxin, a protein inhibitor of single Ca^{2+} -activated K^+ channels from mammalian skeletal muscle, *Nature* **313** (6000), 316–318, 1985.
- [340] Floriano, R.S. et al. The neuromuscular activity of *Bothriopsis bilineata smaragdina* (forest viper) venom and its toxin Bbil-TX (Asp49 phospholipase A) on isolated mouse nerve-muscle preparations, *Toxicon* **96C**, 24–37, 2015.
- [341] Stahl, W.L. (Na^+ + K^+)-ATPase: function, structure, and conformations., *Annals of neurology* **16 Suppl**, S121–S127, 1984.
- [342] Werman, R. The specificity of molecular processes involved in neural transmission, *J Theor Biol.* **9** (3), 471–477, 1965.
- [343] Kostial, K. The role of sodium, potassium, calcium, magnesium and lead ions in synaptic transmission, *Arhiv za higijenu rada* **6** (3), 193–200, 1955.
- [344] Dreyer, F. Peptide toxins and potassium channels, *Rev Physiol Biochem Pharmacol.* **115**, 93–136, 1990.
- [345] Rettig, J. et al. Inactivation properties of voltage-gated K^+ channels altered by presence of beta-subunit, *Nature* **369** (6478), 289–294, 1994.
- [346] Morales, M.J. et al. A novel β subunit increases rate of inactivation of specific voltage-gated potassium channel α subunits, *J Biol Chem.* **270** (11), 6272–6277, 1995.
- [347] Majumder, K. et al. Molecular cloning and functional expression of a novel potassium channel beta-subunit from human atrium, *FEBS Lett.* **361** (1), 13–16, 1995.
- [348] Dufton, M.J. Proteinase inhibitors and dendrotoxins. Sequence classification, structural prediction and structure/activity, *Eur J Biochem.* **153** (3), 647–654, 1985.
- [349] Harvey, A.L., & Anderson, A.J. Dendrotoxins: snake toxins that block potassium channels and facilitate neurotransmitter release, *Pharmacol Ther.* **31** (1-2), 33–55, 1985.
- [350] Aneiros, A. et al. A potassium channel toxin from the secretion of the sea anemone *Bunodosoma granulifera*. Isolation, amino acid sequence and biological activity, *Biochim Biophys Acta.* **1157** (1), 86–92, 1993.
- [351] Castañeda, O. et al. Characterization of a potassium channel toxin from the Caribbean sea anemone *Stichodactyla helianthus*, *Toxicon* **33** (5), 603–613, 1995.
- [352] Schwartz, E.F. et al. OcyKTx2, a new K^+ -channel toxin characterized from the venom of the scorpion *Opisthacanthus cayaporum*, *Peptides* **46**, 40–46, 2013.

- [353] Rauer, H. et al. Structure-guided transformation of charybdotoxin yields an analog that selectively targets Ca²⁺-activated over voltage-gated K⁺ channels, *J Biol Chem.* **275** (2), 1201–1208, 2000.