CONTENTS

Contents	Page No.
Abstract	i-iv
Declaration	
Certificate of Supervisor	
Certificate of Co-supervisor	
Certificate of the external examiner and ODEC	
Acknowledgement	v-vii
Contents	viii-xi i
List of Tables	xiii-xv
List of Figures	xvi-xvii
List of Abbreviations	xviii-xxi
List of Addreviations	XVIII-XXI
Chapter 1: Introduction	1-6
1.1 Background	1
1.2 Factors associates with cancer	2
1.3 Cancer therapeutics	3
1.4 Chemoprevention	3
1.5 Phytochemicals as therapeutic agent	4
1.6 Aims and objectives of the present study1.7 Expected outcomes	6 6
1.7 Expected outcomes	O
Chapter 2: Review of Literature	7-34
2.1 Background	7
2.2 Present scenario of cancer	8
2.2.1 Global cancer statistics	8
2.2.2 Indian cancer statistics	11
2.3 Currently available therapies and their limitations	12 13
2.4 Chemoprevention: promise for cancer2.5 Mechanism of chemoprevention	15
2.6 Biochemical and molecular targets in chemoprevention	15
2.6.1 Role of xenobiotic metabolizing enzymes	15
2.6.2 Role of antioxidants	16
2.6.3 Role of mitogen activated protein kinase (MAPKs)	17
2.6.4 Role of mammalian target of rapamycin (mTOR)	18
2.6.5 Role of nuclear factor kappa B(NF-κB)	19
2.6.6 Role of Kelch-like ECH-associated protein 1 (KEAP1)	20
2.6.7 Role of signal transducers and activators of transcription (STATs)	20
2.7 Clinical basis of chemoprevention	21
2.8 Model system for chemoprevention study	22 23
2.8.1 DMBA induced and croton oil promoted skin papillomagenesis2.8.2 Benzo(a)pyrene induced forestomach papillomagenesis	23
2.9 Phytochemicals as chemopreventive and anticancer agent	24
2.9.1 Phenolic compounds	26

2.9.1.1 Gallic acid	26
2.9.1.2 Caffeic acid	27
2.9.1.3 Ferulic acid	27
2.9.2 Flavonoids	28
2.9.2.1 Quercetin	28
2.9.2.2 Kaempferol	28
2.9.2.3 Apigenin	29
2.9.2.4 Luteolin	29
2.9.2.5 Epigallocatechin gallate (EGCG)	30
2.9.2.6 Silibinin	30
2.9.3 Alkaloids	30
2.9.3.1 Vinca alkaloid	31
2.9.3.2 Taxen	31
2.9.4 Saponin	31
2.10 Traditional knowledge and drug discovery	32
2.11 Biodiversity of North East India	33
2.12 Conclusion	34
Chapter 3: Selection and identification of medicinal plant	35-41
3.1 Background	35
3.2 Method	36
3.1.1 Study area	36
3.1.2 Selection and collection of medicinal plants	36
3.1.3 Taxonomic identification and authentication	37
3.3 Results	37
3.4 Discussion and conclusion	38
5.1 Biseussion and conclusion	50
Chapter 4: Protective function of <i>Phlogacanthus tubiflorus</i> and <i>Nyctanthes</i>	
arbor-tristis against oxidative stress induced by H ₂ O ₂	42-52
4.1 Background	42
4.2 Materials and methods	43
4.2.1 Chemicals and reagents	43
4.2.2 Preparation of extract	43
4.2.3 Evaluation of protective function of REPT and FENA	43
4.2.3.1 Isolation of lymphocytes	44
4.2.3.2 Lymphocyte culture, treatment and preparation of cell lysate	44
4.2.3.3 MTT based cell viability assay	44
4.2.3.4 Estimation of Lactate Dehydrogenase (EC 1.1.1.27)	45
4.2.3.5 Estimation of Lipid Peroxidation	45
4.2.3.6 Estimation of Glutathione-S-transferase (EC 2.5.1.18)	45
4.2.3.7 Estimation of reduced glutathione	46
4.2.3.8 Estimation of Nitric oxide	46
4.2.3.9 Estimation of protein	46
4.2.3.10 Preliminary phytochemical screening	46
4.2.3.11 Statistical analysis	46
4.3 Results	46
4.3.1 Effects of oxidative stress induced by H ₂ O ₂	47
4.3.2 Modulatory effect of REPT	47
4.3.3 Protective function of REPT	47 50
4.3.4 Modulatory effect of FENA	50
4.3.5 Protective function of FENA	50

4.3.6 Preliminary phytochemical screening	50
4.4 Discussion and conclusion	50
Chapter 5: Antioxidative and radical scavenging activity of <i>Nyctanthes</i> arbor-tristis and <i>Phlogacanthus thyrsiflorus</i>	53-63
5.1 Background	53
5.2 Materials and methods	54
5.2.1 Chemicals and reagents	54
5.2.2 Preparation of extract	54
5.2.3 Evaluation of antioxidative and free radical scavenging activity of FENA and LEPT	54
5.2.3.1 ABTS radical scavenging activity	54
5.2.3.2 DPPH radical scavenging activity	55
5.2.3.3 Superoxide radical scavenging activity	55
5.2.3.4 Hydroxyl radical scavenging activity	55
5.2.3.5 Fe ²⁺ chelating inhibitory activity	56
5.2.3.6 Reducing power assay	56
5.2.3.7 Detection of characteristic peaks of phytochemical	56
5.2.3.8 Estimation of total phenolic and total flavonoid content	57
5.2.3.9 Statistical analysis	57
5.3 Results	57
5.3.1 Antioxidative and free radical scavenging activity of FENA and LEPT	57
5.3.2 UV-VIS and FTIR spectroscopic analysis	59
5.4 Discussion and conclusion	61
Chapter 6: Chemopreventive potential of Nyctanthes arbor-tristis and Phlogacanthus thyrsiflorus	64-138
	64-138 64
and Phlogacanthus thyrsiflorus	
and <i>Phlogacanthus thyrsiflorus</i> 6.1 Background	64
and <i>Phlogacanthus thyrsiflorus</i> 6.1 Background 6.2 Materials and methods	64 65
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related	64 65 65
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis	64 65 65 65
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against	64 65 65 65 65
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.3 Histological studies	64 65 65 65 65
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis	64 65 65 65 65 66
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.3 Histological studies	64 65 65 65 65 66 66
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.3 Histological studies 6.2.4 Preparation of homogenate of hepatic tissue 6.2.5 Preparation of homogenate of extra-hepatic organs and papilloma	64 65 65 65 65 66 68 70 70
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.3 Histological studies 6.2.4 Preparation of homogenate of hepatic tissue 6.2.5 Preparation of homogenate of extra-hepatic organs and papilloma bearing tissues	64 65 65 65 65 66 68 70 70 71
6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.4 Preparation of homogenate of hepatic tissue 6.2.5 Preparation of homogenate of extra-hepatic organs and papilloma bearing tissues 6.2.6 Biochemical estimation in hepatic, extra-hepatic and papilloma	64 65 65 65 65 66 68 70 70 71
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.4 Preparation of homogenate of hepatic tissue 6.2.5 Preparation of homogenate of extra-hepatic organs and papilloma bearing tissues 6.2.6 Biochemical estimation in hepatic, extra-hepatic and papilloma bearing tissues 6.2.6.1 Estimation of Cytochrome P450 reductase (EC 1.6.2.4) 6.2.6.2 Estimation of Cytochrome b5 reductase (EC 1.6.2.2)	64 65 65 65 65 66 68 70 70 71
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.4 Preparation of homogenate of hepatic tissue 6.2.5 Preparation of homogenate of extra-hepatic organs and papilloma bearing tissues 6.2.6 Biochemical estimation in hepatic, extra-hepatic and papilloma bearing tissues 6.2.6.1 Estimation of Cytochrome P450 reductase (EC 1.6.2.4)	64 65 65 65 65 66 68 70 70 71 71
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.4 Preparation of homogenate of hepatic tissue 6.2.5 Preparation of homogenate of extra-hepatic organs and papilloma bearing tissues 6.2.6 Biochemical estimation in hepatic, extra-hepatic and papilloma bearing tissues 6.2.6.1 Estimation of Cytochrome P450 reductase (EC 1.6.2.4) 6.2.6.2 Estimation of Cytochrome b5 reductase (EC 1.6.2.2)	64 65 65 65 65 66 68 70 71 71 71 72 72
and Phlogacanthus thyrsiflorus 6.1 Background 6.2 Materials and methods 6.2.1 Chemicals and reagents 6.2.2 Experimental design 6.2.2.1 Modulatory effect of FENA and LEPT on the enzymes of xenobiotic metabolism, antioxidant system, toxicity related parameters and detoxification of reactive species 6.2.2.2 Chemopreventive potential of FENA and LEPT against skin papillomagenesis 6.2.2.3 Chemopreventive potential of FENA and LEPT against forestomach papillomagenesis 6.2.4 Preparation of homogenate of hepatic tissue 6.2.5 Preparation of homogenate of extra-hepatic organs and papilloma bearing tissues 6.2.6 Biochemical estimation in hepatic, extra-hepatic and papilloma bearing tissues 6.2.6.1 Estimation of Cytochrome P450 reductase (EC 1.6.2.4) 6.2.6.2 Estimation of Glutathione S-transferase (EC 2.5.1.18)	64 65 65 65 65 66 68 70 70 71 71 72 72 72

6.2.6.7 Estimation of Glutathione Peroxidase (EC 1.11.1.9)	73
6.2.6.8 Estimation of Catalase (EC 1.11.1.6)	73
6.2.6.9 Estimation of Superoxide dismutase (EC 1.15.1.1)	74
6.2.6.10Estimation of Lipid peroxidation	74
6.2.6.11Estimation of Lactate dehydrogenase (EC 1.1.1.27)	74
6.2.6.12Estimation of Reactive oxygen species	74
6.2.6.13Estimation of Nitric oxide	74
6.2.6.14Estimation of Protein	74
6.2.6.15Statistical analysis	75
6.2.7 GC-MS analysis of FENA and LEPT	75
6.2.8 In silico ADME/Tox analysis	76
6.3 Results	76
6.3.1 Modulatory influence and chemopreventive potential of FENA	76
6.3.1.1 General observation and weight gain profile	76
6.3.1.2 Tumor quantification	76
6.3.1.3 Histopathological examination of tumor	78
6.3.1.4 Biochemical estimation in hepatic tissue	80
6.3.1.5 Biochemical estimation in papilloma bearing tissues	81
6.3.1.6 Biochemical estimation in extra-hepatic tissues	82
6.3.2 Phytochemical characterization of FENA by GC MS	83
6.3.3 Pharmacokinetic profile of phytochemicals present in FENA	83
6.3.4 Modulatory influence and chemopreventive potential of LEPT	103
6.3.2.1 General observation and weight gain profile	103
6.3.2.2 Tumor quantification	104
6.3.2.3 Histopathological examination of tumor	107
6.3.2.4 Biochemical estimation in hepatic tissue	107
6.3.2.5 Biochemical estimation in papilloma bearing tissues	108
6.3.2.6 Biochemical estimation in extra-hepatic tissues	109
6.3.5 Phytochemical characterization of LEPT by GC MS	109
6.3.6 Pharmacokinetic profile of phytochemicals present in LEPT	109
6.4 Discussion and conclusion	133
Chapter 7: Anticancer potential of Nyctanthes arbor-tristis	139-162
7.1 Background	139
7.2 Materials and methods	140
7.2.1 Chemicals and reagents	140
7.2.2 Antibodies	140
7.2.3 Cell lines and culture	141
7.2.4 Preparation of fractionated fractions	141
7.2.4.1 Free and bound phenolics rich fraction	141
7.2.4.2 Free and bound flavonoids rich fraction	141
7.2.4.3 Alkaloids rich fraction	142
7.2.5 Screening of fractions	142
7.2.6 Anticancer efficacy of F ³ NAF against human prostate cancer (PCa) cells	143
7.2.6.1 Cell growth and viability assay	143
7.2.6.2 Cell proliferation assay	143
7.2.6.3 Apoptosis assay	143
7.2.6.4 Cell cycle analysis	144
7.2.6.5 Clonogenic assay	144
7.2.6.6 Wound healing assay	144

7.2.6.7 Transwell invasion and migration assay	145
7.2.6.8 Western blotting	145
7.2.6.9 Statistical analysis	146
7.2.7 Phytochemical characterization of F ³ NAF by HR LCMS analysis	146
7.3 Results	146
7.3.1 Selection of active fraction	146
7.3.2 F ³ NAF exhibit anticancer activity against human prostate cancer cells	148
7.3.2.1 F ³ NAF inhibit the cell growth of PCa cells	148
7.3.2.2 F ³ NAF inhibit the proliferation of PCa cells	150
7.3.2.3 Effects of F ³ NAF on apoptosis in PC3 cells	151
7.3.2.4 F ³ NAF arrest progression of cell cycle at G1 and G2 phase	151
in PC3 cells	
7.3.2.5 F ³ NAF modulate the regulators of G1 and G2 phase of cell cycle in PC3 cells	153
7.3.2.6 F ³ NAF inhibits the clonogenicity of PC3 cells	154
7.3.2.7 F ³ NAF modulate the regulators of growth, proliferation and survival in PC3 cells	155
7.3.2.8 F ³ NAF exerts strong anti-migratory and anti-invasive efficacy against PC3 cells	156
7.3.3 Chemical characterization of F ³ NAF	156
7.4 Discussion and conclusion	156
Chapter 8: Summary and Conclusion	163-166
Chapter 9: Bibliography	167-196

Appendix

Annexures (I to VIII)

List of publications

Reprint of papers

Report of plagiarism checking