

# LIST OF TABLES

---

## CHAPTER 2

2.1	Types of traditional fermented soybean products consumed in various Asian countries, including China, Indonesia, Japan, Korea, and Vietnam	9-10
2.2.	Traditional fermented soybean products consumed in India	11-12
2.3	Some traditional fermented milk products consumed worldwide	14
2.4	Some traditional fermented milk products consumed in India	15

## CHAPTER 3

3.1	Antimicrobial activity of <i>L. lactis</i> AMD 17 cell free culture supernatant against different indicator strains	39
3.2	Autoaggregation and Co-aggregation	40
3.3	Total phenolic content, flavonoid content and antioxidant activity of honey samples	43
3.4	Sensory evaluation scores of curd formulations with different concentration of honey (1%- 5%)	45

## CHAPTER 4

4.1	PCR conditions for amplification of BSH gene	59
4.2	Double digestion of pBluescript KS(+)	60
4.3	Restriction digestion of BSH gene	61
4.4	Ligation reactions	61
4.5	Restriction digestion of recombinant plasmid	64
4.6	Restriction digestion of recombinant pBluescript KS(+)	65
4.7	Restriction digestion of pET-22b(+)	66
4.8	Ligation reaction	66
4.9	Composition of experimental high-cholesterolaemic diet	71

4.10A	Simulated gastric fluid (SGF) tolerance test	73
4.10B	Simulated intestinal fluid (SIF) tolerance test	74
4.10C	Effect of bile salt (Oxgall) concentration on the viability of AMD6 and MTCC	75
4.11	Autoaggregation, co-aggregation and hydrophobicity of <i>L. plantarum</i> AMD6 and <i>L. plantarum</i> MTCC 1407	76
4.12	Deconjugation assay	80
4.13	The specific activity of BSH-His using different bile substrates	86
4.14	<i>In vitro</i> cholesterol assimilation by AMD6 and MTCC 1407 with 0.3% bile salts (oxgall) after 48 h	87
4.15	Faecal cholic acid concentration and body weight gain of rats fed with cholesterol-enriched experimental diets or supplemented with isolates AMD6 and reference probiotic strains	88
4.16	Total cholesterol (TC), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C) and low-density lipoprotein cholesterol (LDL-C) levels in the serum of rats fed a high-cholesterol diet alone (HFD) or supplemented with HFD+AMD6 and HFD+MTCC probiotic isolates for 6 weeks	89

## CHAPTER 5

5.1A	Percentage viability of <i>B. amyloliquefaciens</i> AMS1 and <i>L. rhamnosus</i> under simulated gastric fluid (SGF) transit	104
5.1B	Percentage viability of <i>B. amyloliquefaciens</i> AMS1 and <i>L. rhamnosus</i> under simulated Intestinal fluid (SIF) transit	105
5.2	Antimicrobial activity	105
5.3	Antibiotic susceptibility test	109

## CHAPTER 6

6.1	Effect of additives on CMCase activity of cellulase AMS1	133
-----	--	-----

## CHAPTER 7

7.1	Percentage viability of <i>B. subtilis</i> (AMS6) and <i>L. rhamnosus</i> (L.r) under simulated gastric fluid (SGF) and simulated Intestinal fluid (SIF) transit	148
7.2	Antibacterial activity	149
7.3	Antibiotic susceptibility Test	152

---