List of Figures

| 1.1 | Feynmann diagram of Deep Inelastic Scattering | 6 |
|-----|---|----|
| 1.2 | Ladder diagram for DIS in LLQ^2 | 17 |
| 1.3 | Helicity conservation at hadronic vertex in QPM (a) and QCD improved parton model (b). The arrows represent the spin orientations [57] | 19 |
| 1.4 | Leading order Boson Gluon Fusion (BGF) diagram for heavy quark production in <i>ep</i> -collisions | 22 |
| 1.5 | Gluon distribution function extracted at HERA [11] | 24 |
| 1.6 | The reduced DIS cross section as a function of $y^2/(1+(1-y)^2)$ [11] . | 27 |
| 2.1 | $T^2(t) = T_0.T(t)$ and $T^3(t) = T_1.T(t)$ versus $Q^2(GeV^2)$ | 45 |
| 2.2 | t -evolution results of F_L^g structure function up to NNLO using Taylor expansion method in comparison with the H1, ZEUS data and results of | |
| | DL model | 53 |
| 2.3 | x -evolution results of F_L^g structure function up to NLO using Taylor expansion method in comparison with the H1, ZEUS data and results of | 54 |
| | DL model | 34 |
| 2.4 | x -evolution results of F_L^g structure function up to NLO using Taylor expansion method in comparison with the H1, ZEUS data and the the- | |
| | oretical prediction of MSTW08 | 55 |

| 2.5 | x -evolution results of F_L^g structure function up to NLO using Taylor | |
|-----|---|-----------|
| | expansion method in comparison with the H1, ZEUS data and the theoretical prediction of CT10 | 56 |
| 2.6 | x -evolution results of F_L^g structure function up to NLO using Taylor expansion method in comparison with the H1, ZEUS data and the theoretical prediction of ABM11 | 57 |
| 2.7 | x -evolution results of F_L^g structure function up to NLO using Taylor expansion method in comparison with the H1, ZEUS data and the the- | 01 |
| 2.8 | oretical prediction of NNPDF2.3 | 58 |
| | prediction of Boroun et al. (GRB) | 59 |
| 3.1 | t -evolution results of F_L^g structure function up to NLO using Regge theory in comparison with the H1, ZEUS data and results of DL model | 71 |
| 3.2 | x -evolution results of F_L^g structure function up to NLO using Regge theory in comparison with the H1, ZEUS data and results of DL model | 72 |
| 3.3 | x -evolution results of F_L^g structure function up to NLO using Regge theory in comparison with the H1, ZEUS data and the theoretical prediction | |
| 3.4 | of MSTW08 | 73 |
| | of CT10 | 74 |
| 3.5 | x -evolution results of F_L^g structure function up to NLO using Regge theory in comparison with the H1, ZEUS data and the theoretical prediction | |
| | of ABM11 | 75 |

| 3.6 | x-evolution results of F_L^g structure function up to NLO using Regge the- | |
|------|---|----|
| | ory in comparison with the H1, ZEUS data and the theoretical prediction | |
| | of NNPDF2.3 | 76 |
| 3.7 | x -evolution results of ${\cal F}_L^g$ structure function up to NLO using Regge the- | |
| | ory in comparison with the H1 data and the theoretical prediction of | |
| | Boroun (GRB) [13] | 77 |
| 3.8 | Comparison of t-evolution results of ${\cal F}_L^g$ structure function predicted by | |
| | Regge theory approach and Taylor expansion method and DL model. $% \left(1\right) =\left(1\right) \left(1\right) $. | 79 |
| 3.9 | Comparison of x-evolution results of ${\cal F}_L^g$ structure function predicted by | |
| | Regge theory approach and Taylor expansion method and DL model | 80 |
| 3.10 | Comparison of x-evolution results of ${\cal F}_L^g$ structure function predicted by | |
| | Regge theory approach and Taylor expansion method and MSTW08. $$. | 81 |
| 3.11 | Comparison of x-evolution results of ${\cal F}_L^g$ structure function predicted by | |
| | Regge theory approach and Taylor expansion method and CT10 | 82 |
| 3.12 | Comparison of x-evolution results of \mathcal{F}_L^g structure function predicted by | |
| | Regge theory approach and Taylor expansion method and ABM11. $$ | 83 |
| 3.13 | Comparison of x-evolution results of F_L^g structure function predicted by | |
| | Regge theory approach and Taylor expansion method and NNPDF2.3 | 84 |
| 4.1 | Sensitivity of our results of F_L^g structure function in LO with respect | |
| | to the expansion point of gluon density at $z = 0.5, 0.6, 0.7, 0.8, 0.9$ in | |
| | comparison with H1 data and DL model | 94 |
| 4.2 | Sensitivity of our results of ${\cal F}_L^g$ structure function in NLO and NNLO | |
| | with respect to the expansion point of gluon density at $z = 0.5, 0.6, 0.7,$ | |
| | 0.8, 0.9 in comparison with H1 data and DL model | 96 |

| 4.3 | x -evolution results of F_L^g structure function up to NNLO using Taylor expansion method in comparison with the H1, ZEUS data and results of | |
|-----|---|-----|
| | DL model | 98 |
| 4.4 | x -evolution results of F_L^g structure function up to NNLO using Taylor expansion method in comparison with the H1, ZEUS data and the theoretical prediction of MSTW08 | 99 |
| 4.5 | x -evolution results of F_L^g structure function up to NNLO using Taylor expansion method in comparison with the H1, ZEUS data and the the- | |
| | oretical prediction of CT10 | 100 |
| 4.6 | x -evolution results of F_L^g structure function up to NNLO using Taylor expansion method in comparison with the H1, ZEUS data and the the- | |
| | oretical prediction of ABM11 | 101 |
| 4.7 | x -evolution results of F_L^g structure function up to NNLO using Taylor expansion method in comparison with the H1, ZEUS data and the theoretical prediction of NNPDF2.3 | 102 |
| 4.8 | comparison of our x -evolution results of F_L^g structure function with the results of Sarkar et al (CS) and Boroun et al (GRB) | 103 |
| 5.1 | t -evolution results of F_L^g structure function up to NNLO using Regge theory in comparison with the H1, ZEUS data and results of DL model. | 115 |
| 5.2 | x -evolution results of F_L^g structure function up to NNLO using Regge theory in comparison with the H1, ZEUS data and results of DL model. | 116 |
| 5.3 | x -evolution results of F_L^g structure function up to NNLO using Regge theory in comparison with the H1, ZEUS data and the theoretical pre- | |
| | diction of MSTW08 | 117 |

| 5.4 | x-evolution results of F_L^g structure function up to NNLO using Regge | |
|-----|---|-----|
| | theory in comparison with the H1, ZEUS data and the theoretical pre- | |
| | diction of CT10 | 118 |
| 5.5 | x -evolution results of F_L^g structure function up to NNLO using Regge | |
| | theory in comparison with the H1, ZEUS data and the theoretical pre- | |
| | diction of ABM11 | 119 |
| 5.6 | x -evolution results of F_L^g structure function up to NNLO using Regge | |
| | theory in comparison with the H1, ZEUS data and the theoretical pre- | |
| | diction of NNPDF2.3 | 120 |
| 5.7 | x-evolution results of R in comparison with the H1 data and the theo- | |
| | retical prediction of ACOT fit | 121 |
| 5.8 | Comparison of x-evolution results of ${\cal F}_L^g$ structure function up to NNLO | |
| | using Regge theory (RT) and Taylor expansion (TE) method in compar- | |
| | ison with the H1, ZEUS data and the of DL model | 122 |
| 6.1 | x -evolution results of F_L^c structure function using Taylor expansion method | |
| | with the input gluon distribution from DL model | 136 |
| 6.2 | x -evolution results of F_2^c structure function using Taylor expansion method | |
| | in comparison with the H1, ZEUS data | 137 |
| 6.3 | x-evolution results of the ratio of the charm quark structure functions | |
| | R^c using Taylor expansion method | 138 |
| 6.4 | x -evolution results of charm quark reduced cross section σ_r^c using Taylor | |
| | expansion method in comparison with the H1, ZEUS data | 139 |
| 6.5 | Comparison of our results of F_L^c at $Q^2=20,200 GeV^2$ using Taylor expan- | |
| | sion method with the results of colour dipole model (CDM) and Boroun | |
| | et al (GRB) | 140 |

| 6.6 | Comparison of our results of F_2^c at $Q^2=20,200 GeV^2$ using Taylor ex- | |
|------|---|-----|
| | pansion method with the results of DL, colour dipole model (CDM) and | |
| | Boroun et al (GRB) | 140 |
| 6.7 | Results of the charm content of F_L structure function K_L^c with respect | |
| | to x at $Q^2=20,200 GeV^2$ using Taylor expansion method | 141 |
| 6.8 | x -evolution results of F_L^c structure function using Regge theory with the | |
| | input gluon distribution from DL model | 143 |
| 6.9 | x-evolution results of F_2^c structure function using Regge theory in com- | |
| | parison with the H1, ZEUS data | 144 |
| 6.10 | x -evolution results of the ratio of the charm quark structure function R^c | |
| | using Regge theory | 145 |
| 6.11 | x -evolution results of the charm quark reduced cross section σ_r^c using | |
| | Regge theory in comparison with the H1, ZEUS data | 146 |
| 6.12 | Comparison of our results of F_L^c at $Q^2=20,200 GeV^2$ using Regge theory | |
| | with the results of CD model and Boroun et al (GRB) | 147 |
| 6.13 | Comparison of our results of F_2^c at $Q^2=20,200 GeV^2$ using Regge theory | |
| | with the results of DL, CD model and Boroun et al (GRB) | 147 |
| 6.14 | Results of the charm content of F_L structure function K_L^c with respect | |
| | to x at $Q^2=20,200 GeV^2$ using Regge theory | 148 |
| 6.15 | x -evolution results of F_L^b structure function using Taylor expansion method | |
| | with the input gluon distribution from DL model | 150 |
| 6.16 | x -evolution results of \mathbb{F}_2^b structure function using Taylor expansion method | |
| | in comparison with the H1, ZEUS data | 151 |
| 6.17 | x-evolution results of the ratio of the beauty quark structure functions | |
| | R^b using Taylor expansion method | 152 |
| 6.18 | x -evolution results of the beauty quark reduced cross section σ_r^b using | |
| | Taylor expansion method in comparison with the H1, ZEUS data | 153 |

| 6.19 | Comparison of our results of F_2^b at $Q^2=200 GeV^2$ using Taylor expansion | |
|------|---|-----|
| | method with the results of MSTW 08 | 154 |
| 6.20 | Results of the beauty content of F_L structure function K_L^b with respect | |
| | to x at $Q^2=25,200 GeV^2$ using Taylor expansion method | 154 |
| 6.21 | x -evolution results of F_L^b structure function using Regge theory with the | |
| | input gluon distribution from DL model | 156 |
| 6.22 | x -evolution results of \mathbb{F}_2^b structure function using Regge theory in com- | |
| | parison with the H1, ZEUS data | 157 |
| 6.23 | x-evolution results of the ratio of the beauty quark structure functions | |
| | R^b using Regge theory | 158 |
| 6.24 | x -evolution results of the beauty quark reduced cross section σ_r^b using | |
| | Regge theory in comparison with the H1, ZEUS data | 159 |
| 6.25 | Comparison of our results of F_2^b at $Q^2=200 GeV^2$ using Regge theory | |
| | with the results of MSTW08 parameterization | 160 |
| 6.26 | Results of the beauty content of F_L structure function K_L^b with respect | |
| | to x at $Q^2=25,200 GeV^2$ using Regge theory | 160 |
| 6.27 | Comparison of our results of σ_r^c obtained by Taylor expansion (TE) | |
| | method and Regge theory (RT) | 161 |
| 6.28 | Comparison of our results of σ_r^b obtained by Taylor expansion (TE) | |
| | method and Regge theory (RT) | 162 |
| 6.29 | Sensitivity of our results of σ_r^b and σ_r^c with mass renormalization scale μ | |
| | obtained by Taylor expansion (TE) method and Regge theory (RT) | 163 |