

## List of Tables:

| Chapter  | Table | Title   | Page No |
|----------|-------|---|---------|
| <b>2</b> | 2.1   | PBE0/6-31+G* calculated C–C–C bond angles (in degrees) and C <sub>C</sub> –C bond lengths (in Å) of <b>1-34</b> for both singlet and triplet states.  | 31      |
|          | 2.2   | PBE0/6-31+G* computed singlet-triplet ( $\Delta E_{S-T}$ , kcal mol <sup>-1</sup> ) and HOMO–LUMO ( $\Delta E_{H-L}$ , in eV) gaps of <b>1-34</b> .   | 34      |
|          | 2.3   | PBE0/6-31+G* computed hydrogenation energies ( $E_{hydro}$ , kcal mol <sup>-1</sup> ) and stabilization energies (SE, kcal mol <sup>-1</sup> ) of <b>1-34</b> (excluding <b>2</b> and <b>24-27</b> ).   | 36      |
|          | 2.4   | PBE0/6-31+G* calculated energies of $\sigma$ -symmetric lone pair orbital ( $E_{\sigma}$ in eV) and the $\pi$ -symmetric unoccupied orbital ( $E_{\pi^*}$ in eV) concentrated on the carbenic carbon atom (C <sub>C</sub> ) as well as natural charge at C <sub>C</sub> ( $q(C_C)$ ) of <b>1-34</b> (excluding <b>2</b> and <b>24-27</b> ). | 39      |
|          | 2.5   | PBE0/6-31+G* calculated nucleophilicity indices ( $N$ , in eV) and electrophilicity indices ( $\omega$ , in eV) for <b>1-34</b> (excluding <b>2</b> and <b>24-27</b> ).   | 43      |
|          | 2.6   | PBE0/6-31+G* calculated C <sub>C</sub> –P and P–C <sub>Ph</sub> bond lengths (in Å) and <sup>31</sup> P NMR chemical shifts ( $\delta(^{31}P)$ ) for the adducts L–PPh (L= <b>1-34</b> , excluding <b>2</b> and <b>24-27</b> ).   | 47      |
| <b>3</b> | 3.1.1 | Comparison of the singlet-triplet gaps ( $\Delta E_{S-T}$ , in kcal mol <sup>-1</sup> ) using different basis sets and functionals.   | 59      |
|          | 3.1.2 | Comparison of the calculated and observed <sup>13</sup> C NMR chemical shifts (ppm) of experimentally known molecules.  | 60      |
|          | 3.1.3 | PBE0/6-31+G* calculated C <sub>C</sub> –E bond lengths ( $r_1/r_2$ in Å) and E–C <sub>C</sub> –E bond angles (in degrees) in the singlet state geometry of <b>1-22</b> (where E = C or N).  | 63      |
|          | 3.1.4 | PBE0/6-31+G* calculated singlet–triplet separations ( $\Delta E_{S-T}$ , in kcal mol <sup>-1</sup> ) and stabilization energies (SE, in kcal mol <sup>-1</sup> ) of <b>1-22</b> .   | 64      |

## List of Tables:

| Chapter | Table | Title   | Page No |
|---------|-------|---|---------|
| 3       | 3.1.5 | PBE0/6-31+G* calculated energies ( $E_{\sigma}$ , in eV) and hybridization (%s-character) of the $\sigma$ -symmetric lone-pair orbital concentrated at the carbene carbon atom and the respective proton affinity (PA, kcal mol <sup>-1</sup> ) values of <b>1–22</b> .   | 67      |
|         | 3.1.6 | PBE0/6-31+G* calculated energies of the $\sigma$ -symmetric lone-pair orbital concentrated at the carbene carbon atom ( $E_{\sigma}$ , in eV) and the average carbonyl stretching frequencies ( $\nu_{\text{CO}(\text{avg})}$ , in cm <sup>-1</sup> ) of L–Rh(CO) <sub>2</sub> Cl (L: <b>1–22</b> ) complexes.  | 70      |
|         | 3.1.7 | PBE0/6-31+G* calculated nucleophilicity indices ( $N$ , in eV), bond dissociation energies of the Ga–C <sub>C</sub> bond (BDE, kcal mol <sup>-1</sup> ), degree of pyramidalization at the Ga atom ( $\theta_{\text{Ga}}$ , in degree) and Ga–C <sub>C</sub> bond lengths ( $r_{\text{Ga-C}_C}$ , in Å) of the GaCl <sub>3</sub> adducts of <b>1–22</b> . | 71      |
|         | 3.2.1 | PBE0/6-31+G* calculated values of C <sub>C</sub> –C <sub><math>\alpha</math></sub> bond lengths (in Å) and C <sub><math>\alpha</math></sub> –C <sub>C</sub> –C <sub><math>\alpha'</math></sub> bond angles (in degree) for both singlet and triplet states of <b>1–6</b> .  | 80      |
|         | 3.2.2 | PBE0/6-31+G* calculated values of singlet-triplet separations ( $\Delta E_{\text{S-T}}$ , in kcal mol <sup>-1</sup> ) and natural charges at C <sub>C</sub> ( $q(\text{C}_C)$ ) for molecules <b>1–6</b> .  | 81      |
|         | 3.2.3 | Calculated values of singlet-triplet gaps ( $\Delta E_{\text{S-T}}$ , in kcal mol <sup>-1</sup> ) of <b>6PMe<sub>3</sub></b> using different basis sets and functionals.  | 83      |
|         | 3.2.4 | PBE0/6-31+G* calculated NICS(0) and NICS(1) values of molecules <b>1–6</b> .  | 85      |
|         | 3.2.5 | Average isotropic [ $\chi(\text{C}/\Omega)$ ] and out-of-plane [ $\chi_{\text{ZZ}}(\text{C}/\Omega)$ ] interatomic magnetizability values in cgs-ppm units (where $\Omega = \text{C/N}$ ).  | 87      |
|         | 3.2.6 | PBE0/6-31+G* calculated values of hydrogenation energies ( $E_{\text{hydro}}$ , in kcal mol <sup>-1</sup> ) for molecules <b>1–6</b> .  | 88      |
|         | 3.2.7 | PBE0/6-31+G* calculated dimerization reaction energies ( $\Delta E_{\text{dimer}}$ , in kcal mol <sup>-1</sup> ) and dimerization gibbs free energies ( $\Delta G_{\text{dimer}}$ , kcal mol <sup>-1</sup> ) of some representative molecules.  | 89      |

## List of Tables:

| Chapter | Table  | Title   | Page No |
|---------|--------|---|---------|
| 3       | 3.2.8  | PBE0/6-31+G* calculated energies of the $\sigma$ -symmetric lone pair orbital ( $E_{\sigma}$ , in eV) concentrated at the carbene carbon of molecules <b>1–6</b> .  | 90      |
|         | 3.2.9  | PBE0/6-31+G* calculated energies of the $\sigma$ -symmetric lone pair orbital ( $E_{\sigma}$ , in eV) concentrated at the carbene carbon, carbonyl stretching frequencies ( $\nu_{\text{CO}}$ in $\text{cm}^{-1}$ ) for both symmetrical and unsymmetrical stretching mode and average carbonyl stretching frequencies ( $\nu_{\text{CO}(\text{avg})}$ , in $\text{cm}^{-1}$ ) of some representative L-Rh(CO) <sub>2</sub> Cl (L = <b>1Me</b> , <b>2PMe<sub>3</sub></b> , <b>3Me</b> , <b>4PMe<sub>3</sub></b> , <b>5Me</b> , <b>6PMe<sub>3</sub></b> ) complexes. | 91      |
|         | 3.2.10 | PBE0/6-31+G* calculated nucleophilicity index values of molecules <b>1–6</b> .  | 92      |
| 4       | 4.1    | Calculated values of key geometrical parameters for some representative molecules at different levels of theories.  | 108     |
|         | 4.2    | Calculated zero-point corrected energies of the parent and flipped isomers (in hartree) for some representative molecules and their energy differences ( $\Delta E$ , in $\text{kcal mol}^{-1}$ ) at different levels of theories.  | 109     |
|         | 4.3    | BP86/Def2-SVP, Def2-TZVP(Fe) Calculated values of energies (in hartree) for some representative molecules in their respective singlet and triplet spin states as well as the energy difference between singlet and triplet states ( $\Delta E_{\text{S-T}}$ , in $\text{kcal mol}^{-1}$ ).  | 109     |
|         | 4.4    | BP86/Def2-SVP, Def2-TZVP(Fe) calculated zero point corrected energies (in hartree) of the rotated and unrotated isomers with chelated 5NHCs as well as their energy differences ( $\Delta E_{\text{rot-unrot}}$ , in $\text{kcal mol}^{-1}$ ).  | 113     |
|         | 4.5    | BP86/Def2-SVP, Def2-TZVP(Fe) calculated zero point corrected energies (in hartree) of the rotated and unrotated isomers of <b>2<sub>x</sub>-6NHC</b> , <b>2'<sub>x</sub>-6NHC</b> , <b>2<sub>x</sub>-a<sub>NHC</sub></b> , <b>2'<sub>x</sub>-a<sub>NHC</sub></b> , <b>2<sub>x</sub>-BNHC</b> and <b>2'<sub>x</sub>-BNHC</b> complexes as well as their energy differences ( $\Delta E_{\text{rot-unrot}}$ , in $\text{kcal mol}^{-1}$ ).  | 114     |

## List of Tables:

| Chapter | Table | Title   | Page No |
|---------|-------|---|---------|
| 4       | 4.6   | BP86/Def2-SVP, Def2-TZVP(Fe) calculated zero point corrected energies (in hartree) of the rotated and unrotated isomers with chelated CAACs as well as their energy differences ( $\Delta E_{\text{rot-unrot}}$ , in kcal mol <sup>-1</sup> ). Negative sign indicates unrotated isomer is more stable.   | 115     |
| 5       | 5.1   | M062X/Def2-TZVP calculated values of key geometrical parameters for both the complexes <b>[1]</b> <sup>0</sup> and <b>[2]</b> <sup>0</sup> and the intermediates involved in the catalytic cycle. The iron centers are denoted by Fe <sub>p</sub> and Fe <sub>d</sub> whereas the apical and bridging carbonyl groups are indicated by CO <sub>ap</sub> and CO <sub>bridg</sub> respectively. | 133     |
|         | 5.2   | Calculated values of Mössbauer isomer shift ( $\delta$ ) and Mulliken spin densities ( $\rho$ ).  | 140     |
|         | 5.3   | BP86/Def2-TZVP calculated values of reduction potentials of the reduction events for both <b>[1]</b> <sup>0</sup> and <b>[2]</b> <sup>0</sup> .   | 141     |
|         | 5.4   | M062X/Def2-TZVP calculated $\Delta pK_a$ values for the protonated species obtained from different protonation pathways.  | 142     |

## List of Figures:

| Chapter | Figure | Title   | Page No |
|---------|--------|---|---------|
| 2       | 2.1    | Optimized singlet state geometries for the mesoionic NHCs ( <b>1–18</b> ).  | 30      |
|         | 2.2    | Optimized singlet state geometries for the six-membered remote NHCs ( <b>19–34</b> )  | 32      |
|         | 2.3    | Correlation plot between hydrogenation energies ( $E_{\text{hydro}}$ ) and singlet-triplet gaps ( $\Delta E_{\text{S-T}}$ ) of six-membered rNHCs.  | 37      |
|         | 2.4    | Plot of the energies of $\sigma$ -symmetric lone pair orbitals ( $E_{\sigma}$ ) and $\pi$ -symmetric unoccupied orbitals ( $E_{\pi^*}$ ) concentrated on the central carbon atom of (a) five-membered rNHCs ( <b>1, 3-18</b> ) and (b) six-membered rNHCs ( <b>19-23, 28-34</b> ).                            | 38      |
|         | 2.5    | Correlation plot between the energies of the $\pi$ -symmetric unoccupied MO ( $E_{\pi^*}$ ) and the values of natural charge at carbenic carbon ( $q(\text{C}_c)$ ) of five-membered rNHCs (omitting the point corresponding to <b>10</b> ).  | 41      |
|         | 2.6    | Correlation plot between the energies of the $\pi$ -symmetric unoccupied MO ( $E_{\pi^*}$ ) and the values of natural charge at carbenic carbon ( $q(\text{C}_c)$ ) of six-membered rNHCs (omitting the points corresponding to <b>29</b> and <b>32</b> ).  | 42      |
|         | 2.7    | Correlation plot between the energies of the $\sigma$ -symmetric lone pair orbital ( $E_{\sigma}$ ) and the values of nucleophilicity index ( $N$ ) of five-membered rNHCs.   | 43      |
|         | 2.8    | Correlation plot between the (a) energies of the $\sigma$ -symmetric lone pair orbital ( $E_{\sigma}$ ) and nucleophilicity index values ( $N$ ) and (b) energies of the $\pi$ -symmetric unoccupied molecular orbital ( $E_{\pi^*}$ ) and electrophilicity index values ( $\omega$ ) for six-membered rNHCs. | 45      |
|         | 2.9    | Optimized geometry and key bond lengths (in Å) of the <b>1</b> -PPh adduct.   | 46      |
|         | 2.10   | Correlation plot between the energy of the $\pi$ -symmetric unoccupied MO ( $E_{\pi^*}$ ) and $^{31}\text{P}$ chemical shifts ( $\delta^{31}\text{P}$ ) of the phosphinidene adducts of (a) five- and (b) six-membered rNHC   | 49      |

## List of Figures:

| Chapter | Figure | Title   | Page No |
|---------|--------|---|---------|
| 3       | 3.1.1  | Optimized singlet state geometries of all the carbenes ( <b>1–22</b> ).   | 62      |
|         | 3.1.2  | Correlation plot between calculated singlet-triplet separations ( $\Delta E_{S-T}$ ) and stabilization energies (SE) of <b>1–22</b> .   | 66      |
|         | 3.1.3  | Plot of the energies of the $\sigma$ -symmetric lone-pair orbitals ( $E_\sigma$ , in eV) concentrated at the central carbon atom of <b>1–22</b> .   | 66      |
|         | 3.1.4  | Correlation plot between the energies of the $\sigma$ -symmetric lone-pair orbitals concentrated at the carbene center ( $E_\sigma$ , in eV) and percentage of s-character of the lone pair of <b>1–22</b> .  | 68      |
|         | 3.1.5  | Correlation plot between the energies of the $\sigma$ -symmetric lone-pair orbitals ( $E_\sigma$ , in eV) and proton affinity (PA, kcal mol <sup>-1</sup> ) values of <b>1–22</b> .   | 69      |
|         | 3.1.6  | Correlation plot between the energies of the $\sigma$ -symmetric lone pair orbitals concentrated at the central carbene carbon atom ( $E_\sigma$ , in eV) and the $\nu_{CO(\text{avg})}$ values (in cm <sup>-1</sup> ) of the L–Rh(CO) <sub>2</sub> Cl complexes (L: <b>1–22</b> ).   | 70      |
|         | 3.1.7  | Correlation plot between energies of the $\sigma$ -symmetric lone pair orbitals ( $E_\sigma$ , in eV) and nucleophilicity indices ( $N$ , in eV) for <b>1–22</b> .  | 72      |
|         | 3.1.8  | Correlation plots between (a) the pyramidalization angle at the gallium atom ( $\theta_{\text{Ga}}$ , in degrees) and the nucleophilicity indices ( $N$ , in eV), and (b) the Ga–C <sub>C</sub> bond dissociation energies (BDE, in kcalmol <sup>-1</sup> ) and the Ga–C <sub>C</sub> bond lengths (in Å) of the GaCl <sub>3</sub> adducts of <b>1–22</b> . | 73      |
|         | 3.1.9  | Correlation plot between the calculated proton affinities (kcal mol <sup>-1</sup> ) and pyramidalization angle at gallium atom ( $\theta_{\text{Ga}}$ ) of the GaCl <sub>3</sub> adducts of <b>1–22</b> .   | 73      |
|         | 3.2.1  | Optimized singlet-state geometries of all the remote carbenes ( <b>1–6</b> ).   | 79      |

## List of Figures:

| Chapter | Figure | Title  | Page No   |
|---------|--------|--|---|
| 3       | 3.2.2  | (a) Contour plots of molecular orbitals showing cyclic delocalization of electrons within the five atom ring of <b>6PMe<sub>3</sub></b> and (b) Contour line diagram of the Laplacian of electron density in the ring plane of <b>6PMe<sub>3</sub></b> . Solid blue lines indicate regions of charge depletion [ $\nabla^2\rho(r)>0$ ] and dashed maroon lines indicate regions of charge concentration [ $\nabla^2\rho(r)<0$ ]. Green and red spheres denote bond critical points (bcp) and ring critical points (rcp), respectively.                         | 83  |
|         | 3.2.3  | (a) Contour plots of molecular orbitals showing cyclic delocalization of electrons within the five atom ring of <b>6P<sup>i</sup>Pr<sub>3</sub></b> and (b) Contour line diagram of the Laplacian of electron density in the ring plane of <b>6P<sup>i</sup>Pr<sub>3</sub></b> . Solid blue lines indicate regions of charge depletion [ $\nabla^2\rho(r)>0$ ] and dashed maroon lines indicate regions of charge concentration [ $\nabla^2\rho(r)<0$ ]. Green and red spheres denote bond critical points (bcp) and ring critical points (rcp), respectively. | 84  |
|         | 3.2.4  | Contour plots showing cyclic delocalization of electrons involving all the five ring atoms of <b>2PMe<sub>3</sub></b> , (a) top view (b) side view (PMe <sub>3</sub> groups are omitted for clarity).  | 86  |
|         | 3.2.5  | Graphical representation of singlet-triplet separations ( $\Delta E_{S-T}$ , kcal mol <sup>-1</sup> ) and hydrogenation energies ( $E_{\text{hydro}}$ , kcal mol <sup>-1</sup> ) of rNHCs <b>1–6</b> .   | 88  |
|         | 3.2.6  | Correlation plot between energies of the $\sigma$ -symmetric lone pair orbitals ( $E_{\sigma}$ , in eV) concentrated at the carbene carbon and nucleophilicity index ( $N$ ) values for rNHCs <b>1</b> and <b>3–6</b> .  | 92  |
|         | 4      | 4.1  | Optimized rotated structures for the model complexes with chelated 5NHCs. |
| 4.2     |        | Figure showing the metallacycle formed at Fe <sub>p</sub> as a result of coordination of the chelated 5NHC rings in complex <b>2H-5NHC</b> .   | 112   |
| 4.3     |        | Molecular orbital representing Fe <sub>p</sub> -Fe <sub>d</sub> bonding interaction in (a) <b>1H-5NHC</b> and (b) <b>2H-5NHC</b> .   | 112   |

## List of Figures:

| Chapter | Figure | Title   | Page No |
|---------|--------|---|---------|
| 4       | 4.4    | Molecular orbital representing the overlap between the lone pairs of CAAC and metal d-orbitals in (a) <b>1H-CAAC</b> and (b) <b>2H-CAAC</b> .   | 117     |
|         | 4.5    | Molecular orbital showing the interaction between the proximal iron center ( $\text{Fe}_p$ ) and bridging carbonyl group in <b>2H-CAAC</b> .  | 118     |
|         | 4.6    | Molecular orbitals representing the stepwise change of $\text{Fe}_p$ - $\text{Fe}_d$ bonding interaction as well as formation of new $\text{Fe}_p$ - $\text{CO}_{\text{bridg}}$ bond during rotation of $\text{Fe}_d(\text{CO})_3$ unit in <b>2H-CAAC</b> . The values represented by the arrows are natural charges at $\text{Fe}_p$ (left) and $\text{Fe}_d$ (right). $\theta_{rot}$ ( $\angle \text{C}_{\text{CO}}\text{-Fe}_p\text{-Fe}_d\text{-C}_{\text{CO}}$ ) represents the dihedral angle reflecting rotation at $\text{Fe}_d(\text{CO})_3$ unit ( $\text{C}_{\text{CO}}$ represents carbon atom of the carbonyl group). The values of $\text{Fe}_p$ - $\text{Fe}_d$ bond lengths and $\theta_{rot}$ are given in Å and degrees respectively. | 118     |
| 5       | 5.1    | Lowest Unoccupied Molecular Orbital (LUMO) of <b>[1-<i>t</i>H]<sup>+</sup></b> .  | 132     |
|         | 5.2    | Proposed mechanism for production of $\text{H}_2$ using <b>[1]<sup>0</sup></b> (phenyl groups at the nitrogen atoms of CAAC are omitted for clarity.)   | 136     |
|         | 5.3    | Proposed mechanism for production of $\text{H}_2$ using <b>[2]<sup>0</sup></b>  | 138     |



## List of Schemes:

| Chapter  | Scheme | Title   | Page No |
|----------|--------|---|---------|
| <b>1</b> | 1.1    | Schematic representation of the mesoionic ( <b>1–18</b> ) and remote NHCs ( <b>19–34</b> ) considered in this study.  | 12      |
|          | 1.2    | Schematic representation of the range of carbenes considered in this study (Dipp: 2,6-diisopropylphenyl).   | 14      |
|          | 1.3    | Schematic representation of the range of cyclic carbenes considered in this study.  | 15      |
|          | 1.4    | Schematic representation of the active site structure of [FeFe]-hydrogenase.  | 15      |
|          | 1.5    | Schematic representation of the molecules considered in the present study (X= H/Cyclohexyl).  | 16      |
|          | 1.6    | Schematic representation of the catalytic cycle of H <sub>2</sub> production catalyzed by [ <b>1</b> ] <sup>0</sup> .   | 17      |
|          | 1.7    | Schematic representation of the catalytic cycle of H <sub>2</sub> production catalyzed by [ <b>2</b> ] <sup>0</sup> .   | 18      |
| <b>2</b> | 2.1    | Schematic representation of the mesoionic ( <b>1–18</b> ) and remote NHCs ( <b>19–34</b> ) considered in this study.  | 28      |
|          | 2.2    | Resonance forms of carbene-phosphinidene adducts. Resonance form <b>A</b> dominates over <b>B</b> with increasing $\pi$ acidity of carbenes. <b>C</b> represents the orbital interaction involved in back donation from the phosphorus center to the formally vacant p orbital at the carbenic carbon atom. | 46      |
| <b>3</b> | 3.1.1  | Schematic representation of the range of carbenes considered in this study (Dipp: 2,6-diisopropylphenyl).   | 58      |
|          | 3.1.2  | Schematic representation of the back-bonding interaction between the metal center and the CO antibonding ( $\pi^*$ ) orbital.   | 69      |
|          | 3.2.1  | Schematic representation of experimentally known aminoylidecarbenes   | 76      |

## List of Schemes:

| Chapter | Scheme | Title   | Page No |
|---------|--------|---|---------|
| 3       | 3.2.2  | Range of molecules considered for this study.   | 76      |
| 4       | 4.1    | Schematic structures for the active site of [FeFe]-hydrogenase, left: oxidized or mixed valence state ( $H_{ox}$ ), right: reduced state ( $H_{red}$ ).                                 | 104     |
|         | 4.2    | Crystallographically characterized structures for the model complexes with inverted square pyramidal geometry at one Fe center in the $Fe^I$ - $Fe^I$ state.                            | 105     |
|         | 4.3    | Schematic representation of the molecules considered in the present study ( $X=H/Cyclohexyl$ )  | 106     |
|         | 4.4    | Schematic representations of the difference in orientation of the chelated 5NHC ligands in (a) $[(\mu-pdt)[Fe(Che-5NHC)(CO)_4]$ , (b) $2_X-5NHC$ and (c) $2'_X-5NHC$ , where $X=H/Cy$ . | 111     |
| 5       | 5.1    | Schematic representation of the active site of [FeFe]-hydrogenase.  | 128     |
|         | 5.2    | Schematic representation of the model complexes used as catalyst in the present study for production of dihydrogen.   | 129     |
|         | 5.3    | Schematic representation of the catalytic cycle of $H_2$ production catalyzed by $[1]^0$ .  | 132     |
|         | 5.4    | Schematic representation of the catalytic cycle of $H_2$ production catalyzed by $[2]^0$ .  | 134     |
|         | 5.5    | Schematic depiction of different protonation pathways considered in this study for complex $[1]^0$ . Similar pathways are also considered for $[2]^0$ .                                 | 142     |