

List of Figures

| | | |
|-----|---|----|
| 1.1 | A schematic representation of the cosmic dust life cycle | 3 |
| 1.2 | Average Milky Way and Magellanic cloud extinction curves | 5 |
| 1.3 | An interplanetary dust particle | 8 |
| 1.4 | <i>GALEX</i> spacecraft before launch | 11 |
| 1.5 | <i>Spitzer</i> telescope before launch | 12 |
| 1.6 | <i>AKARI</i> telescope instruments | 13 |
| 1.7 | Aperture photometry technique | 15 |
| 2.1 | FUV all-sky map showing our dust locations in the MW | 23 |
| 2.2 | 8 μm vs. N(H) and FUV intensity at low MW latitude | 26 |
| 2.3 | 24 μm vs. N(H) and FUV intensity at low MW latitude | 27 |
| 2.4 | 8 μm vs. N(H) and FUV intensity at high MW latitude | 28 |
| 2.5 | 24 μm vs. N(H) and FUV intensity at high MW latitude | 29 |
| 2.6 | DIRBE maps explaining our observed correlations | 31 |
| 3.1 | Observed dust locations around N11 and 30D in the LMC | 35 |
| 3.2 | FUV vs. all <i>Spitzer</i> IR intensities in the LMC | 38 |
| 3.3 | FUV vs. N11 IR intensities in the LMC | 39 |
| 3.4 | FUV vs. all <i>AKARI</i> IR intensities in the LMC | 40 |
| 3.5 | FUV vs. 30D IR intensities in the LMC | 41 |
| 4.1 | Observed diffuse dust locations in the SMC | 48 |
| 4.2 | 8 μm vs. FUV intensities in the SMC | 54 |

| | | |
|-----|--|----|
| 4.3 | 22 μm vs. FUV intensities in the SMC | 55 |
| 4.4 | 65 μm vs. FUV intensities in the SMC | 56 |
| 4.5 | 90 μm vs. FUV intensities in the SMC | 57 |
| 4.6 | MIR vs. FIR (65 μm) intensities in the SMC | 58 |
| 4.7 | MIR vs. FIR (65, 90 μm) intensities in the SMC | 59 |
| 5.1 | Observed dust locations around Orion center | 65 |
| 5.2 | FUV vs. FIR intensities in Orion | 70 |
| 5.3 | 3D dust distribution observed towards Orion | 73 |
| 6.1 | Oblate spheroidal composite dust grain model | 84 |
| 6.2 | Best fit grain model to star MWC 1080A | 86 |
| 6.3 | Best fit grain model to star MWC 297 | 87 |
| 6.4 | Best fit grain model to star HL Tau | 88 |
| 6.5 | Best fit grain model to star SVS13 | 92 |

List of Tables

| | | |
|-----|--|----|
| 2.1 | Low latitude rank correlation coefficients in the MW | 25 |
| 2.2 | High latitude rank correlation coefficients in the MW | 25 |
| 3.1 | Details of <i>Spitzer</i> observations in the LMC | 36 |
| 3.2 | Details of <i>AKARI</i> observations in the LMC | 37 |
| 3.3 | Rank correlations for FUV vs. <i>Spitzer</i> IR in the LMC | 38 |
| 3.4 | Rank correlations for FUV vs. <i>AKARI</i> IR in the LMC | 39 |
| 4.1 | Flux details for 21 locations observed by <i>IUE</i> | 50 |
| 4.2 | IR intensities for 24 locations convolved with <i>FUSE</i> | 51 |
| 4.3 | IR intensities for 24 locations convolved with <i>IUE</i> | 52 |
| 4.4 | FUV vs. IR rank correlations in the SMC | 53 |
| 4.5 | MIR vs. FIR rank correlations in the SMC | 58 |
| 5.1 | Central stars contributing to emission in Orion | 66 |
| 5.2 | Details of <i>GALEX</i> FUV observations around Orion | 68 |
| 5.3 | Details of <i>AKARI</i> FIR observations around Orion | 69 |
| 5.4 | FUV vs. FIR correlations in Orion | 71 |
| 5.5 | Best fit model parameters for dust around Orion | 72 |
| 6.1 | Sample of young stars | 80 |
| 6.2 | Measured polarization angles | 81 |
| 6.3 | DDA best fit parameters | 89 |

| | |
|--|----|
| 6.4 EMA T-Matrix best fit parameters | 89 |
|--|----|