

# List of Figures

---

<b>Figure No.</b>		<b>Page No.</b>
<b>CHAPTER 2</b>		
2.1	ER $\alpha$ structure and phosphorylation sites. ER $\alpha$ consists of six structural domains that contain a number of functional domains: Activation function-1 (AF-1), DNA-binding domain (DBD), hinge, ligand binding domain (LBD), dimerization, activation function-2 (AF-2), and the F domain considered to repress ER $\alpha$ function.	9
2.2	Mechanisms of estrogen receptor (ER) action in breast cancer	9
2.3	Structure of conserved protein kinase core	14
2.4	(A) Protein kinase A, representing activation loop in yellow colour; (B) Representation of activation loop showing DFG and APE motif.	15
2.5	Active and inactive conformation of kinase	16
2.6	The structure of LMTK family: The LMTK family members (LMTK1A, LMTK2 and LMTK3) contain relatively conserved transmembrane domains and catalytic/kinase domains.	17
2.7	Sequence alignment of human LMTK1A, LMTK2 and LMTK3. The sequence similarity alignment was carried using Clustal Omega ( <a href="https://www.ebi.ac.uk/Tools/msa/clustalo/">https://www.ebi.ac.uk/Tools/msa/clustalo/</a> ) Similarity scores are represented by asterisk mark	19
2.8	Schematic representation of the order and disorder score in LMTK3. LMTK3 is composed of ordered domains that is a kinase domain, and variable intrinsically disordered domains depicted in red colour. Data obtained using DICHOT ( <a href="http://spock.genes.nig.ac.jp/~genome/DICHOT/">http://spock.genes.nig.ac.jp/~genome/DICHOT/</a> ).	21
2.9	LMTK3 positively controls ESR1 gene transcription, protein	24

# List of Figures

---

levels and phosphorylation, developing resistance against endocrine therapies. LMTK3 increases the transcriptional activation of ESR1 gene by inhibiting PKC activity, whose activity leads to the phosphorylation of AKT, which in turn phosphorylates FOXO3 and degrades. LMTK3 also directly phosphorylates ER $\alpha$  protein, which protects ER $\alpha$  from ubiquitin (Ub)-mediated proteasomal degradation.

- |      |  |    |
|------|--|----|
| 2.10 | LMTK3 promotes invasion and metastasis in Breast cancer through GRB2-Mediated induction of Integrin $\beta$ 1. | 26 |
| 2.11 | Structure of GRB2 (PDB ID: 1GRI) visualized in Chimera   | 27 |

## CHAPTER 3

- |     |  |    |
|-----|--|----|
| 3.1 | Schematic representation of the interactions that contribute to the potential energy function for MD simulation.   | 36 |
| 3.2 | Periodic boundary conditions in two dimensions. The simulation cell (solid) is surrounded by translated copies of itself (dashed).   | 37 |
| 3.3 | Schematic representation of TIP3P water model.   | 41 |
| 3.4 | Flowchart showing the steps involved in MD Simulation.   | 41 |
| 3.5 | The process of energy minimization changes the geometry of the molecule in a step-wise fashion until a minimum is reached.   | 43 |
| 3.6 | A line search is used to locate the minimum in the function in the direction of the gradient.  | 44 |
| 3.7 | Computational schemes of the binding free energies based on MM-PBSA/GBSA. The free energies colored in black are directly calculated, while the free energy of interest colored in blue is indirectly did using the thermodynamic cycle of other free energies | 53 |

## List of Figures

---

- 3.8 Viewing Grids in AutoDockTools. The protein is shown on the left in white bonds, and the grid box is shown on the right side. The blue contours surround areas in the box that are most favorable for binding of carbon atoms, and the red contour areas that favor oxygen atoms. A ligand is shown inside the box at upper right 57
- 3.9 (A) Surface topology graphs for trypsin inhibitor (PDB code: 1ba7). The caps, belts and pits are connected with edges. (B) Geometric patches: the patches are in light colors and the protein is dark. 60
- 3.10 The PatchDock user interface: The receptor molecule and the ligand molecule are given either by the PDB code of the molecule (chain IDs are optional) or by uploading a file in PDB format 61
- 3.11 The solutions page presents the geometric score, interface area size and desolvation energy of the 20 top scoring solutions 62
- 3.12 Representation of the ClusPro algorithm, the number of structures retained after each step is shown in a blue box 63
- 3.13 Schematic diagrams from the PDBsum for entry 1a5z: (A) The 'wiring diagram' shows the protein's secondary structure elements ( $\alpha$ -helices and  $\beta$ -sheets) together with  $\beta$ - and  $\gamma$ -turns, and  $\beta$ -hairpins. The yellow linking bars labelled 1 and 2 represent disulphide bonds. The single-letter amino acid codes showing the protein's sequence are coloured red or blue depending on whether they belong to CATH [251] structural domain. Red dots above the single-letter codes signify residues that interact with any bound ligand(s) while coloured lines underneath represent residues belonging to a PROSITE pattern, the redder the colour the more highly conserved the residue in the pattern. (B) 64

# List of Figures

---

- Topology diagram illustrates the  $\beta$ -strands by the large arrows joined side-by-side (pink colour), forming central  $\beta$ -sheet. The  $\alpha$ -helices represented by the red cylinders. The small arrows indicate the directionality of the protein chain, from the N- to the C-terminus.
- 3.14 Protein–protein interaction diagrams in PDBsum for PDB entry 1mmo: (A) Thumbnail image of the 3D structural model which contains six protein chains (B) Schematic diagram showing the interactions between the chains. The area of each circle is proportional to the surface area of the corresponding protein chain. The joining lines are coloured light blue for hydrogen bonds and orange for non-bonded contacts. (C) A schematic diagram showing the numbers of interactions across one of the interfaces, namely the B–G protein interface, and the numbers of residues involved. (D) Detail of the individual residue–residue interactions across this interface. Hydrogen bonds (blue lines), non-bonded contacts (orange tick-marks), and salt bridges (red lines) between residues on either side of the protein-protein interface. 66
- 3.15 PDBsum interaction plot for PDB entry 5trd in LIGPLOT (A) LIGPLOT diagram showing the protein residues that interact with the CTP (cytidine-5'-triphosphate) ligand, with hydrogen bonds shown by the green dashed lines and non-bonded contacts by the brown rays, and; (B) as in (A), but for residues interacting with the bound sodium ion; (C) Diagram of protein-DNA interactions, with H-bonds as blue dashed lines and non-bonded contacts as brown dashed lines 67

## CHAPTER 4

- 4.1 3D structure of LMTK3 domain obtained from I-TASSER 78

## List of Figures

---

4.2	Representation of superimposed LMTK3 domain model with Insulin receptor (InsR)	78
4.3	Ramachandran plot of LMTK3 domain: 87% of residues are in most favored region (243 residues) and 5.1% in disallowed region	79
4.4	Quality index for the model structure of LMTK3 domain by ProSA server. The index value was represented by Z-score	79
4.5	Verify 3D plot of the modelled structure of LMTK3 domain	80
4.6	Superimposed structure of modelled LMTK3 domain with PDK1 protein	80
4.7	Architecture of LMTK3 domain and PDK1 (pdb ID:3HRF) in cartoon representation. The three panels use the following color scheme: $\alpha$ C helix (red), P-loop: glycine rich (green), activation loop (magenta).	81
4.8	Multiple sequence alignment of LMTK3 domain with different kinases	82
4.9	Structural comparison of LMTK3 domain with different active kinases	83
4.10	Structural comparison of LMTK3 domain with different inactive kinases	83
4.11	RMSD plot of LMTK3 domain for C $\alpha$ atoms as a function of time	84
4.12	Rg plot of LMTK3 domain as a function of time	85
4.13	B-factor and RMSF plot showing flexibility in loop region in LMTK3 as a function of residue index	85

## List of Figures

---

4.14	SASA of LMTK3 domain plot as a function of time	86
4.15	DSSP plot: secondary structure variation of each residue as a function of simulation time period.	87
4.16	Equilibrated structure of LMTK3 domain with the percentage of secondary structure content	87
4.17	Secondary structure topology, $\beta$ sheet topology and hydrogen bond pattern of LMTK3 domain. (A) Secondary structural elements ( $\beta$ sheets, $\beta$ hairpins, $\beta$ turns, strands, helices, helix–helix interactions, and $\gamma$ turns). (B) Topology diagram illustrating the $\beta$ strands, represented by the large arrows (grey), join up side-by-side. The figure also illustrates the relative locations of the helices (black cylinders). The small arrows specify the directionality of the protein chain, from the N- to the C-terminus. (C) Intra- and inter-hydrogen bond pattern between residues in $\beta$ sheet topology	88
4.18	Number of hydrophobic contacts in LMTK3 domain as a function of time	90
4.19	Hydrophobic plot analysis of LMTK3 domain	90
4.20	Hydrophobic cluster analysis of LMTK3 domain	91
4.21	(A) Hydrophobic patches as determined with Swiss pdb viewer for equilibrated structure of LMTK3 domain. Surface representation of hydrophobic patches shown in different colours along with area. (B) Electrostatic potential distributions on the LMTK3 domain surface from $-10.0 \text{ kcal mol}^{-1}$ (red) to $+10.0 \text{ kcal mol}^{-1}$ (blue)	92
4.22	Number of intra-molecular hydrogen bonds in LMTK3 domain as a function of time	92

# List of Figures

---

4.23	Five binding pockets in LMTK3 domain denoted as spheres from LIGSITE <sup>csc</sup> server	95
4.24	LMTK3-ATP interaction (A) LMTK3-ATP complex with least binding free energy (-1.71 k cal mol <sup>-1</sup> ), ATP is shown in red sphere form (B) Ligplot+ analysis: Residues of LMTK3 involved in the formation of hydrogen bonds (green colour) and hydrophobic contacts are represented by an arc with spokes radiating towards the ATP atoms they contact.	96

## CHAPTER 5

5.1	ER $\alpha$ structure and phosphorylation sites: Activation function-1 (AF-1), DNA-binding domain (DBD), hinge, ligand binding domain (LBD), dimerization, and activation function-2 (AF-2) domain	100
5.2	Modelled structure of ER $\alpha$	102
5.3	Quality index for the model structure of ER $\alpha$ generated by ProSA server. The index value was represented by Z-score of -7.23.	105
5.4	ER $\alpha$ -LMTK3 complex model structures obtained from patchdock server based on high geometric shape complementary score	106
5.5	Summary of the total number of residues involved in different types of interactions between ER $\alpha$ and LMTK3 domain	107
5.6	Summary of interacting residues that are involved in bonded and non-bonded interactions between ER $\alpha$ and LMTK3 interface.	108
5.7	Representation of LMTK3-ER $\alpha$ interaction: C-terminal lobe of LMTK3 interacting with N-terminal region of ER $\alpha$ at Ser104 and Ser106	109
5.8	Summary of the total number of residues involved in different	110

# List of Figures

---

	types of interactions between ER $\alpha$ and MAPK.	
5.9	Summary of interacting residues that are involved in bonded and non-bonded interactions between ER $\alpha$ and MAPK interface.	110
5.10	Root Mean Square Deviations (RMSD) of ER $\alpha$ and LMTK3 as a function of simulation time	111
5.11	Snapshots of ER $\alpha$ -LMTK3 complex at different time intervals of simulation	112
5.12	Temperature and energy of ER $\alpha$ -LMTK3 complex as a function of time	112
5.13	Intermolecular distance between ER $\alpha$ and LMTK3 domain as a function of time	113
5.14	Number of inter-molecular hydrogen bonds in ER $\alpha$ - LMTK3 complex model structure as a function of simulation time period. Here we have considered inter-molecular hydrogen bonds involving two cases: ER $\alpha$ as acceptor and LMTK3 as donor (black line) and the other one, ER $\alpha$ as donor and LMTK3 as acceptor (red line)	113

## CHAPTER 6

6.1	GRB2-LMTK3 docked complex from ClusPro	123
6.2	Representation of GRB2-LMTK3 complex, where SH2 domain is interacting with the C-lobe of LMTK3 domain	123
6.3	Activation loop (in red) of LMTK3 interacting with the GRB2	123
6.4	Number of interface residues, bonded and non-bonded contacts	124
6.5	Interface residues and their interactions at GRB2-LMTK3 interface	125

# List of Figures

---

6.6	Potential energy, kinetic energy and total energy of the GRB2-LMTK3 complex as a function of time	127
6.7	RMSD of GRB2-LMTK3 complex as a function of simulation time	128
6.8	Snapshots of GRB2-LMTK3 complex at different time intervals of MD simulation	128
6.9	Number of inter-molecular hydrogen bonds in GRB2-LMTK3 complex as a function of simulation time period. Here we have considered inter-molecular hydrogen bonds involved in two cases: GRB2 as acceptor and LMTK3 as donor (depicted in black) and the other one, GRB2 as donor and LMTK3 as acceptor (depicted in red).	129
6.10	(A) Representation of hot spot and null spot residues at GRB2 and LMTK3 interface. Red color representing the null-spot residues. Whereas the light blue and green color depicted as hot spots. (B) Representation of the interaction of hot spot (green and blue) residues in a form of spheres without secondary structure involved in GRB2-LMTK3 interface	131

## CHAPTER 7 (a)

7a.1	Superimposed structure of LMTK3 domain with ERK2	141
7a.2	Superimposed structure of LMTK3 domain with PKA	142
7a.3	CDK5-LMTK3 docked complex from PatchDock server	143
7a.4	CDK5-LMTK3 complex structures obtained from PatchDock, ranked based on: high geometric shape complementary score, surface area and atomic contact energy	143
7a.5	Summary of the total number of residues involved in different	144

# List of Figures

---

	types of interactions in CDK5-LMTK3 complex obtained from PatchDock based on high geometric shape complementary score.	
7a.6	(A) Number of interacting residues that are involved in bonded and non-bonded interactions between CDK5 and LMTK3 interface. (B) Representation of CDK5-LMTK3 complex: C terminal residues of CDK5 interact with activation segment of LMTK3 domain.	145
7a.7	CDK5-LMTK3 complex representing probable phosphorylation sites in LMTK3 activation loop	146
7a.8	CDK5-LMTK3 complex showing N and C terminal lobe of CDK5 interacting with LMTK3. The binding pocket for ATP analogue is present between these two lobes which is almost at the center of CDK5 and LMTK3 interface	147
7a.9	Temperature and energy plot of CDK5-LMTK3 complex as a function of simulation time	148
7a.10	RMSD of CDK5 and LMTK3 as a function of simulation time.	148
7a.11	Snapshots of CDK5-LMTK3 complex at different time intervals of MD simulation	149
7a.12	Number of inter-molecular hydrogen bonds in CDK5-LMTK3 complex model structure as a function of simulation time period. Here we have considered inter-molecular hydrogen bonds involved in two cases: LMTK3 as acceptor and CDK5 as donor (black line) and the other one, LMTK3 as donor and CDK5 as acceptor (red line)	150

## CHAPTER 7 (b)

7b.1	Snapshots of (A) unphosphorylated (B) phosphorylated (C) mutated LMTK3 domain	156
------	---	-----

## List of Figures

---

7b.2	RMSD plot of C $\alpha$ atom over time course of MD simulation for phosphorylated (red), unphosphorylated (black) and mutated (green) LMTK3 domain	157
7b.3	Loop region RMSD plot of C $\alpha$ atom over time course of MD simulation for phosphorylated (red), unphosphorylated (black) and mutated (green) LMTK3 domain	157
7b.4	Potential energy as a function of RMSD for loop region of LMTK3 domain in (A) unphosphorylated, (B) phosphorylated (C) mutated LMTK3 domain	158
7b.5	RMSF plot of C $\alpha$ atom over time course of MD simulation for phosphorylated (red), unphosphorylated (black) LMTK3 and mutated (green) LMTK3 domain	159
7b.6	Flexibility and conformational change in activation segment of (A) unphosphorylated (B) phosphorylated and (C) mutated LMTK3 domain during the course of MD simulation.	160
7b.7	Radius of gyration as a function of time course of simulation for unphosphorylated, phosphorylated and mutated LMTK3 domain.	160
7b.8	Potential energy plot as a function of time for unphosphorylated (black), phosphorylated (red) and mutated LMTK3 (green) domain	161
7b.9	Electrostatic interaction for (A) unphosphorylated (B) phosphorylated and (C) mutated LMTK3 domain as a function of time.	162

## CHAPTER 8

8.1	Flow chart of structure-based virtual screening of inhibitors obtained from ZINC database	169
-----	---	-----

## List of Figures

---

8.2	Chemical structures of ATP and lead compounds with zinc Iids	169
8.3	Molecular docking results of LMTK3 using Ligplot+: Hydrogen bonds are indicated by green lines between the atoms involved, while hydrophobic contacts are indicated by an arc with spokes radiating towards the ligand atoms they contact.	173
8.4	ATP binding pocket, where activation loop is in open conformation and form a pocket where ATP binds	174
8.5	Binding modes of LMTK3 with ATP and lead compounds: ligands are shown in red color, hydrogen bonds between ligand and protein denoted in navy blue color	177
8.6	Surface diagram of binding cavity of complexes: LMTK3-ATP and LMTK3- lead compounds	178
8.7	Superimposed LMTK3-ATP and inhibitor complex: (A) LMTK3 is DYG-D in conformation, open activation loop forming the binding pocket for ATP/inhibitors (B) Zoomed binding pocket, residues of activation loop protrude out and interacting with inhibitors	179
8.8	RMSD of ATP and six lead compounds to LMTK3 derived by molecular dynamics simulations	181
8.9	RMSF of ATP and six lead compounds to LMTK3 derived by molecular dynamics simulations	182
8.10	Number of intermolecular hydrogen bonds in LMTK3-ATP and LMTK3-lead compound complexes as a function of simulation time.	183
8.11	Unbinding process of ATP/lead compounds dissociating from the ATP binding site of LMTK3 domain	185

# List of Figures

---

## CHAPTER 9

9.1	The modelled structure of the kinase domain of LMTK3 obtained from I-TASSER server	194
9.2	The quality index for the model structure of LMTK3 domain generated by ProSA server. The index value is represented by Z-score	195
9.3	Verify 3D plot of the modelled structure of LMTK3 domain	196
9.4	Ramachandran plot of LMTK3 domain: 87.7% of the residues are in most favored regions, 8.7% of the residues are in allowed region and 3.6% of residues are in disallowed regions	196
9.5	The chemical structures of top three hit compounds with database IDs	198
9.6	E-pharmacophoric features for LMTK3 inhibitor	199
9.7	Superimposed binding pose of LMTK3-hit complexes	200
9.8	The binding mode of the three hits (BAS1294510, BDF24025570 and BAS01313675) with LMTK3 domain: Residues involved in the formation of hydrogen bonds (depicted in cyan color), hydrogen bonds are in red color, inhibitors are shown in ball and stick form. And the residues involved in hydrophobic interaction with the compounds are depicted in yellow color	201
9.9	LMTK3 domain complexed with top three hits visualized in Ligplot+: Hydrogen bonds with distance are indicated by green lines between the atoms involved, while hydrophobic contacts are represented by an arc with spokes radiating towards the ligand atoms they contact.	201

## List of Figures

---

9.10	Representation of inhibitors binding to a region distinct from the ATP binding site. Secondary structure shown in grey color, ATP, hit1, hit2, and hit3 depicted in red, green, blue and magenta colour (A-D) respectively. (E) Superimposed LMTK3-hit complexes with LMTK3-ATP complex	203
9.11	RMSD of apo LMTK3 domain and LMTK3-hit complexes as a function of time.	204
9.12	Rg plot of apo LMTK3 domain and LMTK3-hit complexes as a function of time	204
9.13	RMSF of apo LMTK3 domain and LMTK3-hit complexes as function of residue index	205
9.14	Graphical representation of per residue binding energy decomposition analysis of interacting residues of LMTK3 domain with (A) hit1 (B) hit2 and (C) hit3	210
9.15	Unbinding process of hits dissociating from the LMTK3 domain	211

# List of Tables

---

<b>Table No.</b>		<b>Page No.</b>
<b>CHAPTER 4</b>		
4.1	Solvent accessible surface area and Molecular accessible surface area of the modelled LMTK3 domain structure determined using Surface Racer software	86
4.2	Details of the five binding Pockets (determined using LIGSITE <sup>csc</sup> server) in the model structure of LMTK3 domain	93
4.3	List of residues along with the residue index involved in the formation of five binding pockets in human LMTK3 domain structure as determined by LIGSITE <sup>csc</sup> server	95
4.4	List of residues along with the residue index involved in the formation of five binding pockets in human LMTK3 domain structure as determined by LIGSITE <sup>csc</sup> server	95
<b>CHAPTER 5</b>		
5.1	Interface statistics for ER $\alpha$ -LMTK3 complex	107
5.2	Interface statistics for ER $\alpha$ -MAPK complex	111
<b>CHAPTER 6</b>		
6.1	Summary of interface statistics for GRB2-LMTK3 complex determined from PDBsum server	125
6.2	Total number of H-bonds involved in GRB2-LMTK3 interface identified from PDBsum.	126
6.3	Number of salt bridges at GRB2-LMTK3 interface obtained from PDBsum	127
6.4	Intermolecular H-bond occupancy of GRB2-LMTK3 complex from MD simulation	129
6.5	Comparison of results from six different computational methods used to detect probable hotspot residues at GRB2-LMTK3 interface	132
6.6	Binding free energy calculation of GRB2-LMTK3 complex using MM-GBSA method	133

# List of Tables

---

## CHAPTER 7 (a)

7a.1	Interface statistics for CDK5-LMTK3 complex	144
------	---	-----

## CHAPTER 8

8.1	The Autodock docking score and amino acids of LMTK3 involved in hydrogen bonding interactions with ATP and other lead-compounds	175
8.2	AutoDock results of ATP and lead compounds	176
8.3	Physico-chemical properties of the lead compounds	180
8.4	Binding free energy results of LMTK3-ATP/lead-compound complexes from MM-PBSA/GBSA calculations	183

## CHAPTER 9

9.1	Top 10 hits obtained from Glide XP docking	197
9.2	Glide XP docking scores for top 3 hits and the interacting amino acids of LMTK3	198
9.3	Features of the E-Pharmacophore Model	199
9.4	Summary of the Hydrogen bonding interactions of LMTK3 residues with top three hits.	202
9.5	MM-PBSA binding free energy calculation for LMTK3-inhibitor complexes	206
9.6	Decomposition of binding energy on per residues on LMTK3-inhibitor complexes in dynamic system.	208

# List of Abbreviations

---

Å	:	Angstrom
ACE	:	Atomic contact Energy
AF-1	:	Activation function-1
AFM	:	Atomic Force Microscopy
AIB1	:	Amplified in Breast Cancer
AMBER	:	Assisted Model Building with Energy Refinement
ATP	:	Adenosine Triphosphate
BPTI	:	Bovine Pancreatic Trypsin Inhibitor
CAPRI	:	Critical Assessment of Prediction of Interaction
COM	:	Centre of Mass
CDK5	:	Cyclin-dependent kinase 5
3-D	:	3-Dimensional
DBD	:	DNA-binding domain
DNA	:	Deoxyribonucleic Acid
EPKs	:	Eukaryotic protein kinases
ER	:	Estrogen receptor
FFT	:	Fast Fourier Transform
FTIR	:	Fourier-transform Infrared Spectroscopy
GAFF	:	General Amber force field
GBSA	:	Generalized Born generalized Born and surface area continuum solvation
GRB2	:	Growth factor receptor bound protein
HER2	:	Human epidermal growth factor receptor 2
HR	:	Hormone receptors
HTVS	:	High throughput virtual screening
IDP	:	Intrinsically Disordered Protein
KFC	:	Knowledge-based FADE and Contacts
LBD	:	Ligand binding domain
LGA	:	Lamarckian genetic algorithm
LMTK3	:	Lemur Tyrosine Kinase 3
LTP	:	Long-term Potentiation
I-	:	Iterative Threading ASSEMBLY Refinement
TASSER	:	
MD	:	Molecular Dynamics
MM	:	Molecular Mechanics
ns	:	nanosecond
NMR	:	Nuclear Magnetic Resonance Spectroscopy
ps	:	picosecond
PBC	:	Periodic boundary condition
PBSA	:	Poisson–Boltzmann surface area continuum solvation
PDB	:	Protein Data Bank
PKA	:	Protein kinase A
PME	:	Particle Mesh Ewald
PMF	:	Potential of Mean Force
PPIs	:	Protein-protein interaction
PR	:	Progesterone receptor
QM	:	Quantum Mechanics
RBVI	:	Resource for Biocomputing Visualization, and

# List of Abbreviations

---

		Informatics
RC	:	Reaction Coordinate
Rg	:	Radius of Gyration
RMSD	:	Root Mean Square Deviation
RMSF	:	Root Mean Square Fluctuation
RTKs	:	Receptor tyrosine kinases
SH	:	SRC homology
SP	:	Standard precision
SPC	:	Simple Point Charge
SRF	:	Serum response factor
TIP3P	:	Transferable Intermolecular Potential Three-point
US	:	Umbrella Sampling
VMD	:	Visual Molecular Dynamics
WHAM	:	Weighted Histogram Analysis Method
XP	:	Extra precision