

## LIST OF TABLES

Chapter	Table	Title	Page No.
2	2.1	List of chemicals	36
3	3.1	Comparison of the ORR performance parameters of Pd <sub>3</sub> Fe <sub>0.5</sub> Cu <sub>0.5</sub> /C with recently reported some seminal works of Pd-based electrocatalysts in 0.1 M KOH solution.	68-69
6	6A.1	Comparison of k <sub>app</sub> of Cu, Fe <sub>2</sub> O <sub>3</sub> and Cu/Fe <sub>2</sub> O <sub>3</sub> for the reduction of 4-NP	140
	6A.2	Optimization of Reaction Condition for the Reduction of 4-NP	141
	6A.3	Catalytic performances of the Cu/Fe <sub>2</sub> O <sub>3</sub> NPs for reduction of various nitroaromatics using NaBH <sub>4</sub> .	142-143
	6A.4	Reusability Studies of the Cu/Fe <sub>2</sub> O <sub>3</sub> NPs upto 5 <sup>th</sup> cycles.	146
	6B.1	Catalytic performances of the CuNi NPs for reduction of various nitroaromatics using 2-propanol KOH as base	151-152
	6B.2	Recyclability test of the reduction of 4-iodo 1-nitrobenzene with CuNi catalyst system	153
	6B.3	Comparison table for previously reported catalytic system for the reduction of nitrobenzene through transfer hydrogenation	155

## LIST OF FIGURES

Chapter	Figure	Title	Page No.
1	1.1	The Lycurgus cup in reflected light (left) and in transmitted light (right)	2
	1.2	Five original bottled samples prepared by Micheal Faraday	3
	1.3	Various types of carbon based materials.	5
	1.4	Some of the core areas of applications of metal NPs.	8
	1.5	Schematic presentation of proton exchange membrane fuel cell	15
	1.6	(a) SEM image of Pd/N-G, (b, c) Low and high resolution TEM images of Pd/N-G. Inset in (b) shows the diffraction pattern of Pd/N-G and the scale bar corresponds to 5 nm <sup>-1</sup> . Inset in (c) shows the histogram of the Pd NPs distribution in Pd/N-G electrocatalyst. (d) EDX spectra of Pd/N-G.	17
	1.7	ORR on Pd-CN <sub>x</sub> composite in 0.5 M KOH medium: (a) CVs of Pd-CN <sub>x</sub> composite in O <sub>2</sub> and N <sub>2</sub> -saturated 0.5 M KOH solution with scan rate 100 mV/s, (b) Comparison of LSV curves of porous Pd-CN <sub>x</sub> , Pt/C and Pd/C modified GC electrode in O <sub>2</sub> -saturated 0.5 M KOH with 1600 rpm rotation at 100 mV/s scan rate, (c) LSV curves of Pd-CN <sub>x</sub> modified electrode in O <sub>2</sub> -saturated 0.5 M KOH solution with rotating speed varying from 600 rpm to 2700 rpm, (d) The corresponding K-L plots at different potentials, (e) Mass transfer corrected Tafel slope of Pd-CN <sub>x</sub> composite in comparison with Pt/C and Pd/C catalyst in basic medium, (f) The mass activity and specific activity of Pd-CN <sub>x</sub> , Pt/C and Pd/C modified	18

		electrode at different potentials, (g) Steady state chronoamperometric response of Pd-CN <sub>x</sub> composite, Pt/C and Pd/C at a constant potential of 0.7 V.	
1.8		XRD pattern of different Cu-compounds (a); UV-visible spectra of Cu <sub>2</sub> Cl(OH) <sub>3</sub> (b); TEM image of in-situ generated Cu NPs (c); particle size distribution of Cu NPs (d); UV-visible absorption spectra (e) of 4-nitrophenol before (black line) and after (red line) addition of NaBH <sub>4</sub> and 4-aminophenol (blue line) and plot of ln(A <sub>t</sub> /A <sub>0</sub> ) against reaction time (f) derived from absorption spectra (inset) of 4-nitrophenol reduction in presence of in-situ Cu nanocatalyst.	20
1.9		Time dependent UV-visible absorption spectra over CuNi (a) and CuNi/Co <sub>3</sub> O <sub>4</sub> (b); plots of ln (A <sub>t</sub> /A <sub>0</sub> ) against reaction time derived from absorption spectra (c) and recyclability test over CuNi/Co <sub>3</sub> O <sub>4</sub> nanocatalyst for the 4-NP reduction, respectively.	21
2	2.1	Schematic illustration of Bragg's law	41
3	3.1	EDS elemental mapping of Pd <sub>3</sub> Fe <sub>0.5</sub> Cu <sub>0.5</sub> /C NPs.	53
	3.2	TGA profile of Pd <sub>3</sub> Fe <sub>0.5</sub> Cu <sub>0.5</sub> /C NPs in air atmosphere	53
	3.3	(a) XRD patterns of the Pd <sub>3</sub> Fe <sub>0.5</sub> Cu <sub>0.5</sub> /C, Pd <sub>3</sub> Fe/C, PdFe/C, and PdFe <sub>3</sub> /C electrocatalysts, and (b) N <sub>2</sub> adsorption-desorption isotherm of Pd <sub>3</sub> Fe <sub>0.5</sub> Cu <sub>0.5</sub> /C. Inset of (b) shows the corresponding pore size distribution	55
	3.4	Typical (a) low, (b,c) medium, and (d-f,h) high resolution-TEM images of Pd <sub>3</sub> Fe <sub>0.5</sub> Cu <sub>0.5</sub> /C, (g) corresponding SAED pattern, (e) is magnified HR-TEM image taken from (d), marked by the rectangle.	55
	3.5	TEM and HR-TEM images of Pd <sub>3</sub> Fe/C NPs.	56
	3.6	(a) C 1s, (b) Pd 3d, (c) Fe 2p, and (d) Cu 2p XP spectra of Pd <sub>3</sub> Fe <sub>0.5</sub> Cu <sub>0.5</sub> /C.	58

- 3.7 The core-level XP spectrum of (a) Pd 3d and (b) Fe 2p of Pd<sub>3</sub>Fe/C NPs. 58
- 3.8 (a) CV plots of Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C, N<sub>2</sub>- and O<sub>2</sub>-saturated 0.1 M KOH solution with a scan rate of 50 mV s<sup>-1</sup>, (b) Rotating rate-dependent ORR polarization curves for respective NPs with the scan rate of 10 mV s<sup>-1</sup>, (c) K-L plots of  $j^{-1}$  vs.  $\omega^{-1/2}$  for Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C NPs at different potential obtained from (b), and (d) The plot of the number of transferred electrons vs. potential for Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C NPs. 59
- 3.9 Electrochemical data for ORR over PdFe<sub>3</sub>/C NPs. (a) CV plots in the N<sub>2</sub>- and O<sub>2</sub>-saturated 0.1 M KOH solution with a scan rate of 50 mV s<sup>-1</sup>, (b) Rotating rate-dependent ORR polarization curves with the scan rate of 10 mVs<sup>-1</sup>, (c) K-L plots of  $j^{-1}$  vs.  $\omega^{-1/2}$  at different potential obtained from ORR results, and (d) The plot of the number of transferred electrons vs. potential. 60
- 3.10 Electrochemical data for ORR over PdFe/C NPs. (a) CV plots in the N<sub>2</sub>- and O<sub>2</sub>-saturated 0.1 M KOH solution with a scan rate of 50 mV s<sup>-1</sup>, (b) Rotating rate-dependent ORR polarization curves with the scan rate of 10 mV s<sup>-1</sup>, (c) K-L plots of  $j^{-1}$  vs.  $\omega^{-1/2}$  at different potential obtained from ORR results, and (d) The plot of the number of transferred electrons vs. potential. 60
- 3.11 Electrochemical data for ORR over Pd<sub>3</sub>Fe/C nanocatalyst. (a) CV plots in the N<sub>2</sub>- and O<sub>2</sub>-saturated 0.1 M KOH solution with a scan rate of 50 mV s<sup>-1</sup>, (b) Rotating rate-dependent ORR polarization curves with the scan rate of 10 mV s<sup>-1</sup>, (c) K-L plots of  $j^{-1}$  vs.  $\omega^{-1/2}$  at different potential obtained from ORR results, and (d) The plot of 61

the number of transferred electrons vs. potential.

- 3.12 (a) CV plots of all the electrocatalysts at N<sub>2</sub>-saturated 0.1 M KOH solution at a scan rate of 50 mV s<sup>-1</sup>, (b) Comparison of rotating rate-dependent ORR polarization curves for different catalyst at 1600 rpm, (c) Tafel plots, and (d) Mass activities of NPs under different potential at 1600 rpm. (These values are calculated as per the loading mass of the NPs). 62
- 3.13 Comparison of Tafel plots Pd<sub>3</sub>Fe/C, PdFe/C and PdFe<sub>3</sub>/C NPs. 63
- 3.14 (a) CA curves of different catalysts, recorded at -0.3 V in a O<sub>2</sub>-saturated 0.1 M KOH solution with a rotation rate of 1600 rpm towards ORR, (b) ORR polarization curves of Pt/C and (c) ORR polarization curves of Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C at 1600 rpm before and after the stability test at different potential scans, (d) Comparison of mass activity after the stability test at different potential scans at 1600 rpm. 64
- 3.15 Electrochemical data for ORR over Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C NPs. 65  
(a) CV plots of Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C, N<sub>2</sub>- and O<sub>2</sub>-saturated 0.5 M H<sub>2</sub>SO<sub>4</sub> solution with a scan rate of 50 mV s<sup>-1</sup>, (b) Rotating rate-dependent ORR polarization curves for respective NPs with the scan rate of 10 mV s<sup>-1</sup>, (c) K-L plots of j<sup>-1</sup> vs. ω<sup>-1/2</sup> for Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C NPs at different potential obtained from (b), and (d) The plot of the number of transferred electrons vs. potential for Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C NPs.
- 3.16 TEM and HR-TEM images of Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C after CA test. distribution curve (inset) for Pd<sub>3</sub>Ni<sub>7</sub> nanoparticles 66
- 3.17 Correlation of binding energy change of Pd 3d to ORR activity in terms of mass activity for Pd<sub>3</sub>Fe<sub>0.5</sub>Cu<sub>0.5</sub>/C 67

		(red), Pd <sub>3</sub> Fe/C (black), and Pd/C (green) NPs.	
4	4A.1	TGA plots of Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C NPs at air atmosphere	79
	4A.2	(a) XRD pattern of Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C and Pd/C NPs. (b) N <sub>2</sub> adsorption-desorption isotherm of Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C. inset of (b) corresponding pore size distribution.	80
	4A.3	TEM (a-d) and HRTEM (e,f) images of Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C.	81
	4A.4	XP spectrum of C 1s (a), Pd 3d (b), Cu 2p (c) and Ni 2p (d) of Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C.	82
	4A.5	(a) CV plots of Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C, N <sub>2</sub> - and O <sub>2</sub> -saturated 0.1 M KOH solution with a scan rate of 50 mV s <sup>-1</sup> , (b) Rotating rate-dependent ORR polarization curves for respective NPs with the scan rate of 10 mV s <sup>-1</sup> , (c) Koutecky–Levich (K-L) plots of $j^{-1}$ vs. $\omega^{-1/2}$ for Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C at different potential obtained from (b), and (d) The plot of the number of transferred electrons vs. potential for Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C.	83
	4A.6	CV plots of various NPs in the N <sub>2</sub> -saturated 0.1 M KOH solution with a scan rate of 50 mV s <sup>-1</sup> .	84
	4A.7	Comparison of rotating rate-dependent ORR polarization curves for different NPs at 1600 rpm	84
	4A.8	Tafel plots of the Pd <sub>3</sub> Cu <sub>0.5</sub> Ni <sub>0.5</sub> /C, Pd <sub>3</sub> Cu/C and Pd <sub>3</sub> Ni/C NPs.	85
	4A.9	(a) Mass activities and (b) Specific activity of electrocatalyst under different potential at 1600 rpm. (These values are calculated as per the loading mass of the electrocatalyst).	86
	4A.10	CA curves of different catalysts, recorded at -0.3 V in a O <sub>2</sub> -saturated 0.1 M KOH solution with a rotation rate of 1600 rpm towards ORR	86

4B.1	EDS spectrum of the Pd <sub>2</sub> CuCo/C NFs.	88
4B.2	TGA profile of Pd <sub>2</sub> CuCo/C NFs in air atmosphere.	89
4B.3	XRD patterns (a), XP spectrum of Pd 3d (b), Cu 2p (c) and Co 2p (d) of Pd <sub>2</sub> CuCo/C NFs.	89
4B.4	Correlation of binding energy change of Pd 3d in Pd <sub>2</sub> CuCo/C NFs and Pd/C.	90
4B.5	TEM (a) and HRTEM (b-d) images of Pd <sub>2</sub> CuCo/C NFs.	91
4B.6	EDS elemental mapping of (a) Carbon (b) Pd (c) Cu (d) Co and (e) overlap image of Pd <sub>2</sub> CuCo/C NFs.	92
4B.7	(a) N <sub>2</sub> adsorption-desorption isotherm of Pd <sub>2</sub> CuCo/C NFs. (b) corresponding pore size distribution.	92
4B.8	CV plots of various NPs in the N <sub>2</sub> -saturated 0.1 M KOH solution with a scan rate of 50 mV s <sup>-1</sup> .	93
4B.9	(a) CV plots of Pd <sub>2</sub> CuCo/C NFs, N <sub>2</sub> - and O <sub>2</sub> -saturated 0.1 M KOH solution with a scan rate of 50 mV s <sup>-1</sup> , (b) Rotating rate-dependent ORR polarization curves for respective NC with the scan rate of 10 mV s <sup>-1</sup> , (c) Koutecky–Levich (K-L) plots of $j^{-1}$ vs. $\omega^{-1/2}$ for Pd <sub>2</sub> CuCo/C NFs at different potential obtained from (b), and (d) The plot of the number of transferred electrons vs. potential for Pd <sub>2</sub> CuCo/C NFs.	94
4B.10	(a) Comparison of rotating rate-dependent ORR polarization curves for different catalyst at 1600 rpm, (b) Tafel plots.	94
4B.11	(a) Mass activities and (b) Specific activity of electrocatalyst under different potential at 1600 rpm. (These values are calculated as per the loading mass of the Pd or Pt).	95
4B.12	(a) Comparison of rotating rate-dependent ORR polarization curves for different catalyst at 1600 rpm at	96

	0.5 M H <sub>2</sub> SO <sub>4</sub> (b) Mass activities and (c) Specific activity of electrocatalyst under different potential at 1600 rpm. (These values are calculated as per the loading mass of the Pd or Pt).	
	4B.13 CA curves of different catalysts, recorded at -0.3 V in a O <sub>2</sub> -saturated 0.1 M KOH solution with a rotation rate of 1600 rpm.	96
	4B.14 (a) ORR polarization curves of Pd <sub>2</sub> CuCo/C NFs and (b) ORR polarization curves of Pt/C at 1600 rpm before and after 1000 CV cycles.	97
5	5A.1 XRD patterns (a) Cu-CuFe <sub>2</sub> O <sub>4</sub> /C (black line) and bare Cu-CuFe <sub>2</sub> O <sub>4</sub> . (red line) (b) TGA Curve (c) N <sub>2</sub> adsorption-desorption isotherm of Cu-CuFe <sub>2</sub> O <sub>4</sub> /C (d) Pore size distribution curve of Cu-CuFe <sub>2</sub> O <sub>4</sub> /C	107
	5A.2 (a) XPS survey spectrum of Cu-CuFe <sub>2</sub> O <sub>4</sub> /C. High-resolution XP spectra of (b) Cu 2p, (c) Fe 2p, (d) C 1 s for Cu-CuFe <sub>2</sub> O <sub>4</sub> /C.	108
	5A.3 High-resolution XPS spectra of (a) Cu 2p, (b) Fe 2p, for Cu-CuFe <sub>2</sub> O <sub>4</sub> .	109
	5A.4 (a) TEM images of Cu-CuFe <sub>2</sub> O <sub>4</sub> /C, inset of (a) shows the SAED pattern, (b) HRTEM image of Cu-CuFe <sub>2</sub> O <sub>4</sub> /C showing black contrast of metal/metal oxide and carbon, (c,d) HRTEM image of Cu-CuFe <sub>2</sub> O <sub>4</sub> /C with lattice fringes corresponding to various constituents.	109
	5A.5 (a) TEM images of Cu-CuFe <sub>2</sub> O <sub>4</sub> , (b) High magnitude TEM image of Cu-CuFe <sub>2</sub> O <sub>4</sub>	110
	5A.6 (a) CVs of the Cu-CuFe <sub>2</sub> O <sub>4</sub> /C, Cu-CuFe <sub>2</sub> O <sub>4</sub> + C mixture, Cu- CuFe <sub>2</sub> O <sub>4</sub> , and carbon on GCEs in an O <sub>2</sub> -saturated (solid line) or N <sub>2</sub> -saturated (dashed line) 0.1 M KOH solutions and (b) LSVs of the Cu-CuFe <sub>2</sub> O <sub>4</sub> /C, Cu-CuFe <sub>2</sub> O <sub>4</sub> + C, Cu-CuFe <sub>2</sub> O <sub>4</sub> , carbon, and Pt/C in an O <sub>2</sub> -	112



saturated 0.1 M KOH solution at 1600 rpm.

- 5A.7 (a) LSVs of the Cu-CuFe<sub>2</sub>O<sub>4</sub>/C NPs in O<sub>2</sub>-saturated 0.1 M KOH solution at various rotation rates at a scan rate of 10 mV/s, (b) K–L plots in the potential range of –0.4 to –0.7 V, (c) the plot of the number of transferred electrons vs. potential for Cu-CuFe<sub>2</sub>O<sub>4</sub>/C NPs., (d) Tafel plots of Pt/C and Cu-CuFe<sub>2</sub>O<sub>4</sub>/C NPs. 113
- 5A.8 (a) LSVs of the Cu-CuFe<sub>2</sub>O<sub>4</sub>+C in O<sub>2</sub>-saturated 0.1 M KOH solution at various rotation rates at a scan rate of 10 mV/s, (b) K–L plots in the potential range of –0.4 to –0.7 V, (c) the plot of number of transferred electrons vs. potential for Cu-CuFe<sub>2</sub>O<sub>4</sub>+C. 113
- 5A.9 (a) LSVs of the bare Cu-CuFe<sub>2</sub>O<sub>4</sub> NPs in O<sub>2</sub>-saturated 0.1 M KOH solution at various rotation rates at a scan rate of 10 mV/s, (b) K–L plots in the potential range of –0.4 to –0.7 V, (c) the plot of number of transferred electrons vs. potential for bare Cu-CuFe<sub>2</sub>O<sub>4</sub> NPs. 114
- 5A.10 (a) LSVs of the CuFe<sub>2</sub>O<sub>4</sub>/C NPs in O<sub>2</sub>-saturated 0.1 M KOH solution at various rotation rates at a scan rate of 10 mV/s, (b) K–L plots in the potential range of –0.4 to –0.7 V, (c) the plot of number of transferred electrons vs. potential for CuFe<sub>2</sub>O<sub>4</sub>/C NPs. 115
- 5A.11 CV curves of a) Cu-CuFe<sub>2</sub>O<sub>4</sub>/C and c) CuFe<sub>2</sub>O<sub>4</sub>/C NPs with different scan rates, charging current density differences as function of scan rates for b) Cu-CuFe<sub>2</sub>O<sub>4</sub>/C and d) CuFe<sub>2</sub>O<sub>4</sub>/C. 116
- 5A.12 ORR polarization curves of (a) Cu-CuFe<sub>2</sub>O<sub>4</sub>/C and (b) 20 wt% Pt/C at 1600 rpm in 0.1 M KOH solution before and after the stability test of different potential scans. 117
- 5A.13 CA curves of different NPs recorded at –0.3 V in O<sub>2</sub>- 117

	saturated 0.1 M KOH solution with a rotation rate of 1600 rpm.	
5A.14	CV curves of different catalysts recorded in N <sub>2</sub> -saturated 0.1 M KOH solution.	118
5B.1	XRD patterns of CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C and CuO-Co <sub>3</sub> O <sub>4</sub> /C NPs.	120
5B.2	TGA curve of CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C in air atmosphere	121
5B.3	N <sub>2</sub> adsorption/desorption isotherm and the pore size distribution curve (inset) of CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C NPs.	121
5B.4	(a),(b) TEM images of CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C NPs at different resolution.	122
5B.5	XPS analysis for CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C NPs (a) the survey spectrum; and the high resolution core-level XP-spectrum of (b) C 1s, (c) Cu 2p and (d) Co 2p region.	123
5B.6	CV plots of (a) Co/C, (b) Cu/C, (c) CuO-Co <sub>3</sub> O <sub>4</sub> /C, (d) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> + C, (e) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C and (f) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> NPs in both N <sub>2</sub> and O <sub>2</sub> -saturated 0.1 M KOH solution.	124
5B.7	Rotating rate dependent LSVs at the scan rate of 10 mVs <sup>-1</sup> in O <sub>2</sub> -saturated 0.1 M KOH over (a) Co/C, (b) Cu/C, (c) CuO-Co <sub>3</sub> O <sub>4</sub> /C, (d) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> + C, (e) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C and (f) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> , respectively.	125
5B.8	Koutecky-Levich plots at different potentials for (a) Co/C, (b) Cu/C, (c) CuO-Co <sub>3</sub> O <sub>4</sub> /C, (d) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> + C, (e) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C and (f) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> , respectively.	125
5B.9	LSVs of the all electrocatalysts in O <sub>2</sub> -saturated 0.1 M KOH solution at 1600 rpm.	127
5B.10	The Tafel plots of Pt/C and CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C NPs.	127
5B.11	CV curves of a) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C and c) CuCo/CuO-	128

	Co <sub>3</sub> O <sub>4</sub> NPs with different scan rates. (D) Charging current density differences as function of scan rates for b) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> /C and d) CuCo/CuO-Co <sub>3</sub> O <sub>4</sub> .	
	5B.12 CA curves of different catalysts, recorded at -0.3 V in O <sub>2</sub> -saturated 0.1 M KOH solution with a rotation rate of 1600 rpm.	128
6	6A.1 (a) XRD patterns of the Fe <sub>2</sub> O <sub>3</sub> , Cu and Cu/Fe <sub>2</sub> O <sub>3</sub> , (b) TGA analysis of Cu/Fe <sub>2</sub> O <sub>3</sub> in air atmosphere, and (c) N <sub>2</sub> adsorption-desorption isotherm of Cu/Fe <sub>2</sub> O <sub>3</sub> .	137
	6A.2 (a-d) TEM images of Cu/Fe <sub>2</sub> O <sub>3</sub> , inset of (c) shows the SAED pattern and (e-g) HRTEM images of Cu/Fe <sub>2</sub> O <sub>3</sub> showing the lattice fringes of Cu and Fe <sub>2</sub> O <sub>3</sub> .	137
	6A.3 (a) XPS survey spectrum of Cu/Fe <sub>2</sub> O <sub>3</sub> . High-resolution XPS spectra of (b) Cu 2p, (c) Fe 2p, and (d) O 1 s for Cu/Fe <sub>2</sub> O <sub>3</sub> .	138
	6A.4 Time-dependent absorption spectra of the reaction solution of 4-NP reduction to 4-AP over (a) Cu, (b) Fe <sub>2</sub> O <sub>3</sub> , (c) Cu/Fe <sub>2</sub> O <sub>3</sub> and (d) plot of ln(A <sub>t</sub> /A <sub>0</sub> ) against the reaction time of the catalytic reduction of 4-NP to 4-AP over Cu/ Fe <sub>2</sub> O <sub>3</sub> .	140
	6A.5 Conversion against no. of cycles, (b) plot of ln(A <sub>t</sub> /A <sub>0</sub> ) against the reaction time of the catalytic reduction of 4-NP to 4-AP for different cycles.	146
	6A.6 The magnetic hysteresis loops of the Cu/Fe <sub>2</sub> O <sub>3</sub> NPs. Inset shows the separation of Cu/Fe <sub>2</sub> O <sub>3</sub> NPs from aqueous solution accomplish by an external magnet.	146
	6B.1 XRD pattern of CuNi NPs along with diffraction lines from JCPDS database corresponding to Cu and Ni.	148
	6B.2 TEM (a,b) and HR-TEM (c,d) of CuNi NPs. Inset in the figure (d) shows the SAED pattern of the NPs	149
	6B.3 (a) XPS survey spectra of CuNi, core level XP spectra of	150

	(b) Cu 2p and (c) Ni 2p and (d) O1s.	
6B.4	(a) XRD pattern of CuNi NPs after three catalytic cycles.	153
6B.5	Room-temperature magnetization curve of the CuNi NPs.	154

## LIST OF SCHEMES

Chapter	Scheme	Title	Page No.
1	1.1	Schematic representation of 'Top-down' approach	9
	1.2	Schematic representation of 'bottom-up' approach	9
	1.3	Schematic representation of different catalysts.	11
6	6A.1	Reaction mechanism proposed by Haber for the reduction of nitroaromatics compounds	145
	6B.1	Transfer hydrogenation of nitroaromatics to corresponding anilines over bimetallic CuNi NPs	151