

Table of Contents

Abstract	i-vi
Declaration	vii
Certificate of the Supervisor	vii
Acknowledgements	ix-x
Table of Contents	xi-xviii
List of Tables	xix-xxi
List of Figures	xxii-xxiv
List of Abbreviations	xxv-xxvi
List of Symbols and Units	xxvii-xxviii

Chapter 1	1-7
Introduction	
1.1 Introduction	1
1.2 Objectives	4
Bibliography	4
Chapter 2	9-39
Review of literature	
2.1 Introduction	9
2.1.1 Tamarillo tree overview	11
2.1.2 Classification of tamarillo varieties	11
2.2 Physical composition of tamarillo fruit	13
2.3 Nutritional composition for tamarillo	14
2.4 Bioactive components of tamarillo	15
2.4.1 Polyphenols of tamarillo	16
2.4.2 Phenolic acids	17
2.4.3 Flavonoids	18
2.4.4 Anthocyanin	20
2.4.5 Carotenoids	21
2.3. Health benefits of tamarillo	22
2.3.1 Reduced risk of obesity and diabetes	23
2.3.2 Inhibition of P12 cells and LDL oxidation	24
2.3.3 Prebiotic activity	24
2.3.4 Antioxidant Effect	25
2.4 Effect of processing on bioactive compounds present in tamarillo	26
2.4.1 Polyphenol-rich extract	27
2.4.2 Carotenoids enriched puree	27
2.4.3 Pulp powder	28
2.4.4 Carotenoids encapsulated seed oil nanoemulsion	29
Bibliography	29

Chapter 3 41-70
Phytochemical profiling of different varieties of tamarillos and study the effect of ultrasound and high pressure homogenization on bioactive compounds

3.1 Introduction	41
3.2 Materials and methods	42
3.2.1 Raw sample collection	42
3.2.2 Sample preparation for proximate & biochemical analysis	42
3.2.3 Mineral content	42
3.2.4 Colour	43
3.2.5. Sample treatment	43
3.2.5.1 Conventional method	43
3.2.5.2 High pressure homogenization	43
3.2.5.3 Ultrasound-assisted treatment	43
3.2.6 Total monomeric anthocyanin content	44
3.2.7 Total carotenoids content	44
3.2.8 Sample extraction for phenolics and antioxidant activity	44
3.2.8.1 Total phenolic content	45
3.2.8.2 Total flavonoid content	45
3.2.8.3 DPPH radical scavenging activity	45
3.2.8.4 ABTS radical scavenging activity	46
3.2.9 HPLC of phenolic acids	46
3.2.10 HPLC of anthocyanins	46
3.2.11 HPLC of carotenoids	47
3.2.12 <i>In-vitro</i> digestibility study	47
3.2.13 Determination of bioaccessibility	48
3.2.14 Statistical analysis	48
3.3 Results and discussion	48
3.3.1 Proximate and biochemical analysis	48
3.3.2 Effect of HPH and US on bioactive compounds in tamarillos	50
3.3.2.1 Total monomeric anthocyanin content	50
3.3.2.2 Total carotenoids content	51
3.3.2.3 Total phenolic content	52
3.3.2.4 Total flavonoids content	53
3.3.2.5 DPPH radical scavenging activity	54
3.3.2.6 ABTS radical scavenging activity	54
3.3.3 HPLC analysis of the phenolic acids in control, HPH, and US-treated tamarillo samples	56
3.3.4 HPLC analysis of the anthocyanins in control, HPH, and US-treated tamarillo samples	56
3.3.5 HPLC analysis of the carotenoids in control, HPH, and US-treated tamarillo samples	57
3.3.6 <i>In-vitro</i> digestion of control, HPH, and US treated tamarillo samples	58
3.3 Conclusion	67
Bibliography	67

Chapter 4 Section I	72-91
Optimization of bioactive compounds from yellow and red tamarillo using ultrasound assisted extraction	
4.1.1. Introduction	72
4.1.2. Material and methods	73
4.1.2.1. Materials	73
4.1.2.2. Ultrasonic assisted extraction	73
4.1.2.3. Selection of variables	74
4.1.2.4. Experimental Design	74
4.1.2.5. Total phenolic content (TPC)	75
4.1.2.6. Total flavonoid content (TFC)	75
4.1.2.7. DPPH radical scavenging activity	75
4.1.2.7. Statistical analysis	76
4.1.3. Results and discussion	76
4.1.3.1. Selection of independent variables	76
4.1.3.2. Fitting the model in the experimental design	78
4.1.3.3. Effect of independent variables on the extraction of total phenolic content	78
4.1.3.4 Effect of independent variables on the extraction of total flavonoids content	79
4.1.3.5. Effect of independent variables on DPPH radical scavenging activity of the extract	85
4.1.3.6 Optimization	87
4.1.4. Conclusions	87
Bibliography	88
Chapter 4 Section II	92-117
Optimization of supercritical fluid extraction and ultrasound assisted extraction of extraction conditions of purple tamarillo	
4.2.1. Introduction	92
4.2.2. Materials and methods	93
4.2.2.1. Plant Materials & sample preparation	93
4.2.2.2. Supercritical fluid extraction (SCFE)	94
4.2.2.3. Ultrasound-assisted extraction (UAE)	95
4.2.2.4. Extraction of bioactive compounds	95
4.2.2.5. Total phenolic content (TPC)	95
4.2.2.6. Total monomeric anthocyanin content	96
4.2.2.7. HPLC analysis of phenolic acids	96
4.2.2.8 HPLC analysis of anthocyanins	96
4.2.2.9. Stability of the anthocyanin	97
4.2.2.10. Scanning electron microscopy	97
4.2.2.11. Statistical analysis	97
4.2.3. Results and discussion	98
4.2.3.1. Determination of independent variables limits of SCFE	98
4.2.3.1.1. Effect of independent variables on responses using SCFE	98
4.2.3.1.2. Verification of model for SCFE	102
4.2.3.2. Determination of the limits of independent variables of	103

UAE		
4.2.3.2.1. Effect of independent variables on response using UAE technique	104	
4.2.3.2.2. Verification of model for UAE	105	
4.2.3.3. Optimization and comparison in techniques	108	
4.2.3.4. HPLC profile of phenolic acids	110	
4.2.3.5. HPLC profile of anthocyanins	110	
4.2.3.6. Anthocyanin stability	112	
4.2.3.7. Scanning electron microscopy	113	
4.2.4. Conclusion	113	
Bibliography	114	
Chapter 5 Section I		119-133
Study the effect of addition of olive oil and thermal treatment on tamarillo puree		
5.1.1. Introduction	119	
5.1.2. Materials and methods	120	
5.1.2.1. Materials	120	
5.1.2.2. Sample preparation	120	
5.1.2.3. Thermal processing of puree	120	
5.1.2.4. Biochemical analysis	121	
5.1.2.5. Total phenolic content	121	
5.1.2.6. Total flavonoid content	121	
5.1.2.7. DPPH radical scavenging activity	121	
5.1.2.8. Total carotenoids content	122	
5.1.2.9. Colour	122	
5.1.2.10. Statistical analysis	122	
5.1.3. Results and discussion	123	
5.1.3.1. Effect of thermal processing and oil on pH and TSS	123	
5.1.3.2. Effect of thermal processing and oil on total phenol content (TPC)	123	
5.1.3.3. Effect of heating and addition of oil on total flavonoids content (TFC)	125	
5.1.3.4. Effect of heating and addition of oil on <i>in-vitro</i> antioxidant activity	125	
5.1.3.5 Effect of addition of heating and addition of oil on total carotenoids content	125	
5.1.3.6. Effect of heating time and addition of oil on colour values	126	
5.1.4. Conclusion	130	
Bibliography	130	
Chapter 5 Section II		134-165
Study the effect of high pressure homogenization and thermal processing of tamarillo puree		
5.2.1 Introduction	134	
5.2.2. Materials and methods	135	
5.2.2.1. Preprocessing	135	
5.2.2.2. High pressure homogenization	135	

5.2.2.3. Light Microscopy	136
5.2.2.4. Colour measurement	136
5.2.2.5. Particle size analysis	136
5.2.2.6. Determination of bioactive compounds	136
5.2.2.6.1 Total phenolic content	136
5.2.2.6.2. DPPH radical scavenging activity	137
5.2.2.6.3. Determination of total carotenoids concentration (TCC)	137
5.2.2.7. HPLC of phenolic acids	137
5.2.2.8. HPLC of carotenoids	137
5.2.2.9. Thermal inactivation studies	138
5.2.2.10. <i>In-vitro</i> digestion of puree	138
5.2.2.11. Bioaccessibility of bioactive compounds	139
5.2.2.12. Storage study of puree	139
5.2.2.13. Statistical analysis	140
5.2.3. Results and discussion	140
5.2.3.1. Particle size	140
5.2.3.2. Microstructure	140
5.2.3.3. Colour analysis	141
5.2.3.4. Total phenolic content	142
5.2.3.5. DPPH radical scavenging activity	143
5.2.3.6. Total carotenoids content (TCC)	143
5.2.3.7. HPLC of phenolic acids	143
5.2.3.8. HPLC of carotenoids	144
5.2.3.9. D value of puree	151
5.2.3.10. Bioaccessibility of TPC and phenolic acids of tamarillo puree	156
5.2.3.11 Bioaccessibility of TCC and carotenoids present in tamarillo puree	156
5.2.3.12 Effect on pH, TSS, vitamin C, TPC, DPPH scavenging activity, TCC, and total microbial load during storage of HPH processed heat treated bottled tamarillo puree	157
5.2.4. Conclusion	160
Bibliography	160

Chapter 6	167-200
Development and Characterization of Foam Mat Tamarillo Powder	
6.1. Introduction	167
6.2. Materials and Methods	168
6.2.1. Materials	168
6.2.2. Sample and foam preparation	168
6.2.3. Selection of drying temperature and drying kinetics	169
6.2.3.1 Moisture diffusivity	170
6.2.4. Powder analysis	171
6.2.4.1. Bulk density, tapped density, flowability, and cohesiveness	171
6.2.4.2. Wettability and Solubility	172
6.2.4.3. Hygroscopicity	172
6.2.4.4. Colour parameters	172

6.2.5. Scanning electron microscopy	173
6.2.6. Determination of phytochemicals in foam mat dried powder	173
6.2.6.1. Total phenolic content	173
6.2.6.2. Total flavonoid content	173
6.2.6.3. Total carotenoids content	173
6.2.7. DPPH scavenging activity and ABTS radical scavenging activity of foam mat dried powder	174
6.2.7.1. DPPH radical scavenging activity	174
6.2.7.2 ABTS radical scavenging activity	174
6.2.8. HPLC analysis of phenolic acids of foam mat dried powder	175
6.2.9. HPLC of carotenoids of foam mat dried powder	175
6.2.10. Statistical analysis	175
6.3. Results and Discussion	176
6.3.1. Selection of drying temperature	176
6.3.2. Foam properties	176
6.3.2.1 Foam expansion	176
6.3.2.2. Foam density	177
6.3.2.3. Foam stability	178
6.3.3. Drying kinetics and moisture diffusivity	179
6.3.4. Powder characteristics	179
6.3.4.1. Bulk and tapped density of powder	179
6.3.4.2. Flowability and cohesiveness	181
6.3.4.3. Wettability	181
6.3.4.4. Solubility	182
6.3.4.5. Hygroscopicity	182
6.3.4.6. Colour parameters	182
6.3.5. Scanning electron microscopy (SEM)	183
6.3.6. Effect of foaming agents on TPC, TPC, total carotenoids, and <i>in-vitro</i> antioxidant properties of powder samples	186
6.3.7. HPLC analysis of phenolic acids present in foam mat dried powder	189
6.3.8 HPLC of carotenoids present in foam mat dried powder	191
6.4. Conclusion	194
Bibliography	194

Chapter 7 Section I 202-225
Extraction of carotenoids from tamarillo peel using high shear disperser and ultrasound using green extraction solvent

7.1.1. Introduction	202
7.1.2. Materials and methods	204
7.1.2.1. Plant materials and sample preparation	204
7.1.2.2. High shear disperser (HSD)	204
7.1.2.3. Ultrasound assisted extraction (UAE)	204
7.1.2.4. Experimental design	204
7.1.2.5. Extraction of the TCC	205
7.1.2.6. Determination of oil quality parameters	205
7.1.2.6.1. Acid value and peroxide value	205

7.1.2.6.2. Total phenolic content	206
7.1.2.6.3. DPPH radical scavenging activity	206
7.1.2.6.4. Colour values	206
7.1.2.7. HPLC of phenolic acids of optimized extracts	206
7.1.2.8. HPLC of TCC of optimized extract	207
7.1.2.9. Preparation of mayonnaise	207
7.1.2.10. Sensory analysis	207
7.1.2.11. In-vitro digestion of mayonnaise	208
7.1.2.12. Calculation of bioaccessibility	208
7.1.2.13. Statistical analysis	208
7.1.3. Results and discussion	209
7.1.3.1. Effect of the independent variables on TCC using high shear disperser	209
7.1.3.2. Effect of the independent variables on TCC using ultrasound assisted extraction	212
7.1.3.3. Optimization of response surface methodology and comparison of the two extraction methods	213
7.1.3.4. Effect of HPH and UAE on acid value, peroxide value, total phenolic content, in vitro antioxidant activity, and colour properties of oil	215
7.1.3.5. HPLC of phenolic acids	217
7.1.3.6. HPLC of carotenoids	218
7.1.3.7. Sensory characteristics of mayonnaise	219
7.1.3.8. <i>In-vitro</i> digestion of mayonnaise	220
7.1.4. Conclusion	221
Bibliography	222
Chapter 7 Section II	226-248
Development of carotenoids loaded nanoemulsion using ultrasonication	
7.2.1. Introduction	226
7.2.2. Materials and methods	227
7.2.2.1. Plant materials	227
7.2.2.2. Extraction of tamarillo seed oil	227
7.2.2.3. GC-MS analysis of the tamarillo seed oil	227
7.2.2.4. Extraction of carotenoids	228
7.2.2.5. Preparation of carotenoids-loaded NE	228
7.2.2.5.1. Experimental design	228
7.2.2.5.2. Preparation of NE	228
7.2.2.6. Characterization of NE	229
7.2.2.6.1. Determination of PS and PDI	229
7.2.2.6.2. Encapsulation efficiency	229
7.2.2.7. Stability of carotenoids-loaded NE against pH and ionic strength	229
7.2.2.8. Physical stability of carotenoids-loaded NE at stimulated gastrointestinal conditions	230
7.2.2.9. Effect of storage temperature on stability of the carotenoids-loaded NE	230
7.2.2.10. Transmission electron microscopy	230
7.2.2.11. Statistical analysis	230

7.2.3. Results and discussion	231
7.2.3.1. GC-MS analysis of the tamarillo seed oil	231
7.2.3.2. Fitting the model in experimental design	232
7.2.3.3. Effect of independent variables on PS	233
7.2.3.4. Effect of independent variables on PDI	233
7.2.3.5. Effect of independent variables on EE	234
7.2.3.6. Optimization of process conditions	235
7.2.3.7. Zeta Potential of optimized NE	235
7.2.3.8. Effect of pH on PS and carotenoids stability	238
7.2.3.9. Effect of ionic strength on PS and carotenoids stability	238
7.2.3.10. Effect of temperature on PS and carotenoids release during storage	239
7.2.3.11. In-vitro digestion of NE	241
7.2.3.12. Transmission electron microscopy	243
7.2.4. Conclusion	244
Bibliography	244
 Chapter 8	250-255
Summary, conclusions and future scope	
8.1 Salient findings	250
8.1.1 Analysis of the biochemical properties, nutrient composition, and antioxidant activities of the red, yellow, and purple varieties of tamarillos	250
8.1.2 Optimization of the conditions of ultrasonication and supercritical fluid extraction of polyphenols from tamarillo pulp and identification of polyphenols in the optimized extracts	251
8.1.3 Standardization of the bottling of tamarillo puree and study of the in vitro bioaccessibility of carotenoids	252
8.1.4 Development of tamarillo powder using connective drying and study of the drying behaviour and powder properties	253
8.1.5 Study of the physical and chemical properties of tamarillo seed oil nanoemulsions incorporated with extracted carotenoids of the fruit	253
8.2 Future scope	255
 List of Publications	257