

Contents

S. No.	Title	Page No.
I.	Abstract	i-v
II.	Declaration	vi
III.	Certificate of the supervisor	vii
IV.	Acknowledgement	viii-ix
V.	Contents	x-xvi
VI.	List of Tables	xvii-xviii
VII.	List of Figures	xix-xxi
VIII.	Abbreviations	xxii-xxvi

CONTENTS

1.	Chapter I	1-8
	Introduction	
2.	Chapter II	9-57
	Review of Literature	
2.1.	Historical background and Origin	9
2.2.	Bioactive compound present in Haritaki	11-14
2.3.	Drying kinetics of some fruits and vegetables	14
2.4.	Moisture sorption isotherm of some fruits and vegetables	15
2.5.	Degradation of total phenolic content and antioxidant activity of some fruits and vegetables	16-17
2.6.	Pre-treatment for extraction	17-23
2.6.1.	Extraction of bioactive compound	18
2.6.1.1.	Conventional extraction techniques	19
2.6.1.2.	Non- conventional extraction techniques	19-23
2.6.1.2.1.	Ultrasound-assisted extraction (UAE)	20
2.6.1.2.2.	Supercritical fluid extraction (SFE)	21
2.6.1.3.	Integration of different extraction techniques	21-22
2.6.1.3.1	Ultrasound–Microwave-Assisted Extraction (UMAE)	21-22
2.6.1.3.2.	Enzyme-Based–Ultrasound–Microwave-Assisted Extraction (EUMAE)	22-23
2.7.	Therapeutic properties and pharmacological attributes	23-28

2.8.	Traditional use of haritaki	28-29
2.9.	Application of haritaki bioactive compounds in pharmaceutical and food industry	29-30
2.10.	Encapsulation	30-31
2.11.	Yoghurt	31-36
2.11.1.	Hydrocolloids in yoghurt	32-34
2.11.2.	Encapsulated phenolic compounds in yoghurt	34-36
2.12.	Conclusion	36
2.13.	References	37-57
3.	Chapter III	58-92
	To study the drying behaviour of fresh haritaki (<i>Terminalia chebula</i>) fruit	
3.1.	Introduction	58-60
3.2.	Materials and Methods	60
3.2.1.	Materials	60
3.2.2.	Engineering properties of the fruit	60-61
3.2.3.	Drying of sample	61
3.2.4.	Moisture content	61
3.2.5.	Mathematical modelling	61-62
3.2.6.	Moisture diffusivity	62-6.
3.2.7.	Activation energy	63
3.2.8.	Phytochemical determination	63-64
3.2.8.1.	Extract preparation	63
3.2.8.2.	Total phenolic content (TPC)	63-64
3.2.8.3.	Total flavonoid content (TFC)	64
3.2.8.4.	Total antioxidant activity (DPPH radical scavenging activity)	64
3.2.8.5.	Vitamin C	64
3.2.9.	Color measurement	64
3.2.10.	Degradation kinetics	64-65
3.2.11.	Determination of sorption isotherm	65
3.2.12.	Modelling of sorption isotherms	65-67
3.2.13.	Statistical analysis	67
3.3.	Results and discussion	67-84

3.3.1.	Engineering properties of the haritaki fruit	67-69
3.3.2.	Fitting of drying curves	69-70
3.3.3.	Drying model evaluation	70-73
3.3.4.	Effective moisture diffusivity	73-74
3.3.5.	Activation energy	74-75
3.3.6.	Degradation kinetics of TPC, TFC, vitamin C, DPPH during drying	75-77
3.3.7.	Kinetics of change in color during drying	77-78
3.3.8.	Moisture Sorption Characteristics of haritaki pulp powder	78-81
3.3.9.	Mathematical Modelling and Fitting of Moisture Sorption Data	81-84
3.4.	Conclusion	85
3.5.	References	86-92
4.	Chapter IV	93-117
To Comparison of response surface methodology (RSM) and artificial neural network (ANN) modelling for supercritical fluid extraction of phytochemicals from Terminalia chebula pulp and optimization using RSM coupled with desirability function (DF) and genetic algorithm (GA) and ANN with GA		
4.1.	Introduction	93-95
4.2.	Materials and Methods	95
4.2.1.	Preparation of dried haritaki pulp powder	95
4.2.2.	Extraction of phytochemicals by supercritical CO ₂	95
4.2.3.	Experimental design	95-96
4.2.4.	Determination of phytochemical content and antioxidant activity of the extracts	96
4.2.5.	RSM modelling	96-97
4.2.6.	ANN modelling	97
4.2.7.	Analysis of the developed models	97-98
4.2.8.	Optimization of the process	98
4.3.	Results and discussion	98-113
4.3.1.	RSM modelling and Effect of process parameters and their interaction on the responses	98-107
4.3.2.	ANN modelling	107-109

4.3.3.	Comparison of the developed RSM and ANN models	110-112
4.3.4.	Optimization of the process parameters by different approaches	112-113
4.4.	Conclusion	113
4.5.	References	114-117
5.	Chapter V	118-149
	To extract the bioactive compounds from fresh and dried fruit using various green and novel technologies and their combination	
5.1.	Introduction	118-119
5.2.	Materials and Methods	119-125
5.2.1.	Materials	119
5.2.2.	Drying of haritaki pulp using different approaches	119-120
5.2.3.	Pre-treatment optimization of extraction parameters using SFE	120-123
5.2.3.1.	Novel techniques for Pre-treatment	122-123
5.2.3.1.1.	Ultrasound-assisted extraction (UAE)	122
5.2.3.1.2.	Microwave assisted extraction (MAE)	122
5.2.3.1.3.	Enzyme-assisted extraction (EAE)	122
5.2.3.2.	Combination of techniques for Pre-treatment	122-23
5.2.3.2.1.	Microwave assisted -Enzymatic extraction	122
5.2.3.2.2.	Ultrasound assisted -Microwave extraction	123
5.2.3.2.3.	Ultrasound assisted - Enzyme extraction	123
5.2.3.2.4.	Ultrasound- Microwave- Enzyme assisted extraction	123
5.2.4.	Extraction and characterization of bioactive compounds using SFE	123
5.2.5.	Determination of bioactive compounds	123
5.2.5.	Fourier transform infrared spectroscopy (FT-IR)	123
5.2.6.	Stability of pre-treated fresh haritaki extract	124
5.2.6.1.	Effect of pH on the stability of bioactive compounds	124
5.2.6.2.	Effect of heat treatment on the stability of bioactive compounds	124
5.2.7	Liquid Chromatography Mass Spectrometry (LC-MS)	124
5.2.8.	Color properties	124
5.2.9.	Statistical analysis	125
5.3.	Results and Discussion	125-141
5.3.1.	Impact of drying techniques on the extraction of bioactive compounds	125-126

5.3.2.	Pre-treatment of haritaki pulp	127
5.3.2.1.	Effect of novel techniques on extraction of bioactive compound of haritaki	127-130
5.3.2.1.1.	Ultrasound-assisted extraction	127-128
5.3.2.1.2.	Microwave-assisted extraction	128-129
5.3.2.1.3.	Enzyme-assisted extraction	129-130
5.3.2.2.	Effect of combined techniques on extraction of bioactive compound of haritaki	130-133
5.3.2.2.1.	Microwave assisted -Enzymatic extraction	130-131
5.3.2.2.2.	Ultrasound assisted - Microwave extraction	131-132
5.3.2.2.3.	Ultrasound assisted - Enzyme extraction	132-133
5.3.2.2.4.	Ultrasound-Microwave-Enzyme- assisted extraction	133
5.3.3.	Spectral analysis of pre-treated haritaki pulp	133-134
5.3.4.	Extraction and characterization of bioactive compounds using SFE	134-135
5.3.4.1.	Effect of pH and temperature on the stability of bioactive compounds	135-137
5.3.4.2.	LC-MS of microwave assisted ultrasound pre-treated haritaki pulp SFE extract	137-139
5.3.5.	Color properties	139-141
5.4.	Conclusion	142
5.5.	References	143-149
6.	Chapter VI	150-172
	To encapsulate the bioactive compounds extracted from haritaki using a combination of biopolymeric materials	
6.1.	Introduction	150-151
6.2.	Materials and Methods	151-155
6.2.1.	Material	151
6.2.2.	Extraction of bioactive compounds from haritaki	151-152
6.2.3.	Encapsulation of bioactive compounds	152
6.2.3.1.	Conventional technique	152
6.2.3.2.	Advanced technique	152
6.2.4.	Characterization of physical properties of encapsulates	152-154
6.2.4.1.	Moisture content and water activity	152

6.2.4.2.	Process yield (PY)	152
6.2.4.3.	Encapsulation efficiency	153
6.2.5.	Packing and flow properties	153-154
6.2.6.	Color properties	154
6.2.7.	Fourier transform infrared spectroscopy (FT-IR) spectra	154
6.2.8.	Surface morphology	154
6.2.9.	Thermodynamic properties	154-155
6.2.9.1.	Differential scanning calorimetry (DSC)	154
6.2.9.2.	Thermo-gravimetric analysis (TGA)	154-155
6.2.10.	Statistical analysis	155
6.3.	Result and Discussion	155-165
6.3.1.	Encapsulation of bioactive compounds using conventional and advanced method	155-156
6.3.2.	Characterization of physical properties of encapsulates	156-157
6.3.2.1.	Process yield & Encapsulation efficiency	156-157
6.3.2.2.	Moisture content and water activity	157
6.3.2.3.	Bulk, and Tapped density	157
6.3.2.4.	Carr's index and Hausner ratio	157-159
6.3.3.	Color properties	159
6.3.4.	Surface morphology of powder	159-160
6.3.5.	FTIR Spectra of powder	160-161
6.3.6.	XRD	162
6.3.7.	Thermodynamic properties	162-165
6.3.7.1.	DSC	162-164
6.3.7.2.	TGA	164-165
6.4.	Conclusion	166
6.5.	References	167-172
7.	Chapter VII	173-202
	To develop a functional yoghurt by incorporating extract and microencapsulated extract containing bioactive compounds from haritaki	
7.1.	Introduction	173
7.2.	Materials and Methods	174-177

7.2.1.	Raw material	174
7.2.2.	Yogurt preparation	174-175
7.2.3.	pH and titrable acidity of yoghurt during storage	175
7.2.4.	Colour analysis of yoghurt samples during storage	176
7.2.5.	Syneresis of yoghurt during storage	176
7.2.6.	Total phenol content (TPC)	176
7.2.7.	DPPH free radical scavenging activity	176
7.2.8.	Textural properties of yoghurt during storage	176
7.2.9.	Rheological analysis of yoghurt during storage	176
7.2.10.	Sensory analysis of yoghurt samples	177
7.2.11.	Statistical analysis	177
7.3.	Results and discussions	177-195
7.3.1.	Physico-chemical properties	177-180
7.3.2.	Changes in color of yoghurt	180-182
7.3.3.	Phytochemical properties of yoghurt	183-185
7.3.4.	Rheological properties	186-191
7.3.5.	Texture of yoghurt samples during storage period	192-93
7.3.6.	Sensory analysis of yoghurt	194-195
7.4.	Conclusion	196-197
7.5.	References	198-202
8.	Chapter VIII	202-206
	Summary, Conclusions and Future scope	
