

## TABLE OF CONTENTS

Title	Page no
Title page	
Dedication	i
Abstract	iii - x
Declaration	xi
Certificate of the Supervisor	xiii
Acknowledgement	xv
Table of Contents	xvii - xxxi
List of Tables	xxxiii - xxxv
List of Figures	xxxvii - xl
List of Abbreviations	xli – xliii
List of Symbols	xlv - xlvi
<b>Name of the Chapters and the sections under each chapter</b>	<b>Page No.</b>
<b>Chapter 1: Introduction</b>	<b>1 - 13</b>
1.1 Passion fruit juice and its challenges	3 – 4
1.2 Co-products of Passion fruit juice processing industries	4 – 5
1.3 Recovery of bioactive compounds	5 - 6
1.4 Emulsion based approach	6
1.5 Research gap	7
1.6 The objective of the present investigation	8
Bibliography	9 -13
<b>Chapter 2: Review of Literature</b>	<b>14 - 50</b>
2.1 Passion fruit	15 - 17
2.2 Passion fruit juice and its processing	17 - 20
2.3 Passion fruit peels	20 - 21
2.4 Passion fruit seed and its major parts	21

---

2.5	Dietary fibre and its extraction	22 - 23
2.6	Novel nonthermal preservative technique for juices	23 - 28
2.6.1	Ultrasonication	23 - 24
2.6.2	High pressure homogenization	24 - 26
2.6.3	Emulsion	26 - 28
2.7	Different extraction techniques	28 - 29
2.8	Vegetable oils as green solvents	29
2.9	Green extraction	30
2.10	Carotenoids and their extraction	30 - 32
2.11	Mathematical kinetics modelling in food technology	32 - 33
2.12	Optimization	33 -34
2.13	Bio-accessibility of carotenoids	34
	Bibliography	35 - 50
	<b>Chapter 3: Objective 1: To optimize the extraction of carotenoids from passion fruit peel using novel and green techniques and study its kinetics for industrial application</b>	<b>51 - 90</b>
3.1	Introduction	52 - 53
3.2	Materials and Methods	53 - 64
3.2.1	Chemicals	53
3.2.2	Preparation of sample	53 - 54
3.2.3	Ultrasonic assisted extraction (UAE) of PFP using vegetables oils	54
3.2.4	Microwave assisted extraction (MAE) of PFP using olive oil	54
3.2.5	Energy efficiency of UAE and MAE	54-55
3.2.6	Conventional extraction (CE) of PFP using OO	55
3.2.7	Soxhlet extraction of PFP	55
3.2.8	Total carotenoids content and $\beta$ -carotene content	56
3.2.9	Total phenolic content	56

---

---

3.2.10	Antioxidant activity (DPPH scavenging activity)	56 - 57
3.2.11	Experimental design	57 - 58
3.2.12	Optimization	58
3.2.13	Quality parameters of oil	59
3.2.14	Colors parameters	59
3.2.15	Pseudo second-order model carotenoids extraction kinetics	60 - 61
3.2.16	Phenomenological extraction kinetics model	61
3.2.17	Determination of effective diffusion coefficient during extraction	61 - 62
3.2.18	Biot number	62 - 63
3.2.19	Thermodynamics of UAE and CE	63 - 64
3.2.20	Statistical Analysis	64
3.3	Results and Discussion	64 - 84
3.3.1	Extraction yield of carotenoids	64 - 65
3.3.2	Effects of treatment time on carotenoids extraction of UAE	65 - 66
3.3.3	Effects of temperature on carotenoids extraction of UAE	66 - 67
3.3.4	Effects of solid to oil ratio on carotenoids extraction of UAE	67
3.3.5	Modelling and validation	68 - 69
3.3.6	Optimization	69 - 70
3.3.7	The effect of input parameters on recovery of carotenoids using MAE treatment with OO	70 - 72
3.3.8	Conventional extraction	72
3.3.9	Comparison of UAE, MAE, and conventional extraction methods	73 - 74
3.3.10	Physical and chemical properties of UOO and CEOO	74 - 76

---

3.3.11	Pseudo second order kinetics	76 - 78
3.3.12	Phenomenological kinetic modelling of extraction	78 - 81
3.3.13	Effective diffusion coefficient ( $D_e$ ), mass transfer coefficient ( $k_{mt}$ ), and Biot number ( $B_i$ ) estimation of carotenoids extraction process	81 - 83
3.3.14	Thermodynamics properties of UAE and CE	83 - 84
3.4	Conclusion	84
	Bibliography	85 - 90
<b>Chapter 4: To develop and evaluate Pickering nanoemulsion containing passion fruit peel-derived bioactive compounds and stabilized with starch nanoparticles</b>		<b>91 - 136</b>
4.1	Introduction	92 - 94
4.2	Materials and Methods	94 - 103
4.2.1	Materials	94
4.2.2.	Sample preparation	94 - 96
4.2.2.1	Preparation of ultrasonic-treated (US) starch nanoparticles	94 - 95
4.2.2.2	High pressure homogenization (HPH) of US-StNP	95
4.2.2.3	Preparation of carotenoids enriched StNPs based nanoemulsion	95
4.2.2.4	Preparation of carotenoids enriched StNPs	95 - 96
4.2.3	Particle size and polydispersity index of StNPs	96
4.2.4	Colour of StNPs	96
4.2.5	Flow properties of StNPs	96
4.2.6	Swelling power and solubility of StNPs	97
4.2.7	Fourier transform-infrared spectroscopy (FTIR) of StNPs	97
4.2.8	X-ray diffraction (XRD) of StNPs	97
4.2.9	Morphology of StNPs	98

---

4.2.10	Rheological properties of StNPs	98
4.2.11	Differential scanning calorimetry (DSC) of StNPs	98
4.2.12	Thermogravimetric analysis (TGA) of StNPs	98 - 99
4.2.13	The total carotenoids and $\beta$ -carotene content of nanoemulsion	99
4.2.14	Total phenolic content of nanoemulsion	99
4.2.15	Antioxidant activity (DPPH scavenging activity)	99
4.2.16	Heat and freeze-thaw stability of nanoemulsion	100
4.2.17	Oxidative stability of nanoemulsion	100
4.2.18	Effects of storage time and temperature on emulsion physical stability	100
4.2.19	Experimental design for optimization	100 - 101
4.2.20	Optimization	101
4.2.21	Methods adopted to obtain the thermal degradation kinetics	101 - 103
4.2.22	Hydroxyl groups estimation of StNPs using TGA	103
4.3	Results and Discussion	104 - 130
4.3.1	Distribution pattern of particle size	104 - 105
4.3.2	Colour of StNPs	105 - 106
4.3.3	Flow properties of StNPs	106 - 107
4.3.4	Swelling power and solubility of StNPs	107 - 108
4.3.5	FTIR absorbance of StNPs	108 - 109
4.3.6	Powder XRD analysis of StNPs	109 - 110
4.3.7	Morphological studies	110 - 112
4.3.8	Rheological properties	112 - 113
4.3.9	DSC thermograms of StNPs	113 - 115
4.3.10	TGA analysis of StNPs	115 - 117
4.3.11	Thermal decomposition Kinetics of starch samples	117 - 121
4.3.12	Determination of hydroxyl groups in StNPs using	121 - 122

---

---

	second mass loss from TGA	
4.3.13	Properties of the Pickering emulsion stabilized with starch nanoparticles	122 -
4.3.13.1	Effect of high pressure on particle size of the nanoemulsion	122 - 123
4.3.13.2	Effect of oil content on particle size of nanoemulsion	123 - 125
4.3.13.3	Effects of starch content on particle size of the nanoemulsion	125
4.3.13.4	Modelling and validation	125 - 126
4.3.13.5	Optimization	126 - 127
4.3.14	Properties of starch-based nanoemulsion	127 - 128
4.3.15	Heat and freeze thaw stability of nanoemulsion	128
4.3.16	Effects of antioxidants on oxidative stability of nanoemulsion	128 - 129
4.3.17	Quality of nanoemulsion on storage	129
4.3.17.1	Effect on antioxidant activity	129
4.3.17.2	Effect on emulsion stability	129 - 130
4.4	Conclusion	131
	Bibliography	131 - 136
	<b>Chapter 5: To optimize the extraction of oil and dietary fibre from seeds of passion fruit and study their extraction kinetics and properties.</b>	<b>137 - 184</b>
<b>Chapter 5A</b>	<b>Kinetic modelling of combined ultrasonication and soxhlet-based extraction of oil from passion fruit seed and its comparison and evaluation of physicochemical properties of optimized extracted oil</b>	<b>138 - 160</b>
5A.1	Introduction	138 - 140
5A.2	Materials and Methods	140 - 146
5A.2.1	Sample preparation	140

---

---

5A.2.2	Ultrasonic assisted extraction of oil from PFS	140
5A.2.3	Soxhlet method of oil extraction	141
5A.2.4	Dual UAE-Soxhlet extraction (UAES) of oil from PFS	141
5A.2.5	Proximate analysis of passion fruit seed	141
5A.2.5.1	Moisture content of PFS	141
5A.2.5.2	Total ash content of PFS	141
5A.2.5.3	Protein content of PFS	142
5A.2.5.4	Crude fibre content of PFS	142
5A.2.5.5	Total fat content of PFS	142
5A.2.5.6	Yield of oil	142
5A.2.6	Physicochemical properties of seed oil	143
5A.2.7	Total carotenoid content of PFSO	143
5A.2.8	Total phenolic content and HPLC analysis of phenolic acid in PFSO	143
5A.2.9	Fatty acids composition	143 - 144
5A.2.10	Calculated oxidizability value	144
5A.2.11	Antioxidant activity	144
5A.2.11.1	DPPH radical scavenging activity	144
5A.2.11.2	ABTS activity	144 - 145
5A.2.12	Phenomenological kinetics model	145 - 146
5A.2.13	Statistical analysis	146
5A.3.	Results and Discussion	147 - 156
5A.3.1	Proximate analysis of PFS	147
5A.3.2	Effect of the solid-liquid ratio on ultrasonic extraction of PFSO	147 - 148
5A.3.3	Effect of extraction temperature on ultrasonic extraction of PFSO	148
5A.3.4	Effect of extraction time on ultrasonic extraction of PFSO	148

---

---

5A.3.5	Soxhlet extraction of PFSO	148 - 149
5A.3.6	UAE-Soxhlet extraction of PFSO	150
5A.3.7	Kinetic modelling of extraction of PFSO	150
5A.3.8	Phenomenological model	150 - 154
5A.3.9	Physicochemical properties of PFSO	154 - 155
5A.3.10	Fatty acids composition	155 - 156
5A.4	Conclusion	156 - 157
5A.C	Bibliography	157 - 160
<b>Chapter 5B</b>	<b>Properties of dietary fibre from passion fruit seed obtained through individual and combined alkaline and ultrasonication extraction techniques</b>	<b>161 - 184</b>
5B.1	Introduction	161 - 162
5B.2	Materials and methods	163 - 167
5B.2.1	Sample preparation	163
5B.2.2	Methods of extraction of DF from DPFS	163 - 164
5B.2.2.1	Alkaline extraction method of DFs	163
5B.2.2.2	Ultrasound-assisted alkaline extraction method of DFs	163
5B.2.2.3	Ultrasound treatment of DFs	164
5B.2.3	Physicochemical properties of PFS and DPFS	164
5B.2.4	Characterization of DFs	164 - 165
5B.2.4.1	Morphology of DFs	164
5B.2.4.2	Fourier transform-infrared spectroscopy (FTIR) of DFs	164
5B.2.4.3	X-ray diffraction (XRD) of DFs	164
5B.2.4.4	Thermogravimetric analysis (TGA) of DFs	165
5B.2.5	Physicochemical properties of DFs	165 - 166
5B.2.5.1	Swelling capacity (SC), water holding capacity (WHC), and oil holding capacity	165 - 166

---



---

	(OHC)	
	5B.2.5.2	Cation-exchange capacity (CEC) of DFs 165 - 166
	5B.2.5.3	Emulsifying capacity (EC) of DFs 166
5B.2.6		Functional properties of DFs 166 - 167
	5B.2.6.1	Glucose-adsorption capacity (GAC) and $\alpha$ - amylase inhibitory effect of DFs 166
	5B.2.6.2	Cytotoxicity assay of DFs 166 - 167
5B.2.7		Experimental design of alkaline-ultrasonic extraction 167
5B.2.8		Optimization 167
5B.2.9		Statistical Analysis 167
5B.3		Results and Discussion 168 - 180
	5B.3.1	Physicochemical properties of PFS and DFPS 167 - 168
	5B.3.2	Optimization of UAE-alkaline extraction of DFs 168 - 171
	5B.3.3	Characterization of ADF, AUDF and UDF 171 - 177
	5B.3.3.1	SEM analysis of DFs 171 - 172
	5B.3.3.2	FT-IR spectra of DFs 172 - 174
	5B.3.3.3	XRD of DFs 174 - 176
	5B.3.3.4	TGA analysis of DFs 176 - 177
	5B.3.4	Physicochemical properties of extracted dietary fibre 177 - 178
	5B.3.5	Functional properties of extracted DFs 178 - 180
	5B.3.5.1	The glucose-adsorption capacity and $\alpha$ - amylase activity 178 - 179
	5B.3.5.2	Cytotoxicity MTTs assay 179 - 180
5B.4		Conclusion 180
		Bibliography 180 - 184
		<b>Chapter 6: To evaluate the quality of blended fruit juice of passion fruit and other fruits incorporated with the oil and dietary fibre extracts from the seeds of the fruit, and study the bio-accessibility of carotenoids. 185 - 282</b>
		<b>Chapter Effect of thermosonication treatment on passion fruit juice, 186 - 211</b>

---

---

<b>6A</b>	<b>ann/ga optimization and its predictive modelling for shelf life and quality changes during storage</b>	
6A.1	Introduction	186 - 188
6A.2	Material and Methods	188 - 194
6A.2.1	Juice preparation	188
6A.2.2	Thermosonication (TS) of SPFJ	188
6A.2.3	Physicochemical analysis of juices	188 - 191
6A.2.3.1	Yield determination	188
6A.2.3.1	Proximate analysis	188
6A.2.3.3	Total soluble solid (TSS) content	189
6A.2.3.4	pH value	189
6A.2.3.5	Total titratable acidity (TTA)	189
6A.2.3.6	Fat content (Soluble)	189
6A.2.3.7	Ascorbic acid (AA) content	189 - 190
6A.2.3.8	Total phenolic content	190
6A.2.3.9	Total sugars content	190
6A.2.3.10	DPPH scavenging activity	191
6A.2.3.11	Total carotenoids content (TCC) and $\beta$ -carotene content	191
6A.2.3.12	Microbial analysis/ Total plate count (TPC)	191
6A.2.4	Storage study of juices	191
6A.2.4.1	Shelf-life estimation using predictive modelling	191 -
6A.2.4.2	Quality parameters during storage	191
6A.2.5	Artificial neural network (ANN) of TS of SPFJ	191 -192
6A.2.6	ANN/Genetic algorithm (GA) optimization of TS on SPFJ	192
6A.2.7	Predictive modelling / mathematical modelling for shelf-life prediction	192 -193

---

---

6A.2.8	Sensory evaluation	193
6A.2.9	Statistical analysis	193 -194
6A.3	Results and Discussion	194 - 207
6A.3.1	Physicochemical parameters of SPFJ	194 - 195
6A.3.2	Effects of TS on TPC	195
6A.3.3	Effect of TS on total phenolic content	195 -197
6A.3.4	Effect of TS on $\beta$ - carotene content	197
6A.3.5	Optimization of TS on SPFJ	197 - 200
	6A.3.5.1 ANN Prediction	197 - 200
	6A.3.5.2 Validation of ANN	200
6A.3.6	ANN/GA Optimization	200
6A.3.7	Baranyi equation as predictive modelling	200 - 202
6A.3.8	Shelf-life estimation	202 - 203
6A.3.9	Changes in quality parameters of SPFJ and OPFJUS during refrigerated storage	203 - 207
	6A.3.9.1 Effect of TSS	203
	6A.3.9.2 Effect of pH	206
	6A3.9.3 Effect of total titratable acidity	206
	6A.3.9.4 Effect of ascorbic acid (AA)	206
	6A.3.9.5 Effect of total phenolic content	206 - 207
	6A.3.9.6 Effect of DPPH scavenging activity	207
	6A.3.9.7 Effect of $\beta$ -carotene	207
	6A.3.9.8 Sensory analysis	207
6A.4	Conclusion	207 - 208
	Bibliography	208 - 211
<b>Chapter 6B</b>	<b>Fuzzy logic sensory optimization of passion fruit-based blended beverage</b>	212 - 225
6B.1	Introduction	212
6B.2	Materials and Methods	213 - 215

---

6B.2.1	Samples	213
6B.2.2	Juice extraction	213
6B.2.3	Sensory evaluation	213
6B.2.4	Application of fuzzy logic on sensory data	214 - 215
6B.3.	Results and Discussion	216 - 224
6B.3.1	Sensory evaluation of blended juices	216 - 217
6B.3.2	Triplets for sensory quality of blended beverage samples	217 - 218
6B.3.3	Determination of important quality attributes based on triplet values	218 - 219
6B.3.4	Triplets for overall sensory scores of the blended beverage samples	219 - 220
6B.3.5	Ranking of blended beverage on sensory fuzzy scale	220 - 221
6B.3.6	Ranking of blended beverage on the basis of quality attributes	221 - 222
6B.3.7	Similarity analysis for blended beverage samples on standard fuzzy scale	222
6B.3.8	Membership function of overall sensory scores on standard fuzzy scale	222 - 223
6B.3.9	Similarity values for blended beverage samples	223 - 224
6B.4	Conclusion	224
	Bibliography	224 - 225
<b>Chapter 6C</b>	<b>Development of Pickering emulsion from passion fruit seed fibre and seed oil and passion fruit peel derived carotenoids-enriched oil based and its optimization</b>	<b>226 - 236</b>
6C.1	Introduction	226
6C.2	Materials and Methods	227 - 228
6C.2.1	Samples	227
6C.2.2	Characterization of nanoemulsion	227
6C.2.3	Preparation of carotenoids enriched fibre-based	227 - 228

---

	Pickering nanoemulsion	
6C.2.4	Experimental design for optimization	228
6C.2.5	Optimization	228
6C.3	Results and discussion	228 - 234
6C.3.1	Effects of pressure on practice size	228 - 230
6C.3.2	Effects of fibre content on particle size of the nanoemulsion	230 - 231
6C.3.3	Effects of oil amount on emulsion size	232 - 233
6C.3.4	Modeling and validation	233
6B.3.5	Optimization of responses	233 - 234
6C.4	Conclusion	234
	Bibliography	234 - 236
<b>Chapter 6D</b>	<b>Effect of thermal pasteurization and high-pressure homogenization on quality of beverages and development and evaluation of an optimized passion fruit based-beverage enriched with Pickering nanoemulsion of fibre and oil extracted from its byproducts</b>	<b>237 - 282</b>
6D.1	Introduction	237 - 238
6D.2	Materials and Methods	238 - 244
6D.2.1	Sample preparation	238
6D.2.2	HPH treatment of samples	239
6D.2.3	Heat treatment (HT) of blended juice	239
6D.2.4	Blended beverage added with carotenoids enriched fibre-based Pickering nanoemulsions	239
6D.2.5	Physicochemical properties of treated and untreated beverages	239 - 240
6D.2.5.1	ABTS free radical scavenging activity assay of samples	240
6D.2.6	Suspension stability / Cloudiness	240
6D.2.7	Particle size	240

---

---

6D.2.8	HPLC analysis of carotenoids of samples	241
6D.2.9	HPLC analysis of phenolic acid of samples	241
6D.2.10.	Storage study of samples	242
6D.2.11	Dynamic <i>In-vitro</i> digestion model of samples	242 - 243
6D.2.11.1	Mouth phase	242
6D.2.11.2	Simulated gastric fluid phase	242 - 243
6D.2.11.3	Simulated Intestinal phase	243
6D.2.11.4	Bio-accessibility determination	243
6D2.12	Statistical analysis	243 - 244
6D.3	Results and Discussion	244 - 282
6D.3.1	Changes in quality parameters after treatments	244 - 264
6D.3.1.1	pH value	244
6D.3.1.2	Total titratable acidity (TTA)	246
6D.3.1.3	Total soluble solid (TSS)	246
6D.3.1.4.	Total sugar	246 - 247
6D.3.1.5	Ascorbic acid (AA)	247 - 248
6D.3.1.6	Total phenolic content and polyphenol profile	248 - 251
6D.3.1.7	Total carotenoids content (TCC), $\beta$ -carotene content and carotenoids profiles	251 - 253
6D.3.1.8	Antioxidant activity	253 - 255
6D.3.1.9	Colour value	255 - 257
6D.3.1.10	Suspension stability/ cloudiness	257 - 258
6D.3.1.11	Microbiological analysis (Total plate count)	258 - 259
6D.3.1.12	Particle size	259 - 260
6D.3.1.13	<i>In-vitro</i> bio-accessibility of carotenoids	260 - 262
6D.3.1.14	Sensory parameters	262 - 263
6D.3.1.15	Cytotoxicity MTTS assay	263 - 264

---

---

6D.3.2	Storage study of BSPFJT, BSPFJTT, and BNESPFJT	264 - 273
6D.3.2.1	PH and total titratable acidity (TTA)	264 - 265
6D.3.2.2	Total sugar and TSS	265
6D.3.2.3	Ascorbic acid (AA)	265 - 266
6D.3.2.4	Total phenolic content	266 - 269
6D.3.2.5	Total carotenoids content	269 - 270
6D.3.2.6	Storage of antioxidant	270
6D.3.2.7	Colour value	270 - 271
6D.3.2.8	Storage stability cloud stability	271
6D.3.2.9	Microbial load and shelf life	271 - 273
6D.4	Conclusion	273
	Bibliography	273 - 282
	<b>Chapter 7: Conclusion and future scope of work</b>	<b>283 - 291</b>
7.1	Conclusion	284
7.2	Chapter 3	285
7.3	Chapter 4	286
7.4	Chapter 5	287
7.5	Chapter 6	288 - 289
7.6	Future scope of work	289
	List of Publications and Conferences	290 - 291

---