

Contents

Contents	Page No.
Abstract	i-iv
Declaration	v
Certificate	vi
Acknowledgements	vii-viii
Contents	ix-xvi
List of tables	xvii-xviii
List of figures	xix-xxv
List of abbreviations	xxvi-xxvii
List of symbols	xxviii-xxix
Chapter 1 Introduction	
1.1 Overall introduction	1.1
1.2 <i>Euryale ferox</i>	
1.2.1 Distribution of <i>Euryale ferox</i>	1.2
1.2.2 Description of <i>Euryale ferox</i>	1.3
1.2.3 <i>Euryale ferox</i> Seed	1.3
1.3 Starch	1.4
1.4 Resistant Starch	1.4-1.5
1.5 Polyphenols	1.5-1.6
1.6 Starch-polyphenol complex	1.6
1.7 Pre-gelatinization method	1.6-1.7
1.8. Diabetes mellitus	1.7
1.9 DPP IV inhibition	1.7-1.8
1.10 Inhibition of α -amylase and α -glucosidase	1.8
1.11 Glucose uptake assay	1.8-1.9
1.12 Anti-inflammatory activity	1.9
1.13 Bread	1.9
1.14 Gluten free bread	1.10
1.15 Swarm intelligence supervised neural network (SISNN)	1.10
1.16 Hypothesis	1.11
1.17 Gap of Research	1.11

1.18	Objectives	1.11
	Bibliography	1.12-1.17

Chapter 2 To extract and characterize starch of the kernel of *Euryale ferox* seed

2.1	Introduction	2.1-2.2
2.2	Materials and methods	
2.2.1	Sample collection and preparation	2.2
2.2.2	Determination of starch content and isolation of starch from <i>Euryale ferox</i> kernel	2.2
2.2.3	Chemical composition and amylose content	2.3
2.2.4	Morphology of <i>Euryale ferox</i> kernel starch	2.3
2.2.5	X-ray diffraction (XRD) pattern and relative crystallinity	2.3
2.2.6	Fourier transform infra-red (FTIR) spectroscopy analysis	2.3
2.2.7	Gelatinization properties	2.4
2.2.8	Thermogravimetric analysis	2.4
2.2.9	Swelling power and solubility	2.4
2.2.10	Pasting properties	2.4-2.5
2.2.11	Color properties	2.5
2.2.12	<i>In vitro</i> starch digestibility (IVSD)	2.5-2.6
2.2.13	Statistical analysis	2.6
2.3	Results and discussion	
2.3.1	Specimen identification	2.6
2.3.2	Starch content, starch yield, chemical composition and amylose contents	2.6-2.8
2.3.3	Morphology of <i>Euryale ferox</i> kernel starch	2.8-2.9
2.3.4	X-ray diffraction (XRD) pattern and relative crystallinity	2.9-2.10
2.3.5	Fourier transform infra-red (FTIR) spectroscopy analysis	2.10-2.11
2.3.6	Gelatinization properties	2.11-2.12

2.3.7	Thermogravimetric analysis	2.12
2.3.8	Swelling power and solubility	2.13-2.14
2.3.9	Pasting properties	2.14-2.16
2.3.10	Color properties	2.16
2.3.11	<i>In vitro</i> starch digestibility (IVSD)	2.17
2.4	Conclusion	2.18
	Bibliography	2.18-2.22

Chapter 3 To extract polyphenols from the shell of *Euryale ferox* seed and evaluate its bioactivities

3.1	Introduction	3.1-3.2
3.2	Materials and methods	
3.2.1	Sample preparation	3.2
3.2.2	Extraction of phytochemicals	
3.2.2.1	Maceration	3.3
3.2.2.2	Decoction	3.3
3.2.2.3	Ultrasound assisted extraction	3.3
3.2.2.4	Microwave assisted extraction	3.3
3.2.3	Purification of polyphenols	3.3-3.4
3.2.4	Extraction yield	3.4
3.2.5	Determination of phytochemicals	
3.2.5.1	Total phenolic content	3.4
3.2.5.2	Total flavonoid content	3.4-3.5
3.2.6	Antioxidant activity	
3.2.6.1	DPPH radical scavenging activity	3.5
3.2.6.2	ABTS radical scavenging activity	3.5-3.6
3.2.6.3	FRAP reducing activity	3.6
3.2.7	Phytochemicals screening by HR-LCMS analysis	3.6
3.2.8	Estimation of polyphenols by RP-HPLC	3.6-3.7
3.2.9	<i>In-vitro</i> anti-diabetic study	
3.2.9.1	<i>In vitro</i> α -amylase inhibition activity	3.7
3.2.9.2	<i>In vitro</i> α -glucosidase inhibition activity	3.7-3.8

	3.2.9.3	Determination of DPP-IV inhibition activity	3.8
	3.2.10	<i>In vitro</i> glucose uptake assay on L6 rat skeletal muscle cell line	
	3.2.10.1	MTT assay of <i>Euryale ferox</i> seed shell extract treated L6 cell	3.9
	3.2.10.2	Glucose (2-NBDG) uptake assay	3.9-3.10
	3.2.11	<i>In-vitro</i> anti-inflammatory study of <i>Euryale ferox</i> seed shell extract	
	3.2.11.1	MTT assay of <i>Euryale ferox</i> seed shell extract treated THP-1 cell	3.10
	3.2.11.2	<i>In vitro</i> LPS induced inflammation study in THP-1 macrophage	3.10-3.11
	3.2.12	Statistical analysis	3.11
3.3		Results and discussion	
	3.3.1	Extraction yield	3.12
	3.3.2	Effect of extraction method on the total phenolic content (TPC) and total flavonoid content (TFC)	3.12-3.14
	3.3.3	<i>In vitro</i> antioxidant assays	
	3.3.3.1	DPPH radical scavenging activity of <i>Euryale ferox</i> seed shell extract	3.14-3.15
	3.3.3.2	ABTS scavenging activity of <i>Euryale ferox</i> seed shell extract	3.15-3.16
	3.3.3.3	FRAP reducing activity of <i>Euryale ferox</i> seed shell extract	3.16
	3.3.4	Effect of extraction method on the antioxidant activity	3.17-3.18
	3.3.5	Phytochemical screening by HR-LCMS analysis	3.18-3.20
	3.3.6	Estimation of polyphenols by RP-HPLC	3.21-3.22
	3.3.7	<i>In vitro</i> antidiabetic study	
	3.3.7.1	α -amylase inhibition assay of <i>Euryale ferox</i> seed shell extract	3.23

3.3.7.2	α -glucosidase inhibition assay of <i>Euryale ferox</i> seed shell extract	3.24-3.25
3.3.7.3	DPP IV inhibition assay of <i>Euryale ferox</i> seed shell extract	3.25-3.26
3.3.8	Glucose uptake assay of <i>Euryale ferox</i> seed shell extract	
3.3.8.1	Viability of L6 cell treated with <i>Euryale ferox</i> seed shell extract	3.26-3.27
3.3.8.2	Glucose (2-NBDG) uptake assay	3.27-3.28
3.3.9	Anti-inflammatory activity of <i>Euryale ferox</i> seed shell extract	
3.3.9.2	Viability of THP-1 cell treated with <i>Euryale ferox</i> seed shell extract	3.29
3.3.9.3	<i>In vitro</i> LPS induced inflammatory study in THP-1 macrophage	3.30-3.33
3.4	Conclusions	3.33-3.34
	Bibliography	3.34-3.37

Chapter 4 To develop and characterize starch-polyphenol complex from *Euryale ferox* seed

4.1	Introduction	4.1-4.2
4.2	Materials and methods	4.3
4.2.1	Preparation of <i>Euryale ferox</i> kernel starch-polyphenols complex	4.3
4.2.2	Characterization of <i>Euryale ferox</i> kernel starch-polyphenols complexation	
4.2.2.1	X-ray diffraction (XRD) analysis	4.3
4.2.2.2	Fourier transform infrared (FT-IR) spectroscopy	4.4
4.2.2.3	¹ H NMR spectroscopy	4.4
4.2.2.4	Differential Scanning Calorimetry (DSC)	4.4

4.2.2.5	Scanning electron microscopy (SEM) analysis	4.4
4.2.3	<i>In vitro</i> starch digestibility	
4.2.3.1	Determination of starch fraction	4.4-4.5
4.2.3.2	Predicted glycemic index	4.5-4.6
4.2.4	DPPH radical scavenging activity	4.6
4.2.5	Color characteristics	4.6
4.2.6	Swelling power and solubility	4.6-4.7
4.2.7	Statistical analysis	4.7
4.3	Results and Discussion	
4.3.1	Characterization of <i>Euryale ferox</i> kernel starch-polyphenols complexation	
4.3.1.1	X-ray diffraction (XRD)	4.10-4.18
4.3.1.2	Fourier transform infrared (FT-IR) spectroscopy	4.18-4.26
4.3.1.3	¹ H NMR spectroscopy	4.26-4.35
4.3.1.4	Differential Scanning Calorimetry (DSC) analysis	4.36-4.46
4.3.1.5	Scanning electron microscopy (SEM) analysis	4.46-4.50
4.3.2	<i>In vitro</i> starch digestibility	
4.3.2.1	Starch fractions	4.50-4.52
4.3.2.2	Predicted glycemic index (pGI)	4.53-4.55
4.3.3	DPPH radical scavenging activity	4.56-4.57
4.3.4	Color attributes	4.57-4.58
4.3.5	Swelling power and solubility	4.59-4.60
4.4	Conclusion	4.61
	Bibliography	4.61-4.66

Chapter 5 To study the effect of *Euryale ferox* seed shell extract addition on the *in-vitro* starch digestibility and predicted glycemic index of wheat-based bread

5.1	Introduction	5.1-5.2
5.2	Material and methods	

5.2.1	Preparation of bread	5.3
5.2.2	Specific volume of bread	5.3
5.2.3	Texture profile analysis of bread	5.3-5.4
5.2.4	Color properties of bread	5.4
5.2.5	<i>In vitro</i> starch digestibility and predicted glycemic index	5.4-5.5
5.2.5.1	Mathematical modeling approach	5.5
5.2.5.2	Swarm intelligence supervised neural network (SISNN) approach	5.6-5.7
5.2.6	Sensory evaluation of the bread samples	
5.2.6.1	Collection of sensory scores	5.7-5.8
5.2.6.2	Analysis of the sensory scores using POM supervised PCA technique	5.8-5.9
5.2.7	Statistical analysis	5.9
5.3	Results and discussion	
5.3.1	Specific volume of bread	5.9-5.10
5.3.2	Texture profile analysis of bread	5.10-5.11
5.3.3	Color properties of bread	5.11-5.12
5.3.4	<i>In vitro</i> starch digestibility and predicted glycemic index	5.12
5.3.4.1	Mathematical modeling approach for determination of pGI	5.13-5.16
5.3.4.2	SISNN based modeling approach	5.16-5.19
5.3.4.3	Sensitivity analysis	5.19-5.20
5.3.5	Sensory evaluation by POM supervised PCA technique	5.20-5.23
5.4	Conclusion	5.23
	Bibliography	5.24-5.27

Chapter 6 To incorporate starch-polyphenol complex from *Euryale ferox* seed in gluten free bread formulation

6.1	Introduction	6.1
6.2	Material and methods	

6.2.1	Preparation of gluten-free bread	6.1-6.2
6.2.2	Moisture retention, specific volume	6.2
6.2.3	Texture profile analysis	6.2
6.2.4	Color attributes	6.2
6.2.5	<i>In vitro</i> starch digestibility	
	6.2.5.1 Starch fractions	6.2
	6.2.5.2 Predicted glycemic index (pGI)	6.2
6.2.6	Extraction of phytochemicals	6.3
6.2.7	Total phenolic content	6.3
6.2.8	Total flavonoid content	6.3
6.2.9	DPPH radical scavenging activity	6.3
6.2.10	Statistical analysis	6.4
6.3	Results and discussion	
6.3.1	Specific volume of gluten free bread	6.4-6.5
6.3.2	Moisture retention and baking loss of gluten-free bread	6.6
6.3.3	Textural attributes of gluten-free bread	6.7
6.3.4	Color attributes of gluten free bread	6.8
6.3.5	<i>In vitro</i> starch digestibility and predicted glycemic index of gluten free bread	
	6.3.5.1 Starch fractions	6.9
	6.3.5.2 Predicted glycemic index	6.10-6.11
6.3.6	Total phenolic and flavonoid content	6.11-6.12
6.3.7	Antioxidant activity	6.13
6.3.8	Sensory evaluation	6.14
6.4	Conclusions	6.15
	Bibliography	6.15-6.18
 Chapter 7 Conclusions and future scope of work		
7.1	Conclusions	7.1-7.3
7.2	Future scope of the present investigation	7.3
	List of Publications	a-b