Contents

| | | | Contents | Page No. | | |
|----------------------|----------|----------|---|-------------------|--|--|
| Abstract i- | | | | | | |
| Abstract Declaration | | | | | | |
| | tificate | | | vi vii | | |
| | | | 0 | vii viii-ix | | |
| | itents | dgement | 5 | v111-1X X-Xiii | | |
| | of Tab | alog | | x-xiii xiv | | |
| | | | | | | |
| | of Fig | | tuo duoti on | XV | | |
| 1. | Cnap | ner 1 in | troduction | 1-9 | | |
| | Refer | rences | | 6-9 | | |
| 2. | Chap | ter 2 Re | eview of Literature | 10-30 | | |
| | 2.1 | Pigmer | nted rice and its nutritional values | 10 | | |
| | 2.2 | Physico | o-chemical and phytochemical properties | 11 | | |
| | 2.3 | Extract | tion of Phytochemicals | 12 | | |
| | 2.4 | Ultrasc | ound-assisted extraction | 14 | | |
| | 2.5 | Double | emulsion | 15 | | |
| | 2.6 | Compl | ex Coacervation | 16 | | |
| | 2.7 | Applica | ation in food model | 18 | | |
| | 2.8 | Coconi | ut water | 20 | | |
| | Refer | ences | | 21-30 | | |
| 3. | Chap | oter 3 | Varietal differences in physico-chemical and | 31-55 | | |
| | - | | al properties of pigmented rice | | | |
| | 3.1 | Introdu | | 31 | | |
| | 3.2 | Materia | al and methods | 33 | | |
| | | 3.2.1 | Materials | 33 | | |
| | | 3.2.2 | Proximate analysis | 33 | | |
| | | 3.2.3. | Physico-chemical properties | 34 | | |
| | | 3.2.4 | Phytochemical properties of rice and its bran | 35 | | |
| | | | 3.2.4.1 Extract preparation | 35 | | |
| | | | 3.2.4.2 Total phenolic content (TPC) | 35 | | |
| | | | 3.2.4.3 Total flavonoid content (TFC) | 36 | | |
| | | | 3.2.4.4 DPPH radical scavenging activity | 36 | | |
| | | | 3.2.4.5 Total anthocyanin content (TAC) | 37 | | |
| | | 3.2.5 | Cooking properties | 37 | | |
| | | 2.2.3 | 3.2.5.1 Minimum cooking time | 37 | | |
| | | | 3.2.5.2 Elongation ratio | 37 | | |
| | | | 3.2.5.3 Water uptake ratio | 38 | | |
| | | | 3.2.5.4 Gruel solid loss | 38 | | |
| | | 3.2.6 | Texture Analysis | 38 | | |
| | | | • | | | |
| | | 3.2.7 | Color | 38 | | |

| | 3.2.9 | Statistica | 1 Analysis | 20 |
|-------|--|--|---|---|
| | 3.4.7 | Statistica | n Anarysis | 39 |
| 3.3 | Results | and discu | ssion | 39 |
| | 3.3.1 | Proximat | te composition of pigmented rice | 39 |
| | 3.3.2 | Physicoc | hemical properties | 40 |
| | 3.3.3 | Phytoche | emical properties | 43 |
| | 3.3.4 | Cooking | properties | 45 |
| | 3.3.5 | Texture A | Analysis | 47 |
| | 3.3.6 | Color va | lues | 47 |
| | 3.3.7 | Pasting p | properties | 49 |
| 3.4 | Conclu | sions | | 51 |
| Refer | ences | | | 52-55 |
| Chap | ter 4 Ul | trasound- | assisted extraction of anthocyanin-rich extract | 56-79 |
| from | black | rice bra | n and its stability study under different | |
| temp | erature | and pH | | |
| 4.1 | Introdu | ction | | 56 |
| 4.2 | Materia | als and Me | thods | 59 |
| | 4.2.1 | Rice sam | ples, chemicals and reagents | 59 |
| | 4.2.2. | Prelimina | ary trial | 59 |
| | 4.2.3 | Extractio | on process | 60 |
| | | 4.2.3.1 | Ultrasound-assisted extraction of phenolics, | 60 |
| | | | flavonoids and anthocyanin | |
| | | 4.2.3.2 | Experimental design and optimization | 60 |
| | 4.2.4 | Extract a | nalysis | 61 |
| | | 4.2.4.1 | Total phenolic content | 61 |
| | | 4.2.4.2 | Total flavonoid content | 61 |
| | | 4.2.4.3 | Antioxidant activity | 61 |
| | | 4.2.4.4 | Total anthocyanin content | 62 |
| | 4.2.5 | Kinetics | study of anthocyanin extraction | 62 |
| | 4.2.6 | Stability | study of phytochemicals | 63 |
| | | 4.2.6.1 | Effect of temperature | 63 |
| | | 4.2.6.2 | Effect of pH | 63 |
| | | 4.2.6.3 | | 63 |
| | 4.2.7 | Statistica | • | 64 |
| 4.3 | Results | | • | 64 |
| | 4.3.1 | Phytoche | emical extraction and its antioxidant activity | 64 |
| | | - | • | 64 |
| | | | | 67 |
| | | content | r i i i i i i i i i i i i i i i i i i i | |
| | 4.3.4 | Effect of | | 68 |
| | 4.3.5 | | • | 70 |
| | 3.4 Refer Chap from temp 4.1 4.2 | 3.3 Results 3.3.1 3.3.2 3.3.3 3.3.4 3.3.5 3.3.6 3.3.7 3.4 Conclu References Chapter 4 UI from black temperature 4.1 Introdu 4.2 Materia 4.2.1 4.2.2 4.2.3 4.2.4 4.2.4 4.2.5 4.2.6 4.2.6 | 3.3.1 Proximat 3.3.2 Physicoc 3.3.3 Phytoche 3.3.4 Cooking 3.3.5 Texture A 3.3.6 Color val 3.3.7 Pasting p 3.4 Conclusions References Chapter 4 Ultrasound- from black rice bratemperature and pH 4.1 Introduction 4.2 Materials and Me 4.2.1 Rice sam 4.2.2 Prelimina 4.2.3 Extraction 4.2.3.1 4.2.3.2 4.2.4 Extract a 4.2.4.1 4.2.4.2 4.2.4.3 4.2.4.4 4.2.5 Kinetics 4.2.6.1 4.2.6.2 4.2.6.3 4.2.7 Statistical 4.3.1 Phytoche 4.3.2 Model fire 4.3.3 Effect of content 4.3.4 Effect of antioxidal | 3.3.1 Proximate composition of pigmented rice 3.3.2 Physicochemical properties 3.3.3 Phytochemical properties 3.3.4 Cooking properties 3.3.5 Texture Analysis 3.3.6 Color values 3.3.7 Pasting properties 3.4 Conclusions References Chapter 4 Ultrasound-assisted extraction of anthocyanin-rich extract from black rice bran and its stability study under different temperature and pH 4.1 Introduction 4.2 Materials and Methods 4.2.1 Rice samples, chemicals and reagents 4.2.2 Preliminary trial 4.2.3 Extraction process 4.2.3.1 Ultrasound-assisted extraction of phenolics, flavonoids and anthocyanin 4.2.4.2 Experimental design and optimization 4.2.4.1 Total phenolic content 4.2.4.2 Total flavonoid content 4.2.4.3 Antioxidant activity 4.2.4.4 Total anthocyanin content 4.2.5 Kinetics study of anthocyanin extraction 4.2.6 Stability study of phytochemicals 4.2.6.1 Effect of pH 4.2.6.2 Effect of pH 4.2.6.3 Effect of color / Characterization of anthocyanin extracts in various pH 4.2.7 Statistical analysis 4.3 Results and Discussion 4.3.1 Phytochemical extraction and its antioxidant activity 4.3.2 Model fitting 4.3.3 Effect of process variables on phenolic and flavonoid content 4.3.4 Effect of process variables on anthocyanin content and antioxidant activity |

| | | 4.3.6 | Extraction kinetics | 70 |
|----|-------|-----------|--|--------|
| | | 4.3.7 | Stability study | 71 |
| | | | 4.3.7.1 Effect of temperature | 71 |
| | | | 4.3.7.2 Effect of pH on anthocyanin | 73 |
| | | | 4.3.7.3 Color stability of anthocyanin at different pH | 74 |
| | 4.4 | Conclu | isions | 75 |
| | Refer | ences | | 76-79 |
| 5. | Chap | oter 5 I | Microencapsulation of anthocyanin-rich extract using | 80-106 |
| | doub | le emi | ulsion complex coacervation technique and its | |
| | chara | acterizat | tion | |
| | 5.1 | Introdu | action | 80 |
| | 5.2 | Materia | als and methods | 83 |
| | | 5.2.1 | Raw materials and chemicals | 83 |
| | | 5.2.2 | Extraction and concentration of black rice bran | 83 |
| | | | anthocyanin | |
| | | 5.2.3 | Dual emulsion (W/O/W) | 84 |
| | | 5.2.4 | Complex coacervation process | 84 |
| | | 5.2.5 | Characterization of the extracts using HPLC analysis | 85 |
| | | 5.2.6 | Characterization | 86 |
| | | | 5.2.6.1 Moisture content | 86 |
| | | | 5.2.6.2 Hygroscopicity | 86 |
| | | | 5.2.6.3 Solubility | 86 |
| | | | 5.2.6.4 Encapsulation efficiency | 86 |
| | | | 5.2.6.5 Surface morphology | 87 |
| | | | 5.2.6.6 FT-IR | 87 |
| | | | 5.2.6.7 Thermal stability | 87 |
| | | | 5.2.6.8 XRD | 87 |
| | | | 5.2.6.9 Stability study of anthocyanin | 88 |
| | | 5.2.7 | Statistical analysis | 88 |
| | 5.3 | Results | s and discussion | 88 |
| | | 5.3.1 | Preliminary trials | 88 |
| | | 5.3.2 | Effect of hydrocolloids on microcapsules | 91 |
| | | 5.3.3 | • | 91 |
| | | 5.3.4 | Physicochemical characterization of the coacervated | 92 |
| | | | microcapsules | |
| | | | 5.3.4.1 Moisture content | 92 |
| | | | 5.3.4.2 Hygroscopicity | 92 |
| | | | 5.3.4.3 Solubility | 93 |
| | | | 5.3.4.4 Encapsulation efficiency | 93 |
| | | 5.3.5 | Microstructure | 94 |
| | | 5.3.6 | Infrared Spectroscopy of the coacervated microcapsule | 96 |
| | | | powders | |
| | | 5.3.7 | XRD | 97 |

| | | 5.3.8 | Thermal stability | 98 | | |
|----|-------|---|--|----------------|--|--|
| | | 5.3.9 | Stability study of anthocyanin | 99 | | |
| | 5.4 | Conclu | sions | 101 | | |
| | Refer | rences | | 102-106 | | |
| 6. | Chap | oter 6 | Development of functional coconut water jelly by | 107-124 | | |
| | incor | porating | g anthocyanin microcapsules and its characterization | | | |
| | 6.1 | Introdu | ction | 107 | | |
| | 6.2 | Materia | al and methods | 108 | | |
| | | 6.2.1 | Raw materials and chemicals required | 108 | | |
| | | 6.2.2 | Preparation and production of anthocyanin-loaded coconut water jelly | 109 | | |
| | | 6.2.3 | Judging the end point of jelly | 109 | | |
| | | 6.2.4 | Determination of moisture content | 109 | | |
| | | 6.2.5 | Determination of pH value, titratable acidity and total soluble solids | 110 | | |
| | | 6.2.6 | Determination of texture profile | 110 | | |
| | | 6.2.7 | Determination of color | 110 | | |
| | | 6.2.8 | Sensory analysis | 111 | | |
| | | 6.2.9 | Determination of total phenolic content | 111 | | |
| | | 6.2.10 | Determination of total anthocyanin content | 111 | | |
| | | 6.2.11 | Antioxidant activity | 112 | | |
| | | 6.2.12 | Microbial analysis of jelly (Total plate count, yeast and mould count) | 112 | | |
| | | 6.2.13 | Statistical analysis | 112 | | |
| | 6.3 | Results | and discussions | 113 | | |
| | | 6.3.1 | Moisture content | 113 | | |
| | | 6.3.2 | Titrable acidity and pH | 113 | | |
| | | 6.3.3 | Total soluble solids (TSS) | 114 | | |
| | | 6.3.4 | Texture profile | 115 | | |
| | | 6.3.5 | Color | 115 | | |
| | | 6.3.6 | Total phenolic content | 116 | | |
| | | 6.3.7 | Total anthocyanin content | 117 | | |
| | | 6.3.8 | Antioxidant activity | 117 | | |
| | | 6.3.9 | Microbial quality of coconut water jelly | 118 | | |
| | | 6.3.10 | , i | 118 | | |
| | 6.4 | Conclu | sions | 120 120-124 | | |
| _ | | References | | | | |
| 7. | _ | Chapter 7 Summary, conclusions and future scope | | | | |
| | 7.1 | | ary and conclusions | 125 | | |
| | 72 | Future | Scope | 126 | | |