

---

## TABLE OF CONTENTS

---

<b>Contents</b>	<b>Page No.</b>
Abstract	i-iv
Keywords	v
Declaration	vi
Supervisor certificate	vii
External certificate	viii
Acknowledgements	ix - x
List of Contents	xi-xvii
List of Tables	xviii-xx
List of Figures	xxi-xxv
List of Schemes	xxvi
List of Abbreviations	xxvii-xxx
<b>Chapter 1: Introduction</b>	<b>1.1-1.81</b>
1.1 Preamble	1.1
1.2 Background information	1.1
1.2.1 Niobium (Nb)- general comments	1.1
1.2.2 A quick glance at the coordination chemistry of niobium	1.5
1.2.3 Peroxido complexes of niobium – salient features	1.8
1.2.3.1 Homoleptic and mixed-ligand/heteroleptic peroxido complexes of niobium	1.10
1.2.3.2 Applications of peroxidoniobium (pNb) compounds	1.18
1.2.3.3.1 Application as molecular precursors to obtain Nb-oxide materials	1.20
1.2.3.3.2 Application in catalysis	1.21
1.2.4 Polymer immobilized metal complexes -general aspects	1.26
1.2.4.1 Catalytic application of polymer supported metal complexes	1.29

1.2.5 Water-soluble polymers (WSPs) as metal supports – significance in catalysis and biology	1.36
1.2.5.1 Catalytic application of soluble polymer anchored metal complexes	1.40
1.2.5.2 Biological applications of soluble macro complexes	1.42
References	1.45-1.81
<b>Chapter 2: Materials and Methods</b>	<b>2.1-2.7</b>
2.1 Chemicals	2.1
2.2 Elemental analysis	2.1
2.2.1 Niobium	2.1
2.2.2 Peroxide	2.2
2.2.2.1 Permanganometry	2.2
2.2.2.2 Iodometry	2.2
2.2.2.3 By standard Ce(IV) solution	2.2
2.2.3 Carbon, Hydrogen and Nitrogen	2.2
2.2.4 Chlorine	2.3
2.2.5 Sodium	2.3
2.3 Physical and spectroscopic measurements	2.3
2.3.1 pH measurement	2.3
2.3.2 Electronic spectra	2.3
2.3.3 Infrared (IR) and Raman spectra	2.3
2.3.4 Surface morphology analysis by Scanning Electron Microscope (SEM)	2.3
2.3.5 Atomic Absorption Spectroscopy (AAS)	2.4
2.3.6 Inductively coupled plasma-optical emission spectroscopy (ICP-OES)	2.4
2.3.7 Powder X-ray diffraction (PXRD)	2.4
2.3.8 X-ray photoelectron spectroscopy (XPS)	2.4
2.3.9 Surface area analysis	2.4

2.3.10 $^1\text{H}$ NMR spectra	2.5
2.3.11 $^{13}\text{C}$ NMR spectra	2.5
2.3.12 $^{93}\text{Nb}$ NMR spectra	2.5
2.3.13 GC/GC-MS analysis	2.5
2.3.14 High performance liquid chromatography (HPLC) analysis	2.5
2.3.15 Thermogravimetric analysis	2.6
2.3.16 Melting point determination	2.6
2.3.17 Magnetic susceptibility	2.6
References	2.7
<b>Chapter 3: Niobium(V)-Peroxiido Complexes Supported on Organic Polymer as Sustainable Catalysts for Solventless Olefin Epoxidation and Sulfide Oxidation</b>	<b>3.1-3.51</b>
3.1 Introduction	3.1
3.2 Experimental section	3.3
3.2.1 Functionalization of the polymer support with amino acids	3.3
3.2.2 Preparation of sodium tetraperoxidoniobate (TpNb)	3.3
3.2.3 Synthesis of polymer immobilized triperoxidoniobium compounds (3.1-3.3)	3.3
3.2.4 General procedure for catalytic epoxidation of olefins	3.4
3.2.5 General procedure for oxidation of sulfides	3.5
3.2.6 Procedure for control experiments	3.5
3.2.7 Regeneration of the catalyst	3.6
3.3 Results and discussion	3.6
3.3.1 Synthesis	3.6
3.3.2 Characterization	3.7
3.3.2.1 SEM and EDX analysis	3.7
3.3.2.2 Powder X-ray diffraction studies	3.10
3.3.2.3 X-ray photoelectron spectroscopy	3.11

3.3.2.4 BET analysis	3.11
3.3.2.5 FTIR and Raman spectral analysis	3.12
3.3.2.6 Electronic spectral studies	3.15
3.3.2.7 <sup>13</sup> C NMR analysis	3.16
3.3.2.8 Thermogravimetric analysis	3.19
3.3.3 Catalytic performances	3.21
3.3.3.1 Epoxidation of olefins	3.21
3.3.3.2 Oxidation of sulfides	3.29
3.3.3.3 Heterogeneity test	3.31
3.3.3.4 Recyclability of the catalysts	3.34
3.3.3.5 H <sub>2</sub> O <sub>2</sub> utilization efficiency	3.35
3.3.3.6 Comparison with some reported catalysts	3.35
3.3.3.7 The proposed catalytic cycle	3.38
3.4 Conclusions	3.40
References	3.41-3.51
<b>Chapter 4: Selective and Mild Oxidation of 5-Hydroxymethylfurfural to 5-Hydroxymethyl-2-Furancarboxylic Acid over Organic Polymer-Supported Peroxido-niobium(V) Catalysts</b>	4.1-4.32
4.1 Introduction	4.1
4.2 Experimental section	4.4
4.2.1 Synthesis of [Nb(O <sub>2</sub> ) <sub>3</sub> (carboxylate)] <sup>2-</sup> -PMA (NbPMA) (4.1)	4.4
4.2.2 Procedure for the oxidation of HMF to HMFCA	4.4
4.2.3 Procedure for control experiment	4.5
4.2.4 Catalyst regeneration	4.5
4.3 Results and discussion	4.6
4.3.1 Synthesis	4.6
4.3.2 Characterization	4.6
4.3.2.1 SEM and Energy Dispersive X-ray Analysis	4.7

4.3.2.2 FTIR and Raman spectral analysis	4.8
4.3.2.3 Electronic spectral studies	4.9
4.3.2.4 <sup>13</sup> C NMR analysis	4.10
4.3.2.5 <sup>93</sup> Nb NMR analysis	4.11
4.3.2.6 Thermogravimetric Analysis	4.12
4.3.3 Catalytic performances of the polymer-supported catalysts in the conversion of HMF to HMFCFA	4.13
4.3.3.1 Effect of temperature	4.13
4.3.3.2 Effect of oxidant concentration	4.14
4.3.3.3 Effect of catalyst amount	4.16
4.3.3.4 Effect of the amount of base	4.17
4.3.3.5 Effect of reaction time	4.19
4.3.3.6 Effect of solvent	4.19
4.3.3.7 HMF oxidation over homogeneous catalyst NbPMA (4.1)	4.22
4.3.3.8 Recyclability of the catalysts	4.23
4.3.3.9 Heterogeneity test	4.24
4.4 Conclusions	4.25
References	4.26-4.32
<b>Chapter 5: Chitosan Supported Niobium(V) and Tantalum(V) Peroxido Complexes: Versatile Heterogeneous Catalysts for Eco-Compatible Organic Oxidations with H<sub>2</sub>O<sub>2</sub> under Organic-Solvent-Free Conditions</b>	5.1-5.49
5.1 Introduction	5.1
5.2 Experimental section	5.3
5.2.1 Synthesis of Na[Nb(O <sub>2</sub> ) <sub>3</sub> (NH <sub>2</sub> )(OH)]-chitosan (ChpNb) (5.1)	5.3
5.2.2 Phenol hydroxylation	5.3
5.2.3 Styrene epoxidation	5.4
5.2.4 Oxidation of thioanisole (MPS)	5.4

5.2.5	Procedure for control experiments	5.4
5.2.6	Catalyst regeneration	5.5
5.3	Results and discussion	5.5
5.3.1	Synthesis	5.5
5.3.2	Characterization	5.5
5.3.2.1	SEM and EDX Analysis	5.6
5.3.2.2	Powder X-ray diffraction analysis	5.8
5.3.2.3	X-ray photoelectron spectroscopy	5.8
5.3.2.4	BET analysis	5.9
5.3.2.5	FTIR and Raman spectral analysis	5.10
5.3.2.6	Electronic spectral analysis	5.12
5.3.2.7	<sup>13</sup> C NMR analysis	5.12
5.3.2.8	Thermogravimetric analysis	5.13
5.3.3	Catalytic activity	5.15
5.3.3.1	Phenol hydroxylation	5.15
5.3.3.2	Styrene epoxidation	5.21
5.3.3.3	Sulfide oxidation	5.26
5.3.3.4	Stability and reusability of the catalyst	5.28
5.3.3.5	Heterogeneity test	5.31
5.3.3.6	Scalability of the catalytic methodologies	5.32
5.3.3.7	Comparison with some reported catalysts	5.32
5.3.3.8	Proposed catalytic cycle	5.35
5.4	Conclusions	5.36
	References	5.38-5.49
	<b>Chapter 6: <i>In vitro</i> Acid Phosphatase Inhibition by Oxido- and Mixed-Ligand Peroxido-Niobium(V) Complexes</b>	6.1-6.41
6.1	Introduction	6.1
6.2	Experimental section	6.3
6.2.1	Synthesis of the complex, [NbO(malt) <sub>3</sub> ] <sub>2</sub> ·9H <sub>2</sub> O (6.1)	6.3

6.2.2 Synthesis of the complex $\text{Na}_2[\text{Nb}(\text{O}_2)_3(\text{malt})]\cdot\text{H}_2\text{O}$ (6.2)	6.3
6.2.3 Synthesis of the complex $\text{Na}_2[\text{Nb}(\text{O}_2)_3(\text{def})]\cdot 2\text{H}_2\text{O}$ (6.3)	6.4
6.2.4 X-Ray Crystallography	6.4
6.2.5 Computational Details	6.5
6.2.6 Stability of the pNb complexes in aqueous solution	6.5
6.2.7 Acid phosphatase activity measurement	6.6
6.2.8 Determination of kinetic parameters	6.6
6.3 Results and discussion	6.7
6.3.1 Synthesis	6.7
6.3.2 Characterization	6.7
6.3.2.1 FTIR and Raman spectral analysis	6.8
6.3.2.2 Electronic spectral analysis	6.11
6.3.2.3 $^1\text{H}$ and $^{13}\text{C}$ NMR analysis	6.12
6.3.2.4 Thermogravimetric analysis	6.14
6.3.3 Description of the crystal structures	6.19
6.3.4 Theoretical Investigation	6.22
6.3.5 Stability of the compounds in aqueous solution	6.23
6.3.6 Nb(V) compounds as phosphatase inhibitors	6.25
6.3.7 Kinetics of ACP inhibition by the Nb(V) complexes	6.26
6.4 Conclusions	6.30
References	6.31-6.41
Appendix	xxxix-xxxvii
List of Publications	xxxviii
List of Conferences Attended	xl