

Dedicated to

Maa and Deuta

For dust you are and to dust you will return

-- Genesis 3:19

Declaration by the candidate

I, **OLAG PRATIM BORDOLOI**, declare that the thesis entitled “**A multi-wavelength study of the interstellar medium of dwarf irregular galaxies**” is a record of my own work carried out under the supervision of **Dr. Rupjyoti Gogoi** during the period from **July, 2018** to **January, 2024**. I confirm that:

- This work was done wholly or mainly while in candidature for a research degree in Doctor of Philosophy at Tezpur University. Where no part of this thesis has previously been submitted for a degree or any other qualification at this University or any other institution.
- Where I have quoted from the work of others, the source is always given. With the exception of such quotations, this thesis is entirely my own work. I have acknowledged all main sources of help.
- Whereas the thesis is based on the work done by myself jointly with others, I have made clear exactly what was done by others and what I have contributed myself.

Date: 25-07-2024

Place: Tezpur University

Olag Pratim Bordoloi

(Olag Pratim Bordoloi)

Regd. No.: TZ201052 of 2019



TEZPUR UNIVERSITY

Certificate from the Supervisor

This is to certify that the thesis entitled “**A multi-wavelength study of the interstellar medium of dwarf irregular galaxies**”, submitted to the School of Sciences, Tezpur University in partial fulfillment of the requirements for the award of the degree of **Doctor of Philosophy in Physics** is a record of original research work carried out by **Mr. Olag Pratim Bordoloi** under my supervision and guidance.

All help received by him from various sources have been duly acknowledged. No part of this thesis has been submitted elsewhere for award of any other degree.

Rupjyoti Gogoi

(Dr. Rupjyoti Gogoi)

Supervisor, Assistant Professor

Department of Physics

School of Sciences

Tezpur University

Acknowledgements

"Kind words can be short and easy to speak, but their echoes are truly endless"

- Mother Teresa

The completion of this thesis marks a remarkable journey, made possible by the unwavering support and motivation from numerous individuals. I extend my heartfelt thanks to each and every one of them for being a constant presence in my moments of need.

First and foremost, my deepest gratitude goes to my supervisor and mentor, Dr. Rupjyoti Gogoi. Throughout the years of our association, she has consistently demonstrated unwavering faith in my abilities and provided a continuous source of motivation. She has not only encouraged excellence in research but also in life. Her accessibility, whether through email, instant messaging or phone, has been invaluable, offering guidance whenever doubt arose. I consider myself fortunate to have had the opportunity to work under her guidance. She is not just an exceptional Ph.D. advisor but also a friendly and understanding person. The values she has imparted have played a pivotal role in my evolution, both as a researcher and as a human being during the course of this Ph.D.

The contributions of my collaborators – Dr. P. Shalima, Dr. Margarita Safonova, and Prof. Yuri A. Shchekinov have been indispensable to the realization of this thesis. Dr. P. Shalima, in particular, has been my go-to person whenever challenges arose, and her advice has consistently proven invaluable. Her humility, despite possessing immense knowledge, is truly commendable. I am forever grateful for her motivation and guidance throughout this thesis work.

I express my gratitude to the members of my doctoral committee, Prof. Nilakshi Das, and Dr. Moon Moon Devi, for their insightful comments and encouragement, providing the impetus to broaden my research from various perspectives.

Tezpur University, my home for more than five years, and the Department of Physics have provided an excellent environment for study. I am thankful to Tezpur University for the provision of internet and computational facilities, laboratory resources, and infrastructure,

all of which facilitated the swift completion of this research. Special thanks to all faculty members, all the heads of the department during my tenure – Prof. Nilakshi Das, Prof. Mrinal Kr. Das, Prof. Nidhi Saxena Bhattacharya and Prof. Pabitra Nath, for their consistent support, and to Mr. Narayan Sarma for his continuous assistance on all official matters. I express my gratitude for the financial support received during my Ph.D. tenure – thanks to the Science and Engineering Research Board (SERB) - Department of Science and Technology (DST), Govt. of India for the JRF fellowship and Tezpur University for the institutional fellowship.

Heartfelt thanks to my colleagues from the Astronomy and Astrophysics Lab and friends from Tezpur University and beyond. Your companionship has made my time as a Ph.D. student joyous and memorable. So, thank you Rituraj, Bhugendra Daa, Plaban, Gautam, Pranjupriya, Rukaiya, Anshuman, Hritwik, Swagat, Bichitra Daa, Ankur, Amar, Anup, Sritam, Diganta, Ankush, Prankrishna, Tonmoy, Jotin, Abinash, Bidhan, Biprav, Jubaraj, Niranjana, Debasish, Rupam, Anil, Dhanjit, Raju Daa, Parag, Debajit, Bidisha, Sunita, Snigdha, and all whom I have interacted with. I gratefully acknowledge the help received specially from Anshuman and Hritwik in different stages of this research. In addition to my friends, heartfelt gratitude goes to my family members for their unwavering support. Special mention to Riya (Mainee), whose constant companionship has been a pillar of strength throughout this phase of my life. Without her untiring love and support, facing the challenges alone would have been considerably more difficult.

Finally, I want to express my acknowledgment to all the telescope facilities, online astronomical resources, and various other sources that have played an indispensable role in making this thesis work possible. This work uses the data from the AstroSat mission of the Indian Space Research Organisation (ISRO), archived at the Indian Space Science Data Centre (ISSDC). This research has used observations made with the Spitzer Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. Herschel is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This research has made use of the NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

Thank you all for your unwavering and continuous support!

Olag Pratim Bordoloi
Olag Pratim Bordoloi

List of Publications

Published in refereed journals

1. A. Vinokurov, K. Atapin, **O. P. Bordoloi**, A. Sarkisyan, U. Kashyap, M. Chakraborty, P.T. Rahna, A. Kostenkov, Y. Solovyeva, S. Fabrika, M. Safonova, R. Gogoi, F. Sutaria and J. Murthy. Simultaneous X-ray/UV Observations of Ultraluminous X-ray Source Holmberg II X-1 with Indian Space Mission AstroSat, *Astrophysical Bulletin*, vol. 77, no. 3, pp. 231-245, 2022.
2. **Olag Pratim Bordoloi**, B. Ananthamoorthy, P. Shalima, Margarita Safonova, Debbijoy Bhattacharya, Yuri A. Shchekinov and Rupjyoti Gogoi. AstroSat/UVIT Study of the Diffuse Ultraviolet Radiation in the Dwarf Galaxy Holmberg II, *Publications of the Astronomical Society of the Pacific*, vol. 136, no. 7, p. 074101, 2024.

To be submitted

1. **Olag Pratim Bordoloi**, Yuri A. Shchekinov, P. Shalima, M. Safonova and Rupjyoti Gogoi. Unravelling multi-temperature dust populations in the dwarf galaxy Holmberg II.

Conference proceedings

1. K. Atapin, A. Vinokurov, M. Safonova, S. Fabrika, **O. Bordoloi**. Long term X-ray/UV observations of the ultraluminous X-ray source Holmberg II X-1, *Proceedings of Science*, vol. 425, p. 56, 2022.
2. Barenya Kumar Dev, Gautam Saikia, **Olag Pratim Bordoloi**, P. Shalima and Rupjyoti Gogoi. Analysis of Diffuse Far Ultraviolet emissions near R136 cluster in 30 Doradus, *under review in Journal of Physics: Conference Series*.

Book chapters

1. **Olag Pratim Bordoloi** and Rupjyoti Gogoi. Interstellar Dust: Unveiling Cosmic Grains and Their Enigmatic Journey, *Recent Review and Research in Physics (Volume - 6)*, AkiNik Publications, New Delhi, India, ISBN: 978-93-5570-878-6, 2023.
2. **Olag Pratim Bordoloi**, Anshuman Borgohain and Rupjyoti Gogoi. Dwarf Galaxies: Our Bridge to the Early Universe, *Recent Review and Research in Physics (Volume - 8)*, AkiNik Publications, New Delhi, India, ISBN: 978-93-6135-325-3, 2024.

List of figures

1.1	An schematic diagram of the life cycle of interstellar dust grains (Image credit: <i>Herschel Space Observatory</i>)	10
1.2	The Core-Mantle structure of interstellar dust grains (Image credit: https://astronomy.swin.edu.au/cosmos/D/Dust+Grain)	12
1.3	Typical Milky Way extinction curves along different lines of sight, characterized by different values of R_V (adopted from [11])	13
2.1	The different payloads onboard <i>AstroSat</i> (Image credit: ISRO)	27
2.2	An schematic diagram showing the different processes involved in the processing of UVIT data with JUDE	31
2.3	UVIT - FUV co-added image of Ho II (Observation date: December 9, 2016, Filter: F154W)	32
2.4	UVIT - NUV co-added image of Ho II (Observation date: December 9, 2016, Filter: N245M)	33
2.5	(a) <i>AstroSat</i> /UVIT image of the star forming region harboring the ULX Ho II X-1, (b) <i>GALEX</i> image of the same region. The location of the ULX is shown with the reticle. The two images clearly highlight the significant improvement of the resolution of UVIT compared to <i>GALEX</i>	34
3.1	Light curve of synchronous X-ray–UV observations of Ho II X-1. The X-ray fluxes (0.7–7 keV) are marked by black points with dashed bars; the UV flux densities at 148.1 nm (F148W filter), are shown in green	39
3.2	X-ray fluxes against UV flux densities at 148.1 nm (F148W filter). Colors are in the rainbow order from red (observation #1) to violet (#10). Formal Pearson correlation coefficient is $R = 0.45$	40

3.3	The ratio of the flux densities at 1500 Å from the heated and not heated donors (<i>left panels</i>), and the temperatures averaged over the visible star disk (<i>right panels</i>) as functions of the X-ray luminosity of the accretor. Two type of the accretors: a neutron star (<i>top</i>) and a black hole (<i>bottom</i>) are considered. The donor is either a O8 or B2 supergiant. The orbit inclination either $i = 0$ (view along the axis) or the maximum possible angle at which the particular system remains non-eclipsing in X-rays. Grey areas show the range of Ho II X-1 X-ray luminosities in the <i>AstroSat</i> observations	42
4.1	FUV–IR correlation plots for 17 selected locations with $N(\text{HI}) > 1 \times 10^{21} \text{ cm}^{-2}$. The straight lines represent linear fits	53
4.2	NUV–IR correlation plots for 17 selected locations with $N(\text{HI}) > 1 \times 10^{21} \text{ cm}^{-2}$. Straight lines represent linear fits	54
4.3	FUV–IR and NUV–IR intensity plots for 16 locations with $N(\text{HI}) < 1 \times 10^{21} \text{ cm}^{-2}$	57
4.4	<i>Left</i> : Smoothed FUV image of Ho II with isophotes overplotted for five intensity levels. <i>Right</i> : Multiband image of Ho II with FUV (F154W; <i>AstroSat</i> -UVIT) in blue, IR (3.6 μm; <i>Spitzer</i>) in green, and integrated HI map (THINGS) in teal contours. The HI contour corresponds to $N(\text{HI}) = 1 \times 10^{21} \text{ cm}^{-2}$. Selected diffuse UV locations with $N(\text{HI}) > 1 \times 10^{21} \text{ cm}^{-2}$ are indicated by purple circles, and locations with $N(\text{HI}) < 1 \times 10^{21} \text{ cm}^{-2}$ are indicated by white squares. It is evident from the figure that the locations with $N(\text{HI}) < 1 \times 10^{21} \text{ cm}^{-2}$ mostly lie in the cavities devoid of HI gas . . .	58
4.5	Schematic diagram showing the relative geometry of the source (star) and the scattering of starlight by optically thin clouds towards the observer . . .	61
4.6	Locations depicted in Table 4.8 overplotted on the FUV image of Ho II obtained by the UVIT instrument [114]. (a) Locations where intensity peaks at 100 μm, represented by cyan colour, mostly lie in the SFR, (b) Locations where intensity peaks at 70 μm, represented by green colour, lie mostly in the outskirts of SFR, (c) Locations lying in HI voids ($N(\text{HI}) < 1 \times 10^{21} \text{ cm}^{-2}$), represented by red colour. The circles representing the locations are of the same size as circular apertures (5'' radius) used to extract the IR intensities from the locations. The FUV image was taken on December 9, 2016 with an exposure time of around 9 ks	66

4.7	Fits of the spectra of dust at different environmental conditions as classified in Fig. 4.6. They represent spectra of dust: i) spectra peaking at $100 \mu\text{m}$, ii) spectra peaking at $70 \mu\text{m}$, iii) in regions with HI column densities below the detection limit $N(\text{HI}) < 10^{21} \text{ cm}^{-2}$, iv) spectra with emission at $8 \mu\text{m}$, v) spectra with a weak intensity dip at $100 \mu\text{m}$ and vi) spectra with a strong intensity dip at $100 \mu\text{m}$. Specifically, the presented spectra fitted with multi-temperature modified black body with the following temperatures and dust surface mass densities for cold components and dust masses for warm and hot ones: (a) $T = 27, 56, 310 \text{ K}$ ($\Sigma_d = 6 \times 10^{-5} M_\odot \text{pc}^{-2}$, $M_d = 0.009, 6.4 \times 10^{-8} M_\odot$); (b) $T = 10, 25, 53, 220, 320 \text{ K}$ ($\Sigma_d = 3 \times 10^{-3} M_\odot \text{pc}^{-2}$, $M_d = 0.5, 0.08, 4 \times 10^{-7}, 4 \times 10^{-8} M_\odot$); (c) $T = 15, 46, 260 \text{ K}$ ($\Sigma_d = 7 \times 10^{-4} M_\odot \text{pc}^{-2}$, $M_d = 0.025, 3 \times 10^{-7} M_\odot$); (d) $T = 28, 63, 100, 320 \text{ K}$ ($\Sigma_d = 6 \times 10^{-4} M_\odot \text{pc}^{-2}$, $M_d = 63, 0.002, 2 \times 10^{-7} M_\odot$)	68
4.8	Variation of temperature of cold dust with FUV intensity for the considered 27 locations. Colour coding is according to Fig. 4.6	70
5.1	<i>FUSE</i> observed dust locations in FUV (blue squares) and the star locations (red stars) overplotted on DSS2 color image of 30 Doradus at field of view of $1.2^\circ \times 36'$	78
5.2	The scattering optical depth (τ_1) (1157 \AA) at the 30 Doradus <i>FUSE</i> locations is represented by circles. The area of the circles is proportional to τ_1 . These circles are overlaid on a DSS2 IR band image	83

List of tables

2.1	Some general properties of Ho II	26
2.2	Details of <i>AstroSat</i> /UVIT observations of Ho II	28
2.3	Properties of different UVIT filters used in our observation	28
2.4	JUDE modules and their purposes	31
3.1	Observation dates, exposures and measured source fluxes. The X-ray fluxes (0.7-7.0 keV) are in units of 10^{-12} erg s ⁻¹ cm ⁻² . The UV flux densities (background subtracted) are in units of 10^{-16} erg s ⁻¹ cm ⁻² Å ⁻¹	38
4.1	Diffuse FUV, NUV intensities, and neutral hydrogen column densities N(HI) for selected 33 locations having non-zero IR intensities. The N(HI) values have been derived from the THINGS integrated HI map. The UV intensities are in units of photons cm ⁻² s ⁻¹ sr ⁻¹ Å ⁻¹ , and N(HI) values are in units of 10^{21} cm ⁻² . Magenta colour indicates locations with $N(\text{HI}) < 1 \times 10^{21}$ cm ⁻²	50
4.2	IR intensities in selected 33 locations. See Sec. 4.3.1 for details	51
4.3	Best-fit ratios of IR and UV intensities in units of 10^{-5} MJy/(phot/cm ² /s/Å) for locations with $N(\text{HI}) > 1 \times 10^{21}$ cm ⁻²	54
4.4	FUV-IR and NUV-IR correlation values for locations with $N(\text{HI}) > 1 \times 10^{21}$ cm ⁻² (Figs. 4.1 and 4.2)	55
4.5	FUV-IR and NUV-IR correlation values for locations with $N(\text{HI}) < 1 \times 10^{21}$ cm ⁻² (Fig. 4.3)	55
4.6	Properties of the contributing star-forming complexes. The complexes have been adopted from [108]	60
4.7	The values of the best-fit parameters (based on minimum χ^2 -statistic) for locations with $N(\text{HI}) > 1 \times 10^{21}$ cm ⁻² . The range of values within 90% confidence level for each parameter are shown inside parentheses. The location numbers are in tally with Table 4.1	62

4.8	IR intensities in selected 27 locations. Colour coding is according to Fig. 4.6. See Sec. 4.3.1 for details of calculations	65
5.1	Total hydrogen column density $N(\text{H})$ map of 30 Doradus and SN1987A. $N(\text{H})$ is in the units of H atoms per cm^2 [Credits: AIfA Hi Surveys Data Server of the Argelander-Institut für Astronomie and HEASARC $N(\text{H})$ tool]	76
5.2	List of the 22 sources under consideration and their luminosities corresponding to the 7 <i>FUSE</i> bands (in $\text{ergs s}^{-1} \text{\AA}^{-1}$)	79
5.3	<i>FUSE</i> observed dust scattered intensities (in $\text{photons cm}^{-2} \text{s}^{-1} \text{sr}^{-1} \text{\AA}^{-1}$) [96]	80
5.4	<i>FUSE</i> wavelengths and corresponding albedo (α), asymmetry factor (g), extinction cross section (σ_{ext} (in cm^2 / H) [35] and calculated mean optical depth (τ_{avg})	80
5.5	Single scattering model outputs for the <i>FUSE</i> bands (Continued in Table 5.6)	81
5.6	Single scattering model outputs for the <i>FUSE</i> bands (Contd.)	82
5.7	Percentage contribution of RMC 136 to the surface brightness of the diffuse FUV emission at dust locations at 1157\AA	82
5.8	Scattering optical depth (τ_1) of each dust location corresponding to the <i>FUSE</i> wavelengths (μm^{-1})	83

Nomenclature

Acronyms / Abbreviations

AGB Asymptotic Giant Branch

BCD Blue Compact Dwarf

CMB Cosmic Microwave Background

CNM Cold Neutral Medium

CZTI Cadmium Zinc Telluride Imager

dIrr Dwarf Irregular

dSph Dwarf Spheroidal

FIGGS Faint Irregular Galaxies GMRT Survey

FIR Far Infrared

FUSE Far Ultraviolet Spectroscopic Explorer

FUV Far Ultraviolet

FWHM Full Width at Half Maximum

GDL GNU Data Language

HIFI Heterodyne Instrument for the Far Infrared

HIM Hot Ionized Medium

Ho II Holmberg II

HST Hubble Space Telescope

IRAC Infrared Array Camera

IR Infrared

IRS Infrared Spectrograph

ISM Interstellar Medium

JUDE Jayant's UVIT Data Explorer

JWST James Webb Space Telescope

LAXPC Large Area X-ray Proportional Counter

LEGUS Legacy Extragalactic UV Survey

LMC Large Magellanic Cloud

MIPS Multiband Imaging Photometer for Spitzer

NIR Near Infrared

NUV Near Ultraviolet

PACS Photoconductor Array Camera and Spectrometer

PAH Polycyclic Aromatic Hydrocarbon

PSF Point Spread Function

SDSS Sloan Digital Sky Survey

SED Spectral Energy Distribution

SFR Star Formation Rate

SHIELD Survey of HI in Extremely Low Mass Dwarfs

SMC Small Magellanic Cloud

SNe Supernovae

SPIRE Spectral and Photometric Imaging Receiver

SSM Scanning Sky Monitor

SXT Soft X-ray Telescope

THINGS The HI Nearby Galaxy Survey

UIT Ultraviolet Imaging Telescope

ULXP Ultra luminous X-ray Pulsars

ULX Ultra luminous X-ray

UVIT Ultra Violet Imaging Telescope

UV Ultraviolet

VSG Very Small Grains

WIM Warm Ionized Medium

WNM Warm Neutral Medium

XMM X-ray Multi-Mirror Mission

