

Table of contents	Page no.
Abstract	i-ii
Declaration	iii
Certificate	iv
Acknowledgments	v-vi
Contents	vii-xiii
List of tables	xiv-xv
List of Figures	xvi-xvii
List of abbreviations	xviii-xx
List of symbols	xxi

Chapter 1: Introduction	Page no.
1.1 Overall introduction and literature review	1.1-1.3
1.2 Economic importance of pineapple	1.3-1.4
1.3 Impact of Pineapple Waste Utilization	1.4-1.5
1.4 Physico-chemical composition and nutritional value of pineapple	1.6-1.8
1.5 Pineapple waste towards a circular economy	1.9
1.6 Pineapple waste-processing and potential value-added products	1.10-1.13
1.7 Pineapple waste-based products	
1.7.1 Pineapple food-based products	1.14-1.16
1.7.2 Volatile compounds	1.17-1.18
1.7.3 Dietary fiber (soluble & insoluble)	1.19-1.20
1.7.4 Bioactive compounds	1.21
1.7.5 Organic acid	
1.7.5.1 Citric acid	1.22-1.23
1.7.5.2 Lactic acid	1.24
1.7.5.3 Ferulic acid	1.24
1.7.6 Enzyme	1.25
1.7.7 Phenolic compounds	1.25-1.26
1.7.8 Residual sugar recovery	1.27
1.7.9 Hydrogels and cellulose nanofiber production	1.28-1.30
1.7.10 Methane production	1.30

1.8	Health benefits	1.32-1.33
1.9	Dietary fiber as a functional component	1.34
1.10	Extraction of dietary fiber	1.34
1.11	Modification of dietary fiber	1.35
1.12	Antidiabetic study	1.36
1.13	Development of functional food	1.37
1.14	Hypothesis	1.38
1.15	Gap of research	1.38
1.16	Objectives	1.38
	Bibliography	1.40-1.49

Chapter 2: To study different extraction methods on dietary fiber from queen pineapple waste of Northeast India and its process optimization **Page no.**

2.1	Introduction	2.1-2.2
2.2	Materials and Methods	
2.2.1	Raw materials and chemicals	2.3
2.2.2	Extraction of dietary fiber	2.3
2.2.3	Experimental design	2.4
2.2.4	Chemical compositional analysis of extracted dietary fibre	2.5
2.2.5	Mineral Content	2.5
2.2.6	FTIR analysis of extracted dietary fibre	2.5
2.2.7	SEM analysis of extracted dietary fibre	2.5
2.2.8	XRD analysis of extracted dietary fibre	2.5
2.2.9	Thermo-gravimetry analysis (TGA) of extracted dietary fibre	2.5
2.2.10	Functional properties of the extracted dietary fibre	
2.2.10.1	Water holding capacity (WHC) and oil holding capacity (OHC)	2.6
2.2.10.2	Swelling capacity (SC)	2.6
2.2.10.3	Cation exchange capacity (CEC)	2.6

2.2.10.4	Emulsifying activity (EA) and emulsion stability (ES)	2.7
2.2.11	Hypoglycemic activity	
2.2.11.1	Glucose adsorption capacity (GAC)	2.7
2.2.11.2	Glucose dialysis retardation index (GDRI)	2.7
2.2.12	Statistical analysis	2.8
2.3	Results and Discussion	
2.3.1	Optimization of the UAE method	2.8-2.9
2.3.2	Determination of significance factors	2.10
2.3.3	Optimum conditions for extraction	2.11
2.3.4	Chemical composition of pineapple waste and extracted dietary fiber	2.12
2.3.5	FTIR analysis of extracted dietary fibre	2.13
2.3.6	SEM analysis of the extracted dietary fibre	2.15-2.16
2.3.7	XRD analysis of the extracted dietary fibre	2.17
2.3.8	Thermo-gravimetric analysis (TGA) of extracted dietary fibre	2.18-2.19
2.3.9	Functional properties of extracted dietary fibre	2.19-2.21
2.3.10	Glucose adsorption capacity	2.21
2.3.11	Determination of glucose dialysis retardation index	2.22
2.4	Conclusion	2.23
	Bibliography	2.24-2.28

Chapter 3: To evaluate the effect of enzymatic modification of extracted dietary fiber and its characterization **Page no.**

3.1	Introduction	3.1-3.2
3.2	Methodology	
3.2.1	Materials	3.3
3.2.2	Ultrasonic-assisted extraction of IDF	3.3
3.2.3	Modification of IDF	3.3
3.2.4	FTIR analysis of modified dietary fibre	3.4
3.2.5	SEM analysis of modified dietary fibre	3.4
3.2.6	XRD analysis of modified dietary fibre	3.4

3.2.7	Thermo-gravimetry analysis (TGA) of modified dietary fibre	3.4
3.2.8	Functional characterization of the enzyme-modified dietary fibre	
3.2.8.1	Water holding capacity (WHC) and oil holding capacity (OHC)	3.4
3.2.8.2	Swelling capacity (SC)	3.5
3.2.8.3	Emulsifying activity (EA) and Emulsion stability (ES)	3.5
3.2.8.4	Cation exchange capacity (CEC)	3.5
3.2.8.5	Cholesterol absorption capacity (CAC)	3.6
3.2.9	Hypoglycemic activity	
3.2.9.1	Glucose adsorption capacity (GAC)	3.6
3.2.9.2	Glucose dialysis retardation index (GDRI)	3.7
3.2.10	Statistical analysis	3.7
3.3	Result and discussion	3.7
3.3.1	FTIR of modified dietary fibre	3.8-3.9
3.3.2	SEM of modified dietary fibre	3.10-3.11
3.3.3	XRD of modified dietary fibre	3.11
3.3.4	TGA of modified dietary fibre	3.12
3.3.5	Functional properties	3.14-3.15
3.3.6	Hypoglycaemic activity:	
3.3.6.1	GAC of modified dietary fibre	3.16
3.3.6.2	GDRI of modified dietary fibre	3.16
3.4	Conclusion	3.17
	Bibliography	3.19-3.22

Chapter 4: To study the <i>in vitro</i> antidiabetic activity of modified dietary fiber		Page no.
4.1	Introduction	4.1-4.2
4.2	Materials and methods	
4.2.1	Sample preparation	4.3
4.2.2	Extraction and Modification of queen pineapple waste	4.3
4.2.3	Anti-diabetic study of enzyme-modified dietary fiber from queen pineapple waste	
4.2.3.1	α -amylase inhibition activity:	4.4
4.2.3.2	α -glucosidase inhibition activity	4.4
4.2.3.3	DPP IV Inhibition	4.5
4.2.4	Cell culture	4.5
4.2.5	Cytotoxicity assessment by MTT assay in Caco2 and HepG2 cell line	4.6
4.2.6	Glucose (2-NBDG) uptake assay in Caco2 and HepG2 cell line	4.6
4.2.7	RNA extraction of Caco2 and HepG2 cell line	4.7
4.2.8	RNAseq dataset and analysis	4.7
4.2.9	Statistical analysis	4.8
4.3	Results and discussion	
4.3.1	Anti-diabetic study of enzyme-modified dietary fiber	
4.3.1.1	α -amylase inhibition	4.8
4.3.1.2	α -Glucosidase inhibition	4.9
4.3.1.3	DPP IV Inhibition	4.10
4.3.2	Toxicity study of enzyme-modified dietary fiber from queen pineapple waste	
4.3.2.1	Cell viability assay of enzyme-modified dietary fiber in Caco2 and HepG2 cell line	4.11-4.12
4.3.3	<i>In vitro</i> glucose uptake assay of enzyme-modified dietary fiber	4.13-4.14
4.3.4	Effect of EMDF on RNA sequencing and differentially expressed mRNAs	4.15

4.3.5	Functional prediction analysis of RNAseq by GO enrichment	4.16
4.4	Conclusion	4.17
	Bibliography	4.18-4.20

Chapter 5: To develop a fiber-rich functional food and its quality evaluation **Page no.**

5.1	Introduction	5.1-5.3
5.2	Methodology	
5.2.1	Materials	5.4
5.2.2	Preparation of functional bread	5.4
5.2.3	Chemical constituents of functional bread	
5.2.3.1	Water retention capacity of functional bread	5.5
5.2.3.2	Specific volume of functional bread	5.5
5.2.4	Color Characteristics of Functional Bread	5.5
5.2.5	Texture profile analysis of functional bread	5.5
5.2.6	Sensory evaluation	5.6
5.2.7	<i>In vitro</i> analysis of functional bread	
5.2.7.1	<i>In vitro</i> starch digestibility of functional bread	5.6
5.2.7.2	Approach to Kinetic modeling	5.7
5.2.8	Statistical analysis	5.8
5.3	Result and discussion	
5.3.1	Chemical composition of Functional bread	5.9
5.3.2	Physical properties of Functional bread	5.10
5.3.2.1	Water retention capacity and specific volume of functional bread	5.11
5.3.2.2	Color characteristics of functional bread	5.12
5.3.2.3	Textural properties of functional bread	5.13-5.14
5.3.3	Sensory evaluation of functional bread	5.15
5.3.4	<i>In vitro</i> analysis of functional bread	
5.3.4.1	<i>In vitro</i> starch digestibility	5.16-5.17
5.3.4.2	Predicted glycemic index	5.17

5.4	Conclusion	5.18
	Bibliography	5.19-5.22

Chapter 6: Conclusions and future scope of work		Page no.
6.1	Conclusions	6.1-6.2
6.2	Future scope of the present investigation	6.3
	List of publications	6.4-6.5
	Participation in National/International conferences	6.5-6.6