## **CONTENTS**

CHAPTERS	PAGE NO.
1. INTRODUCTION	1 - 5
2. REVIEW OF LITERATURE	6 - 12
3. MATERIALS AND METHODS	13 - 22
4. RESULTS AND DISCUSSIONS	23 - 50
5. SUMMARY AND CONCLUSION	51
6. BIBLIOGRAPHY	52 - 66

## LIST OF FIGURES

FIGURE NO.	. TITLE	PAGE NO.
3.1	Taro tubers locally known as "Panchamukhi Kochu".	13
3.2	Flowchart showing the extraction process of Colocasia starch	14
4.1	% Ash content of native and different types of HMT modified	i
	Taro starches.	24
4.2	% Crude fat content of native and different types of HMT	
	modified Taro starches.	24
4.3	% Crude Protein content of native and different types of	
	HMT modified Taro starches.	25
4.4	% Crude fibre content of native and different types of	
,	HMT modified Taro starches.	25
4.5	% Starch content of native and different types of HMT	
	modified Taro starches.	26
4.6	% Amylose content of native and different types of HMT	
	modified Taro starches.	28
4.7	X-ray diffraction pattern of native and HMT modified	
	C. esculenta starch granules.	29
4.8	FT-IR profile of native and different types of HMT	

	modified Taro starch.	30
4.9	Swelling power of native and different types of HMT	
	modified Taro starches.	33
4.10	Solubility profile of native and different types of HMT	
	modified Taro starches.	34
4.11:	Paste Clarity of native and different types of HMT modified	
	Taro starches with respect to time.	37
4.12	Variation in Freeze-Thaw stability of Taro starch with respect	
	to time.	39
4.13	The RVA viscoamylogram of native and different types of	
	HMT modified Taro.	41
4.14	Scanning electron micrograph (4,000X) of C. esculenta	
•	(native starch) starch granules showing granule size and	
	morphology.	42
4.15	Scanning electron micrograph (4,000X) of C. esculenta	
	(Microwave HMT modified starch) starch granules showing	
	granule size and morphology.	43
4.16	Scanning electron micrograph (4,000X) of C. esculenta	
	(Hot Air Oven HMT modified starch) starch granules showing	
	granule size and morphology.	44
4.17	Scanning electron micrograph (4,000X) of C. esculenta	
	(Autoclave HMT modified starch) starch granules showing	
	granule size and morphology.	44
4.18	Scanning electron micrograph (4,000X) of C. esculenta	
	(Microwave + Hot Air Oven HMT modified starch) starch	
·	granules showing granule size and morphology.	45
4.19	Scanning electron micrograph (4,000X) of C. esculenta	
	(Microwave + Autoclave HMT modified starch) starch	
	granules showing granule size and morphology.	45
4.20	Firmness profile of native and different types of HMT modified	
	Taro starch gel.	48

4.21	Consistency profile of native and different types of HMT	
	modified Taro starch gel.	48
4.22	Cohesiveness profile of native and different types of HMT	
	modified Taro starch gel.	49
4.23	Index of viscosity profile of native and different types of	
	HMT modified Taro starch gel.	49
·		

## LIST OF TABLES

TTTLE	PAGE NO.
Proximate composition of native and different types of HMT	
modified Taro starches.	23
Effect of heat moisture treatment on % amylose content of Taro	
starch.	27
Colour of native and different types of HMT modified Taro	
starches.	31
Effect of temperature on swelling power of Taro starch.	32
Effect of temperature on % solubility of Taro starch.	33
Effect of time on paste clarity of Taro starch after heating at 95 °C	. 36
Variation in Freeze-Thaw stability of Taro starch with respect to	
time.	38
Pasting properties of native and different types of HMT modified	
Taro starches.	40
Gel texture of native and different types of HMT modified	
Taro starches.	47
	67
•	68
	Proximate composition of native and different types of HMT modified Taro starches.  Effect of heat moisture treatment on % amylose content of Taro starch.  Colour of native and different types of HMT modified Taro starches.  Effect of temperature on swelling power of Taro starch.  Effect of temperature on % solubility of Taro starch.  Effect of time on paste clarity of Taro starch after heating at 95 °C. Variation in Freeze-Thaw stability of Taro starch with respect to time.  Pasting properties of native and different types of HMT modified Taro starches.  Gel texture of native and different types of HMT modified