TABLE OF CONTENTS

Page no

Chapter 1: Introduction	1
Chapter 2: Review of Literature	5
2.1 Consumer's perception on minimally processed vegetables	6
2.2 Use of calcium chloride in minimal processing	7
2.3 Use of sodium hypochlorite in minimal processing	11
2.4 Microbiology and nutritional quality of minimally processed vegetables	13
Chapter 3: Materials and Methods	18
3.1 Raw materials	18
3.2 Preparation of solution	18
3.3 Method of minimal processing	18
3.4 Analysis of minimally processed vegetable	19
Chapter 4: Results and Discussion	22
4.1 Effect on colour and texture	22
4.2 Effect on moisture content	24
4.3 Effect on protein	34
4.4 Effect on ash content	34
4.5 Effect on crude fibre	34
4.6 Effect on ascorbic acid	34
4.7 Effect on water activity	42
4.8 Effect on microbial load	44
Chapter 5: Summary and Conclusion	52
Chapter 6: Future scope of work	54
References	55

÷

LIST OF TABLES

y.

Table. no

.

ż .

Table

4.2.1	Effect of cutting knife and sodium hypochlorite dip solution in moisture content (%) of minimally processed vegetables stored in different packaging material	25
4.2.2	Effect of cutting knife and sodium hypochlorite-calcium chloride dip solution in moisture content (%) of minimally processed vegetables stored in different packaging material	25
4.3.1	Effect of cutting knife and sodium hypochlorite dip solution in protein (%) content of minimally processed vegetables stored in different packaging material.	31
4.3.2	Effect of cutting knife and sodium hypochlorite-calcium chloride dip solution in protein (%) content of minimally processed vegetables stored in different packaging material.	31
4.4.1	Effect of cutting knife and sodium hypochlorite dip solution in ash (%) content of minimally processed vegetables stored in different packaging material.	32
4.4.2	Effect of cutting knife and sodium hypochlorite-calcium chloride dip solution in ash (%) content of minimally processed vegetables stored in different packaging material.	32
4.5.1	Effect of cutting knife and sodium hypochlorite dip solution in crude fibre content (g) of minimally processed vegetables stored in different packaging material	33
4.5.2	Effect of cutting knife and sodium hypochlorite-calcium chloride dip solution in crude fibre content (g) of minimally processed vegetables stored in different packaging material	33
4.6.1	Effect of cutting knife and sodium hypochlorite dip solution in ascorbic acid (mg/100g) content of minimally processed vegetables stored in different packaging material.	35
4.6.2	Effect of cutting knife and sodium hypochlorite-calcium chloride dip solution in ascorbic acid (mg/100g) content of minimally processed vegetables stored in different packaging material	35
4.7.1	Effect of cutting knife and sodium hypochlorite dip solution in water activity of minimally processed vegetables stored in different packaging material.	41
4.7.2	Effect of cutting knife and sodium hypochlorite-calcium chloride dip solution in water activity of minimally processed vegetables stored in	41
4.8.1	different packaging material. Effect of sodium hypochlorite dip solution in microbial load (cfu/ml) of	45
4.8.2	minimally processed beans stored in different packaging material Effect of sodium hypochlorite-calcium chloride dip solution in microbial load (cfu/ml) of minimally processed beans stored in different packaging material	45
4.8.3	Effect of sodium hypochlorite dip solution in microbial load (cfu/ml) of minimally processed capsicum stored in different packaging material	46

- 4.8.4 Effect of sodium hypochlorite-calcium chloride dip solution in microbial 46 load (cfu/ml) of minimally processed capsicum stored in different packaging material
- 4.8.5 Effect of sodium hypochlorite dip solution in microbial load (cfu/ml) of 47 minimally processed knol-khol stored in different packaging material
- 4.8.6 Effect of sodium hypochlorite-calcium chloride dip solution in microbial 47 load (cfu/ml) of minimally processed knol-khol stored in different packaging material

LIST OF FIGURES

Fig. no	Figure	Pg. no
4.1.1	Colour and texture of capsicum dipped in sodium hypochlorite- calcium chloride solution and packed in HDPP on fourth day of storage	22
4.1.2	Colour and texture deterioration on tenth day of storage of beans and capsicum dipped in sodium hypochlorite and packed in LDPP.	23
4.2.1	Effect of sodium hypochlorite dip treatment in moisture content of minimally processed vegetables cut with iron knife and stored in HDPP.	26
4.2.2	Effect of sodium hypochlorite dip treatment in moisture content of minimally processed vegetables cut with iron knife and stored in LDPP	26
4.2.3	Effect of sodium hypochlorite dip treatment in moisture content of minimally processed vegetables cut with stainless steel knife and stored in HDPP.	27
4.2.4	Effect of sodium hypochlorite dip treatment in moisture content of minimally processed vegetables cut with stainless steel knife and stored in LDPP.	27
4.2.5	Effect of sodium hypochlorite-calcium chloride dip treatment in moisture content of minimally processed vegetables cut with iron knife and stored in HDPP	28
4.2.6	Effect of sodium hypochlorite-calcium chloride dip treatment in moisture content of minimally processed vegetables cut with iron knife and stored in LDPP	28
4.2.7	Effect of sodium hypochlorite-calcium chloride dip treatment in moisture content of minimally processed vegetables cut with stainless steel knife and stored in HDPP	29
4.2.8	Effect of sodium hypochlorite-calcium chloride dip treatment in moisture content of minimally processed vegetables cut with stainless steel knife and stored in LDPP	29
4.6.1	Effect of sodium hypochlorite dip treatment in ascorbic acid concentration of minimally processed vegetables cut with iron knife and stored in HDPP.	36
4.6.2	Effect of sodium hypochlorite dip treatment in ascorbic acid concentration of minimally processed vegetables cut with stainless steel knife and stored in HDPP.	36
4.6.3	Effect of sodium hypochlorite dip treatment in ascorbic acid concentration of minimally processed vegetables cut with iron knife and stored in LDPP	37
4.6.4	Effect of sodium hypochlorite dip treatment in ascorbic acid concentration of minimally processed vegetables cut with stainless steel knife and stored in LDPP	37

÷

,

	ascorbic acid concentration of minimally processed vegetables cut with iron knife and stored in HDPP	
4.6.6	Effect of sodium hypochlorite-calcium chloride dip treatment in ascorbic acid concentration of minimally processed vegetables cut with iron knife and stored in LDPP	3
4.6.7	Effect of sodium hypochlorite-calcium chloride dip treatment in ascorbic acid concentration of minimally processed vegetables cut with stainless steel knife and stored in HDPP	3
4.6.8	Effect of sodium hypochlorite-calcium chloride dip treatment in ascorbic acid concentration of minimally processed vegetables cut with stainless steel knife and stored in LDPP.	4
4.7.1	Effect of sodium hypochlorite solution in water activity of minimally processed vegetables stored in HDPP.	4
4.7.2	Effect of sodium hypochlorite solution in water activity of minimally processed vegetables stored in LDPP.	4
4.7.3	Effect of sodium hypochlorite-calcium chloride solution in water activity of minimally processed vegetables stored in HDPP.	4
4.7.4	Effect of sodium hypochlorite-calcium chloride solution in water activity of minimally processed vegetables stored in LDPP.	4
4.8.1	Effect of sodium hypochlorite dip solution in microbial load (cfu/ml) of minimally processed beans stored in HDPP	4
4.8.2	Effect of sodium hypochlorite dip solution in microbial load (cfu/ml) of minimally processed beans stored in LDPP	4
4.8.3	Effect of sodium hypochlorite-calcium chloride dip solution in microbial load (cfu/ml) of minimally processed beans stored in HDPP	4
4.8.4	Effect of sodium hypochlorite dip solution in microbial load (cfu/ml) of minimally processed capsicum stored in HDPP	4
4.8.5	Effect of sodium hypochlorite-calcium chloride dip solution in microbial load (cfu/ml) of minimally processed capsicum stored in HDPP	5
4.8.6	Effect of sodium hypochlorite dip solution in microbial load (cfu/ml) of minimally processed knol-khol stored in HDPP	5
4.8.7	Effect of sodium hypochlorite-calcium chloride dip solution in microbial load (cfu/ml) of minimally processed knol-khol stored in HDPP	5

.