

CONTENTS

LIST OF FIGURES

LIST OF TABLES

1. INTRODUCTION	1-2
2. REVIEW OF LITERATURE	3-15
2.1 Anthocyanins	3-8
2.1.1 Classification of Anthocyanins	3-4
2.1.2 Biosynthesis of Anthocyanins	4-5
2.1.3 Antioxidant properties and health effects of anthocyanins	6-7
2.1.4 Anthocyanins in rice	7-8
2.2 Extraction	8-14
2.2.1 Solid Liquid Extraction	9
2.2.2 Extraction methods	10-11
2.2.3 Separation and purification of Anthocyanins	11-12
2.2.4 Characterization and detection of anthocyanins	12-14
2.2.4.1.1 HPLC	12
2.2.4.1.2 UV-Vis spectroscopy	12-13
2.2.4.1.3 Mass spectroscopy	13
2.2.4.1.4 Nuclear magnetic resonance	13
2.2.4.1.5 Hydrolysis technique	13-14
2.3 Antioxidant	14-15
2.3.1 Antioxidant activity	14
2.3.2 Anti-oxidative analysis methods	14-15
2.3.2.1 DPPH method	15
2.3.2.2 FRAP method	15
3. MATERIALS AND METHODS	16-29
3.1 Pigmented rice sample	16
3.2 Chemicals and reagents	16-17
3.3 Instruments used in the experiments	17

3.4 Preparation of anthocyanin extracts from coloured rice bran	18
3.5 Preparation of fruit juices	19
3.6 HPLC analysis of anthocyanins	20
3.7 Antioxidant assay	20-22
3.7.1 DPPH assay of the AECB extracts	20-21
3.7.2 FRAP assay of the AECB extracts	21-22
3.8 Total monomeric anthocyanin content of AECB extracts	22-23
3.9 Degradation studies of anthocyanins	23-25
3.9.1 Total monomeric anthocyanin content	23-24
3.9.2 Kinetics model of anthocyanins in aqueous solutions	24-25
3.10 Extraction of protein from local rice bran	25
3.11 Protein estimation by Lowry's method	25-26
3.12 Spray drying for encapsulated powders	26-28
3.12.1 Moisture content of encapsulated powders	26-27
3.12.2 Hygroscopicity of encapsulated powders	27
3.12.3 Solubility of encapsulated powder materials	27
3.12.4 Encapsulation efficiency of encapsulated powders	28
3.13 Colour measurement of encapsulated powders	28
3.14 Lemon juice fortified with encapsulated powders	28-29
3.14.1 pH of the lemon juices	29
3.14.2 Total anthocyanin content of the lemon juices	29
3.14.3 TSS of the lemon juices	29
3.15 Colour measurement of the lemon juices	29
3.16 Sensory evaluation of the lemon juices	29
4. RESULTS AND DISCUSSION	30-45
4.1 Obtaining rice bran from <i>Chakhao Poiraiton</i>	30
4.2 Extraction of anthocyanins from rice bran	30
4.3 HPLC analysis of the anthocyanin extracted from coloured rice bran AECB extracts.	30-31
4.4 Antioxidant assays	31-32
4.5 Total monomeric anthocyanin content (MAC)	32-33

4.6 MAC (mg/L) content of AECB extracts at pH buffers (1-5) at temperature 60, 70 and 80°C during time interval of 0-120 minutes.	33-34
4.7 Degradation kinetics studies of anthocyanin during heat treatment and at different pH buffer solutions	35-39
4.8 Protein extracted from local rice bran.	39
4.9 Yield of spray dried anthocyanin powders	39-40
4.10 Characteristics of the spray dried encapsulated powder materials	41-42
4.11 Colour measurement of encapsulated powder materials	42
4.12 Various parameters of the lemon juices incorporated with microencapsulates	42-43
4.13 Colour measurement of the lemon juices	44
4.14 Sensory Evaluation of the lemon juices	44-45
5. CONCLUSION	46
6. BIBLIOGRAPHY	47 -55

LIST OF FIGURES

Figure 2.1 The structure of the flavylum ion or 2-phenylbenzopyrilium	3
Figure 2.2 Schematic representation of the flavonoid biosynthetic pathway	5
Figure 2.3 Schematic extraction: before extraction and after extraction	9
Figure 3.1 Rice sample (<i>Chakhao Poiraiton</i>)	16
Figure 3.2 Preparation of fruit juice	19
Figure 4.1 Typical chromatogram of standard anthocyanin pigments and anthocyanin from pigmented rice	31
Figure 4.2 L-ascorbic acid standard curve for FRAP assay	32
Figure 4.3 Degradation of AECEB extracts at pH 1.0 during heating at 60, 70 and 80°C	35
Figure 4.4 Degradation of AECEB extracts at pH 2.0 during heating at 60, 70 and 80°C	35
Figure 4.5 Degradation of AECEB extracts at pH 3.0 during heating at 60, 70 and 80°C	36
Figure 4.6 Degradation of AECEB extracts at pH 4.0 during heating at 60, 70 and 80°C	36
Figure 4.7 Degradation of AECEB extracts at pH 5.0 during heating at 60, 70 and 80°C	37
Figure 4.8 Encapsulated powder (EP1) containing maltodextrine as carrier agent and anthocyanin extract	40
Figure 4.9 Encapsulated powder (EP2) containing maltodextrine as carrier agent, anthocyanin extract and bran proteins	40
Figure 4.10 S1: control lime juice; S2: lime juice containing 1% encapsulated powder EP1; S3: lime juice containing 1% encapsulated powder EP2	43
Figure 4.11 Radar graph representation of the sensory evaluation data	45

LIST OF TABLES

Table 2.1 Common anthocyanidins present in nature	4
Table 4.1 Rice bran obtained from <i>Chakhao Poireiton</i>	30
Table 4.2 DPPH activity and FRAP reducing power of AECB extracts	32
Table 4.3 Initial monomeric anthocyanin content (MAC) of AECB extracts at different pH buffers	33
Table 4.4 Initial monomeric anthocyanin content (MAC) of AECB extracts at different pH buffers after 30 min of heat treatment at 60, 70 and 80°C	34
Table 4.5 Effect of temperature and pH on the k , $t_{1/2}$, E_a and Q_{10} values of anthocyanin degradation during heating at 60, 70 and 80°C	38
Table 4.6 Yield of spray dried encapsulated powders	39
Table 4.7 Various parameters of encapsulated powder materials	41
Table 4.8 Hunter colour Lab results of encapsulated powder materials	42
Table 4.9 pH, TSS and TAC of encapsulated powder materials.	43
Table 4.10 Hunter Colour Lab results of the lemon juices.	44
Table 4.11 Sensory scores of the lemon juices with microencapsulates	44