List of abbreviations

| O ⁰ | Degree Celsius |
|--|--|
| AFLP | Amplified fragment length polymorphism |
| ANOVA | Analysis of Variance |
| ATP | Adenosine Triose Phosphate |
| BLAST | Basic Local Alignment Search Tool |
| C/N ratio | Carbon/ Nitrogen ratio |
| CCD | Central Composite Design |
| CVD | Cardiovascular Disease |
| DNA | Deoxy Ribose Nucleic Acid |
| dNTP | Deoxyribo Nucleotide Triose phosphate |
| | |
| ECLT | Euglobulin Clot Lysis Time |
| ECLT EDTA | Euglobulin Clot Lysis Time Ethylene Diammine Tetraacetatetae |
| | - |
| EDTA | Ethylene Diammine Tetraacetatetae Fibrinolytic activity/ Caseinolytic activity |
| EDTA F/C ratio | Ethylene Diammine Tetraacetatetae Fibrinolytic activity/ Caseinolytic activity ratio |
| EDTA F/C ratio FC reagent | Ethylene Diammine Tetraacetatetae Fibrinolytic activity/ Caseinolytic activity ratio Folin-Ciocalteu's Reagent |
| EDTA F/C ratio FC reagent FFD | Ethylene Diammine Tetraacetatetae Fibrinolytic activity/ Caseinolytic activity ratio Folin-Ciocalteu's Reagent Fractional Factorial Design |

| kDa | Kilo Dalton |
|-----------|---|
| М | Molar |
| MEGA | Molecular Evolutionary Genetic Algorithm |
| MI | Mili Liter |
| mM | Mili Molar |
| MR | Methyl Red |
| NB | Nutient Broth |
| NCBI | National Center for Biotechnological Information |
| NE, India | North-East, India |
| NK | Natto kinase |
| O.D | Optical Density |
| PCR | Polymerase Chain Reaction |
| PEG | Poly Ethylene Glycol |
| RAPD | Rapid Amplification of Polymorphic DNA |
| RFLP | Restriction Fragment Length Polymorphism |
| RNA | Ribose Nucleic Acid |
| RPM | Rotation Per Minute |
| rRNA | Ribosomal Ribose Nucleic Acid |
| RSM | Response Surface Methodology |
| S.D | Standard Deviation |
| SDS | Sodium Dodecyl Sulphate |

| SIM | Sulfur Indole Motility |
|------|------------------------------|
| SMA | Skimmed Milk Agar |
| SmF | Submerged Fermentation |
| TAE | Tris Acetate EDTA |
| ТСА | Tricholoacetic acid |
| t-PA | Tissue plasminogen activator |
| TSI | Triple Sugar Iron |
| U | Unit (Enzyme activity unit) |
| VP | Voges proskaure |
| w/v | Weight/ Volume |
| w/w | Weight/ Weight |
| WHO | World Health Organization |
| hð | micro- gram |
| μί | micro-liter |

.

Contents

| Legends | | Page |
|---------|--|-------|
| | | No |
| | Chapter 1: Introduction | 1-17 |
| 1.0 | Introduction | 2-6 |
| 1.2 | Information on Cardiovascular diseases, available using | 7-8 |
| | thrombolytic drugs and its limitations | |
| 1.3 | Microbial protease (fibrinolytic enzymes) a boon for health sector | 8-11 |
| 1.4 | Classification of protease (fibrinolytic enzymes) | 11-14 |
| 1.4.1 | Type of serine proteases | 11-12 |
| 1.4.2 | Chymotrypsin-like proteases | 12-13 |
| 1.4.3 | Subtilisin-like protease or Subtilases | 13 |
| 1.4.4 | Wheat serine carboxypeptidase II-like protease | 13 |
| 1.4.5 | Prolyloligopeptidae-like serine protease | 13 |
| 1.4.6 | Myxobacter ∝-lytic proteases | 13 |
| 1.4.7 | Staphylococcal protease | 13-14 |
| 1.5 | Protease (fibrinolytic) enzyme fermentation and yield | 14 |
| | improvement | |
| 1.5.1 | Fermentation methods | 14-17 |
| 1.5.2 | Aim and objectives | 16-17 |
| | Chapter 2: Review of literature | 18-25 |
| 2.1 | Non-food sources | 19-20 |

| 2.2 | Food sources | 20-22 |
|-----------|---|-------|
| 2.3 | Comprehensive review on fibrinolytic enzyme production | 23-24 |
| | by submerged fermentation | |
| 2.4 | Application and perspectives | 25 |
| | Chapter 3: Materials and Methods | 26-43 |
| 3.1 | Materials | 27 |
| 3.1.1 | Plastic ware / Glass ware /Columns | 27 |
| 3.1.2 | Chemicals | 27-28 |
| 3.1.2.1 | Analytical grade | 27 |
| 3.1.2.2 | Microbiological grade culture media/chemicals | 27 |
| 3.1.2.3 | Molecular biology grade chemicals /kits | 28 |
| 3.1.2.4 | The bacterial strain | 28 |
| 3.2 | Methods | 28 |
| 3.2.1 | Pure culture preparation of fibrinolytic protease secreting | 28 |
| | bacterial isolates | |
| 3.2.1.1 | Spread plate technique | 28-29 |
| 3.2.1.2 | Streak-plate technique | 29 |
| 3.2.2.3 | Routine maintenance and preservation of Microorganism | 29 |
| 3.2.3 | Taxonomic identification of alkaline protease (Fibrinolytic | 29-36 |
| | enzyme) producing bacteria | |
| 3.2.3.1 | Morphological test | 29-30 |
| 3.2.3.1.1 | Gram staining | 29-30 |

| 3.2.3.2 | Biochemical test | 30-33 |
|------------|--|-------|
| 3.2.3.2.1 | Hydrolysis test for casein, starch, lipid and gelatin | 30 |
| 3.2.3.2.2 | Carbohydrate fermentation test | 30. |
| 3.2.3.2.3 | Triple sugar iron (TSI) agar test | 31 |
| 3.2.3.2.4 | IMVIC test | 31 |
| 3.2.3.2.5 | Hydrogen sulphide test | 32 |
| 3.2.3.2.6 | Urease test | 32 |
| 3.2.3.2.7 | Litmus –milk test | 32 |
| 3.2.3.2.8 | Nitrate reduction test | 32 |
| 3.2.3.2.9 | Catalase test | 32-33 |
| 3.2.3.2.10 | Oxidase test | 33 |
| 3.2.3.3 | Ribotyping using 16S rRNA gene amplification | 33-36 |
| 3.2.3.3.1 | Isolation of DNA | 33-34 |
| 3.2.3.3.2 | PCR amplification of 16S rRNA gene | 34-35 |
| 3.2.3.3.3 | PCR amplification of the 16S-23S inter spacer region (ISR) | 35 |
| | region of the isolated DNA | |
| 3.2.3.4 | Phylogenetic analysis | 36 |
| 3.2.4 | Determination of protease activity | 36-37 |
| 3.2.5 | Optimization of culture condition for optimum growth and | 37 |
| | Maximum fibrinolytic protease production by by selected bacteria | |
| | under SmF systems | |
| 3.2.5.1 | Fibrinolytic protease production under submerged Fermentation | 37 |

(SmF) systems

| 3.2.5.2 | Effect of various carbon sources on fibrinolytic protease | 38 | |
|-----------|--|-------|--|
| | production | | |
| 3.2.5.3 | Effect of various inorganic and organic nitrogen sources on | 38 | |
| | fibrinolytic protease production | | |
| 3.2.5.4 | Effect of pH on fibrinolytic protease production | 38 | |
| 3.2.5.5 | Effect of incubation time on fibrinolytic enzyme Production | 38 | |
| 3.2.6 | Fibrinolytic protease production under submerged fermentation | 38 | |
| | system | | |
| 3.2.6.1 | Statistical optimization of protease production in SmF | 38-43 | |
| 3.2.6.1.1 | Screening of factors effecting fibrinolytic protease Production | 39-40 | |
| | using Plackett-Burman design | | |
| 3.2.6.1.2 | Statistical optimization of fibrinolytic protease production using | 41-42 | |
| | Response Surface Methodology(RSM) | | |
| 3.2.7 | Validation of response surface; Batch fermentation Under | 43 | |
| | optimized condition | | |
| | Chapter 4: Results | 44-83 | |
| Results | | | |
| 4.1 | Pure culture preparation of the bacterial strain: FF02B | 45 | |
| 4.2 | Taxonomic identification of fibrinolytic enzyme producing FF02B | 46-58 | |
| | bacterial isolate | | |
| 4.2.1 | Phenotypic study of FF02B bacterial isolate | 46 | |

| 4.2.1.1 | Morphological identification: Gram staining of The bacterial strain | 46 |
|---------|---|---------|
| 4.2.1.2 | Biochemical identification | 47-50 · |
| 4.3.2 | Genotyping profiling of FF02B bacterial isolate | 51 |
| 4.3.2.1 | PCR amplification of 16S rRNA gene of FF02B Bacterial isolate | 51-52 |
| 4.3.2.2 | Phylogenetic tree construction using 16S-rRNA gene | 53-54 |
| 4.3.2.3 | PCR amplification of 16S-23S rRNA gene of FF02B bacterial | 55-56 |
| | isolate | |
| 4.3.2.4 | Phylogenetic tree construction using 16S-23S rRNA gene | 57 |
| 4.3.2.5 | Bacterial naming and identification | 58 |
| 4.4 | Screening of initial process parameters for fibrinolytic enzyme | 58-73 |
| | production from FF02B bacterial isolate under submerged | |
| | fermentation | |
| 4.5 | Statistical optimization of influencing parameters for Plackett- | 74 |
| | Burman design | |
| 4.5.1 | Screening of influencing parameters by Plackett-Burman design | 74-76 |
| 4.5.2 | Statistical optimization of fibrinolytic enzyme production using | 76-82 |
| | Central composite design (CCD) | |
| 4.5.3 | Validation of the model | 82-83 |
| | Chapter 5: Discussion | 84-90 |
| 5.1 | Isolation and culture maintenance techniques | 85 |
| 5.1.1 | Pure culture techniques | 85 |
| 5.1.2 | Streak plate method | 85-86 |
| | | |

V

| Reference and Appendix | | | |
|------------------------|--|-------|--|
| | Chapter 6: Conclusion | 91-93 | |
| 5.3.1.2 | Response surface method | 90 | |
| 5.3.1.1 | Plackett-Burman Design | 89 | |
| | fibrinolytic protease production using statistical tools | | |
| 5.3.1 | Screening of influencing process parameters for microbial | 89-90 | |
| | fermentation system (SmF) | | |
| 5.3 | Microbial fibrinolytic protease production under submerged | 88-89 | |
| 5.2.2 | Genotypic approach | 87-88 | |
| 5.2.1 | Phenotypic approach | 87 · | |
| 5.2 | Bacterial identification | 86 | |
| 5.1.3 | Pour plate Method | 86 | |

List of Tables

| Table No. | Table Legends | Pages |
|-----------|--|-------|
| | Chapter 1: Introduction | 1-17 |
| 1.1 | Enzymes in various industrial segments and their | 4-6 |
| | applications | |
| 1.2 | Sources of microbial fibrinolytic enzymes | 9 |
| 1.3 | Fibrinolytic enzyme producing Bacilli, isolated from traditional | 10 |
| | fermented food | |
| 1.4 | Statistical method to improve protease production from | 16 |
| | microorganisms | |
| | Chapter 2: Review of Literature | 18-25 |
| 2.1 | Recent reports on isolation of fibrinolytic enzymes from | 21 |
| | various sources from India | |
| 2.2 | Some latest and selected reports on isolation of fibrinolytic | 22 |
| | enzymes from various sources from abroad | |
| 2.3 | Recent reports on statistically optimized process parameters | 24 |
| | for fibrinolytic enzyme prouction under submerged | |
| | fermentation | |
| | | 00 40 |

26-43

Chapter 3: Materials and Methods

- 3.1 Optimal PCR reaction conditions or amplification of 35 conserved region of 16S-rRNA gene of selected fibrinolytic protease secreting bacterial strain
- 3.2 Independent variables for fibrinolytic protease production 40
 under SmF systems using Plackett-Burman design.
- 3.3
 Independent variables for fibrinolytic protease production
 42

 under SmF system using Central Composite Design
- 3.4 Validation of response surface for fibrinolytic protease 43 production

- 4.1 Biochemical and morphological tests of bacteria FF02B 50 isolate. Experiments were repeated thrice to assure reproducibility
- 4.2 Homologous search results of 16S-rRNA gene sequence 52 . using BLAST tool from NCBI.
- 4.3 Homologous search results of 16S-23S ISR gene sequence 56 using BLAST tool from NCBI
- 4.4 Representation of bacterial identification and designated 58 names

| 4.5 | Plackett-Burman store design showing three variables with | 75 |
|-----|---|----|
| | coded values along with the observed results for fibrinolytic | |
| | enzyme production by FF-02B bacterial isolate | |

- 4.6 Statistical analysis of Plackett–Burman design showing 76 coefficient values, t- and p-value for each variable for fibrinolytic enzymes from FF-02B bacterial isolate (p-value <0.05).</p>
- 4.7 Observed responses and predicted values of fibrinolytic 77 enzyme production by FF02B bacterial isolates.
- 4.8 Analysis of Variance (ANOVA) of fibrinolytic enzyme 78 produced by FF02B bacterial isolate
- 4.9 Model coefficients estimated by multiple linear regressions 79 (significance of regression coefficients) for fibrinolytic enzyme production by FF02B bacterial isolate in SmF under shake-flask study (p<0.05).</p>

|) |
|---|
| |

Chapter 6: Conclusion 91-93

| List of Figure legends | | | | |
|-------------------------|---|------|--|--|
| Figure | Figure Legends | Page | | |
| No. | | No. | | |
| Chapter 1: Introduction | | | | |
| 1.1 | The steps involved in classical versus state-of the- art | 3 | | |
| | development of enzymes | | | |
| 1.2 | The proportion of deaths by causes in WHO regions, estimates for | 7 | | |
| | 2000 (WHO, 2001) | | | |
| | Chapter 2: Review of Literature | | | |
| | Chapter 3: Materials and Methods | | | |
| | Chapter 4: Results | | | |
| 4.1 | Preparation of pure culture | 45 | | |
| 4.2 | Gram staining of FF02B cells at a magnification of 1000x under | 46 | | |
| | compound microscope | | | |
| 4.3 | Biochemical fingerprints of FF02B bacterial isolate | 47 | | |
| 4.4 | Biochemical profile of FF02B bacterial isolate | 48 | | |
| 4.5 | Biochemical property of FF02B bacterial isolate | 49 | | |
| 4.6 | PCR amplification of 16S rRNA gene from FF02B bacterial isolates | 51 | | |
| 4.7 | Phylogenetic tree construction of 16S-rRNA gene using Neighbor- | 53 | | |
| | Joining method | | | |
| 4.8 | Phylogenetic tree construction using minimum evolutionary | 54 | | |
| | method | | | |
| 4.9 | PCR amplification of 16s-23s ISR region amplification | 55 | | |
| 4.10 | Phylogenetic tree construction off 16S-23S ISR gene using | 57 | | |
| | Neighbor-Joining method | | | |
| 4.11 | Effect of pH on bacterial growth (\blacksquare) and dry biomass (\blacktriangle) post | 59 | | |
| | incubated for 24h at 37°C. Values are means ± S.D are of | | | |
| | triplicate experiments. | | | |
| | | | | |

| 4.12 | Effect of pH on protein content (♦), Caseinolytic activity (■),and fibrinolytic activity (△) of culture supernatant from FF02B bacterial isolate. Value are means ± S.D are of triplicate experiments. | 60 |
|------|--|----|
| 4.13 | Effect of pH on F/C ratio of culture supernatant FF-02B bacterial isolates. Value are means ± S.D are of triplicate experiments | 61 |
| 4.14 | Effect of carbon sources on bacterial growth (☑) and bacterial dry biomass (◆) from FF02B bacterial isolate. Values are mean ± S.D are of triplicate value. | 63 |
| 4.15 | Effect of carbon sources on protein content (\blacklozenge), Caseinolytic activity (\blacksquare) and fibrinolytic activity ($▲$) production from FF02B bacterial isolate. Values are mean \pm SD are of triplicate value. | 64 |
| 4.16 | Effect on Carbon source on F/C ratio on Fibrinolytic enzyme production by FF02B bacterial isolates. Values are mean \pm S.D of triplicate values. | 65 |
| 4.17 | Effect of nitrogen sources on bacterial growth (▲) and bacterial dry biomass (■) of FF02B bacterial isolates. Values are mean ± S.D are of triplicate experiments. | 67 |
| 4.18 | Effect of nitrogen sources on protein yield (\blacklozenge), Caseinolytic activity (\blacksquare) and fibrinolytic activity (\blacktriangle) of FF02B bacterial isolates. Values are mean ± S.D are of triplicate experiments | 68 |
| 4.19 | Effect of nitrogen sources on F/C ratio of fibrinolytic enzymes from FF02B bacterial isolates. Values are mean \pm S.D are of triplicate experiments | 69 |
| 4.20 | Effect of incubation time on bacterial growth (■)and bacterial dry biomass (▲) of FF02B bacterial isolates. Values are mean ± S.D are of triplicate experiments | 71 |
| 4.21 | Effect of incubation time on protein content (\blacktriangle), Caseinolytic activity (\blacksquare), and fibrinolytic activity (\blacklozenge) of FF2B bacterial isolate. Values are mean ± S.D are of triplicate experiments. | 72 |
| 4.22 | Effect of incubation time on F/C ratio of protease enzymes | 73 |

XI

| | produced by FF-02B bacterial isolates. Values are mean \pm S.D | |
|----------|---|-------|
| | are of triplicate experiments. | |
| 4.23 (a) | Response Surface Methodology plots for F/C ratio for fibrinolytic | 80 |
| | enzymes from FF02B bacterial isolates response surface plot. | |
| 4.23 (b) | RSM plots for F/C ratio for fibrinolytic enzymes from | 81 |
| | FF02B bacterial isolates | |
| | Chapter 5: Discussion | 84-90 |
| | Chapter 6: Conclusion | 91-93 |
| | Reference and Appendix | |

,